



Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

A

MANUAL OF DIETETICS

BY

J. MILNER FOTHERGILL, M.D., EDIN.,

LECTURER ON THE DISEASES OF THE DIGESTIVE ORGANS, IN THE ROYAL COLLEGE OF PHYSICIANS, EDINBURGH;
FELLOW OF THE MEDICAL SOCIETY OF EDINBURGH;
FELLOW OF THE ROYAL SOCIETY OF EDINBURGH;
FELLOW OF THE ROYAL SOCIETY OF MEDICAL PHYSICIANS, PHILADELPHIA.

"WHAT IS ONE MAN'S FORTH IN ANOTHER MAN'S?"

NEW YORK
WILLIAM WOOD & COMPANY

1886

CONTENTS.

PART I.

PAGE

CHAP. I.—Food: Its Object,	1
II.—Forms of Food and their Digestion,	6
III.—Methods of Preparing Food,	23
IV.—Condiments,	39
V.—Beverages,	42
VI.—Stimulants,	49
VII.—Fluid Foods,	64
VIII.—Preserved and Canned Foods,	72
IX.—Prepared Foods,	79
X.—Artificial Digestive Agents,	85

PART II.

XI.—Food in Infancy,	96
XII.—Food in Adolescence,	109
XIII.—Food in Adult Life,	118
XIV.—Food in Old Age,	130
XV.—Food in Acute Disease,	137
XVI.—Food in Convalescence,	147
XVII.—Food in Gastric Affections,	157
XVIII.—Food in Struma,	167
XIX.—Food in Anæmia,	169
XX.—Food in Constipation and Diarrhoea,	167
XXI.—Food in Phthisis,	177
XXII.—Food in Chronic Heart and Lung Disease,	184

	PAGE
CHAP. XXIII. Food in Bright's Disease,	189
XXIV. Food in Albuminuria,	195
XXV. Food in Diabetes,	197
XXVI. Food in Glycosuria,	203
XXVII. Food in Gout,	205
XXVIII. Food in Neurosal Affections,	216
XXIX. Food for Chronic Invalids,	222
XXX. Food in Obesity,	225
XXXI. Food in Indigestion,	230
XXXII. Food in Biliousness,	242
XXXIII. Food given otherwise than by the Mouth,	250
Conclusion,	254

To
FORDYCE BARKER, M.D., LL.D., EDIN.,
AN ACKNOWLEDGED LEADER IN MEDICINE,
THIS WORK
IS AFFECTIONATELY DEDICATED
BY
THE AUTHOR.

PREFACE.

The day of Dietetics has arrived.

Modern advances in our knowledge of the physiology of digestion have been accompanied by a like progress in the preparation of foods. The value of predigested carbohydrates in acute disease and malassimilation among adults, as well as children, is now being gradually realised. The digested albuminoids are making their way ; while oil emulsions have established themselves on a firm footing. The many evils of a dietary too rich in albuminoids are now being generally recognised. The time indeed is at hand when systematic lectures on Food will be a part of Medical education ; while the value of feeding in disease is admitted to be as important as the administration of medicines.

3, HENRIETTA STREET,
CAVENDISH SQUARE,
LONDON, W.

MANUAL OF DIETETICS.

CHAPTER I.

THE OBJECT OF FOOD.

BEFORE the days of chemistry, food was in its pre-scientific stage. Then it was mainly a question of the palate. Kings kept their cooks, and smaller personages imitated them. The cooks cooked by day and devised new dishes by night. In the present work, however, this culinary aspect of food will be subordinated to that of its scientific aspect.

Not that the cook is to be ignored, by any means. Without proper cooking, no food is palatable; and if the palate—the janitor of the stomach—be offended, no matter what is its real value, the food is useless. The palate must be consulted as well as the stomach.

Of old the palate ran riot. The Romans were both epicures and gluttons. The oysters of Britain were conveyed by relays of runners to the Roman villas. One epicure cast a slave into his fishpond to improve the flavour of his lampreys. The most absurd dishes were contrived, up to a pie of nightingales' tongues. Their tastes were not ours. Fowls were boiled in aniseed water, and served with a sauce containing aniseed, mint, mustard seeds and asafoetida. Nor were the middle ages far different. Charles V. of Germany was a notorious gourmand. Besides ordinary meats he had roasted horse, cats in jelly, lizard soup, fried frogs, etc., &c. His cook, when asked for a new dish, could only suggest a compote of warts. In allusion to that monarch's passion for such time-keepers,

Animals were chased because it was found that the flesh of hunted beasts was more tender and palatable than that of those killed without such preliminary preparation. Fowls were driven to death to fit them for the table; while one who had been

"How to roast and eat a goose alive." No cruelty to an animal was too great to gratify the palate. It was merciless!

The amount eaten, too, was enormous. The Roman had five meals a day; and at great feasts when full to repletion, the stomach was emptied in order that the process of filling it might be recommenced.

Charles V. was quite as big a glutton as he was an epicure. At a dinner of the Knights of the Golden Fleece Roger Ascham saw him make his way through sod beef, roast mutton, and baked hare, after which he fed full well on a capon. Nor did he forget to drink with it all. "He had his head in the glass five times as long as any of them, and drank no less at once than a quart of Rhine wine." No wonder he became a gouty man.

Then views as to food varied in the widest manner. The English athlete placed his faith in rare beefsteaks and raw eggs; while the Hindu wrestlers trained on sweetmeats. The English farmer feeds his servants on meat in order that they may be equal to hard work; just as he adds some extra corn to the dietary of his horses when working hard. The hanna of Stamboul carries incredible weights on a dietary which the English athlete would regard as utterly insufficient to support life; and for backsheesh is grateful for a slice of mutton.

Cookery-books were written by men of learning. Even Dr. Johnson, of dictionary fame, did not think the subject beneath him. "I could write a better book of cookery than has ever yet been written," he boasted. "It should be a book on philosophical principles." So important did he deem the subject that he said contemptuously, "A woman may spin, but she cannot make a good book of cookery." Dr. Kebleton, the author of "The Cook's Oracle," had no doubt of cookery related to food and cookery, consisting of no less than three volumes. It cannot then be said that the subject has been regarded as trifling in all ages.

But it was not till recent days, and from the teaching of Liebig, that the subject was regarded from a scientific point of view; both as regards the nutritive value of food, and our knowledge of organic chemistry, and the importance of a more complete comprehension of the utility of the various elements of the organism. Before that day the subject was regarded as *Disculina Disculum!*

The British Government, for instance, is particular in order to give the natives of the Sandwich Islands a taste of the Sandwich Islanders' like their own, that they should eat the same food as soon as caught.

Europeans prefer a fresh egg; in Cochin China the people prefer them "high." The Israelites talked of a land of "corn and wine" and of "oil and honey." The Egyptians lived on lentiles; while the Arab exists upon dates.

The Hindoo and the Negro both delight in boiling rice and butter. Ghee is clarified butter; and the Hindoo in praising anything comestible, conches his highest praise in the phrase "as good as ghee!" On the other hand, the Gauchos of the South American pampas live solely on lean beef and water; and Sir Francis Stead has told us what power of endurance this diet gives. The Eskimo and the natives of Northern Asia eat huge quantities of fat, and drink oil. This diet is a perfect contrast with that of the desert Arab. On the burning desert little is required to maintain the body heat, and a handful of dates is sufficient for the body-needs. On the frozen Arctic wastes, fat, the strongest fuel-food, is required in considerable quantities to keep the body temperature at a point compatible with life. Cut off from farinaceous food, the Eskimo rejoices in the liver of the walrus, with its glycogen, or animal starch; but to give it a higher heating power, he eats it with slices of fat. Instinctively man has adopted the dietary suited to his needs; and the Anglo-Saxon who, on the plains of Bengal, persists in the roast beef and ale of Old England, pays the penalty of his dietary indiscretion in hepatic trouble. Blindly and darkly then, instinct has guided man in his choice of food, before the daylight of organic chemistry came to direct his path aright. Light has dispelled the darkness; but the palate is not to be dethroned. There are other things that cling to us as well as original sin! The belief in meat is ubiquitous with the Anglo-Saxon. No one can be strong without meat, it is asserted.

"Many people seem to look upon meat almost as though it formed the only food that really nourished and supplied what is wanted for work. The physician is constantly coming across an expression of this view. Undoubtedly, a greater feeling of satiety is produced by meat than by other food. It forms a greater stay to the stomach, but this arises from the stomach constituting the seat of its digestion; and a longer time being occupied before it passes on and leaves the organ in an empty condition." (Pavy on Food and Dietetics.)

This fallacy is disproved by the fact that the Sikhs of the Punjab are pulse-eaters. The Italian lives almost solely on rice and macaroni. The dreaded Iroquois were cultivators of maize, of bean, and hunting Indians. The Spaniard munches his onion and digests

crust of bread in oil. The Kaffir, like the Kirghis, lives mainly on milk; as did the Cynri, when Cæsar invaded Britain. The Arab has his dates. The Brahmin prefers the banana. The Hindoo, the Chinese and the Japanese, find their chief sustenance in rice, which with them takes the place of bread with us. Bread or its farinaceous equivalent is the staff of life. But "thou shalt not live by bread alone" tells that something in addition is necessary for the food of man.

What is that something? Milk; certainly! Meat? No, emphatically No! Fat, Yes!—especially butter. Some fluid is, of course, necessary as water. Here we have all that is required. There is starch, body-fuel; there is albumen for tissue-repair in the gluten; there are earthy salts, too, especially if the bread be made of whole meal; and then there is the fat, which partly is burnt as body-fuel, partly engaged in the building up of healthy tissue.

Let us trace the history of a mouthful of bread and butter. While it is being masticated, the saliva is brought into contact with many of the starch granules (already cracked by the heat of the oven and, not only that, but a step forward has thus been made in its solution,) and the conversion of insoluble starch into soluble sugar is begun. Then it is swallowed, when a new action is set up. The soluble grape sugar of the converted starch passes through the wall of the stomach into the gastric vesicles of the portal vein; leaving the undissolved residuum behind. The acid gastric juice next acts upon the gluten, the albuminous stroma, or framework of the mass—and by dissolving it, liberates the rest of the starch granules which had escaped the contact of the saliva. When the softened mass passes the pyloric ring, and comes into contact with the bile and the secretion of the pancreas, the most active part of the digestive net sets in. The liberated starch granules come into contact with the diastase of the pancreatic secretion, and are by it converted into soluble grape sugar; the digestion of the albuminous substance is further carried on by the trypsin; the fat is emulsionised. This is only digestion. The sugar, the dissolved albuminoids, the earthy salts, pass into the portal vein, to be dealt with by the liver; while the fat globules pass by the mouths of the lacteals into the lymphatics. Such is the digestion of a typical mass of food; there are two important variations (starch and sugar), albuminoids, fat, and earthy salts. And such the art and skill of all the cooks who ever lived, ancient, cool, and thorough the various edible substances of the animal and vegetable world, these are the constituents of the food of Man.

The carbo-hydrates are the body-fuel, the surplusage being stored as fat; the albuminoids serve to repair the tissues as they wear out; the salts form the blood-salts; the fat helps to build up normal health-tissues, the excess being burnt as body-fuel. That is the real object of food!

But Nature has veiled her object. She provides an imperious palate, engrafted upon an imperative sense of hunger, and these secure the nutrition of the body. Succulent vegetables, luscious fruits, sapid flesh; such are the materials upon which the cook exercises his art; and such are the food of the savage. While milk, the natural food of the new-born, alone or in combination with (chiefly) vegetable matter, is ever acceptable. Hunger compels the individual to take food to fill the stomach; the palate guides him in his choice; if the food be simple, there is little or no temptation to indulge in excess. But when the cook's art tempts the palate, then comes in the temptation. The first temptation came through the palate—"the tree was good for food;" and Adam hearkened into the voice of Eve, and ate of the tree—and the history of human troubles began. Now, the man has hearkened to the voice of the tempter, and eats—in excess of his body-needs. He must eat to live; but that is not the true equivalent of to live to eat. The object of food is the nutrition of the body—rather than the gratification of the palate. It may be well next to examine these different constituent elements of food in fuller detail both as to their digestion and their destiny.

CHAPTER II.

FORMS OF FOOD WITH THEIR DIGESTION.

Carbo-hydrates -Albuminoids -Fat-salts. Liebig's great division of food into hydro-carbons, or "fuel-food," and albuminoids, or "tissue-food," though at one time subjected to much criticism has, in the main, stood its ground. Modern advance in physiological chemistry certainly has told us that albuminoids are oxidised in their retrograde metamorphosis and so contribute to the body-heat; we know, too, that fat is necessary to the formation of healthy tissue—as seen in the cod liver oil so much in repute for the treatment of phthisis—and so to some extent is a tissue-food, as well as a fuel-food. Even certain salts, as the malates, the citrates and the tartrates are a fuel-food, in that the malic, citric and tartaric acids are oxidised, while the base (largely potash), appears in the urine as a carbonate or may be a lithate. The muscles require potash and the blood soda; why we do not yet quite know. The bones call for lime; and admitting that Liebig's view is not precisely accurate, it is true as a broad division; and is well worth bearing in mind. Why? Because since the days when Cain kept sheep, and gave way to an ungovernable temper and behaved like a carnivorous brute, in his numerous instincts, man has been tempted to take meat to gratify his palate, to an extent far beyond his real tissue-needs. Man is like a steam engine in that he is a piece of mechanism in which fuel is burnt. Certainly the iron framework of the engine needs a slow oxidation, and needs repair; and so do the tissues of man wear out. The engine needs oil to lubricate its bearings, and so do other matters which have to be continuously supplied to it. The man needs far other wants than merely "fuel-food" and "tissue-food," as motor-oil and motor-lubrication. The engine is a means of heating water into steam, of converting that steam into put wheels into motion, of converting heat into motor power. When the engine is working hard it takes in much more water and fuel. When a man is exerting himself he produces more heat by increased oxidation, and he must take in more food. Liebig doubtless was in error in assuming that the heat of the body is produced when fuel tissues are consumed.

No more than the framework of a locomotive when ascending an incline. An engine doing hard work will need more oil for its lubrication than one doing little work; and probably will sooner have to be sent to the repairing shed. A hardworking man may wear out his muscles at a more rapid rate than an idle man. But all the same in the locomotive and the man, the framework and the tissues do not "burn," but rust, or undergo a slower oxidation. (In forming conclusions from the amount of urea—the body ash—the perpetual and ineradicable difficulty has to be met that urea is the final soluble form of nitrogenised waste derived alike from the *luxus consumption*, or surplussage of albuminoid food and from effete or worn-out tissue.) Just as when the body is at rest its temperature is maintained by oxidation of hydro-carbons, so in activity the combustion is the same; only greater. Consequently the tissue-food for tissue-repair actually required is small, and more albuminoids are usually eaten than are absolutely needed. Granting that the general health of convicts deteriorates on an insufficient dietary; and that the system suffers when imperfectly fed; and that when work is hard more food, especially of the stimulating nitrogenised character is given, as the result of the teaching of experience; still the main fact remains unaffected, viz., that the actual wear and tear of the albuminous tissues is small, and can really be met by a dietary far less rich in azotised matter than that taken by most persons. Look at the lusty babies fed on "Pobbs," consisting of bread and water (with little if any milk), to be seen in any agricultural district in England. See the huge navy "Lincolnshire bred," reared on bread and bacon, or dripping and potatoes; or the Scotchman built up on oatmeal; and we can understand that the constructive powers of the human organism can successfully contend with great difficulties. But all the same, when the navy can procure some beefsteak he can do a harder day's work; and, I am informed, that a Highland gillie, when he shares the sportsman's repast on a dietary superior to that to which he is accustomed waxes in vigour and energy, in an obvious and unmistakable manner; while Dr. Ferguson of Bolton has told us of the improvement in the physique of the children of the Lancashire operatives when a certain amount of milk was added to their bread and butter and tea.

Keeping in mind then Liebig's great generalisation (which, notwithstanding that it is not absolutely correct as to minutæ) we find the food of man ranging itself into three main divisions:

1. Carbo-hydrates:

2. Albuminoids:

3. Fat:

With a small sub-class—

4. Salts.

Water too is essential to the body, but it cannot exactly be classed as a "Food."

Carbo-hydrates.— Under the term Carbo-hydrates are classed the starchy, or farinaceous, or amyloid (as we choose an English, a Latin, or a Greek term), and the saccharine elements of our food.

Starch consists of Carbon, Nitrogen and Oxygen (having the formula $C_6H_{10}O_5$) and is found in all the cereals: especially in those used as the food of man, wheat, oats, barley, maize and rye; it is also found in the pulse tribe, or legumes, as the pea, bean, lentil and the haricot bean, as also the dahl of India; also in rice, buckwheat, sago, tapioca and arrowroot, and the quino of Peru. It abounds in the potato, and other roots (which contain the germs of the new plant), in carrots, parsnips and turnips which first produce a root, from which again a seed-bearing stem is developed; and is part of the unripe fruit, which becomes sweet as the starch is changed or converted into sugar.

Starch is the food of the seedling plant. To take the case of a grain of any cereal, it contains starch, albuminoid matter, and earthy salts; while oats and maize also contain notable quantities of fat. When the plant germinates the starch is converted into sugar, (as the baby plant requires it) by the action of the diastase (an albuminous ferment), as the seedling grows. Insoluble starch is converted into soluble sugar in the grain; just as it is in the human mouth. The plant stores up food for its young; and that food man takes for his own use. Why does the store of plant food take the form of insoluble starch? Because if soluble it would be liable to be dissolved by water; and after rain the grain field would be barren, because the food elements were washed away in solution. Insoluble starch can be stored.

A moist germination is artificially set up by the process of "malt'ing"; which we occupy our attention further on.

When man ate a cooked grain, as he did at an early period (as the worm-eaters of grain-nivorous men still tell us when dug up, he gnawed it slowly under his teeth; and during this time the starch came in contact with the saliva, which by its diastase converted this

starch into soluble dextrine (with the starch formula) and grape sugar ($C_6H_{12}O_6 \cdot H_2O$). The action was exercised upon the outside of the starch granule; the remainder passing on through the stomach till it reached the diastase of the pancreatic secretion, when it became fully dissolved. (All starch granules however are not equally acted upon, and where much starch food is taken a certain quantity of them are to be found in the feces.)

When man came first to crush and then to grind and cook his food, a new departure took place. The toil of grinding his grain was done for him in a less laborious manner. Disintegration took place in the mill. He could eat faster. But if no other change had gone on, this might have been a very questionable advantage; for it would have had the result of filling his stomach with insoluble starch granules (a fertile source of indigestion). The effect of heat upon the starch granule is to crack it and thus to form a channel by which the saliva could reach the interior; an important matter. But another change goes on of which the knowledge is far less general. The starch is converted largely into soluble dextrine. (When a portion of well-baked flour is subjected to the iodine test, the conversion of the starch is readily seen.) Consequently the farina is speedily transformed into soluble matters by the saliva. The existing dextrine is carried forward to grape sugar; while the remaining starch is largely changed into dextrine. Such then is the effect of the act of cooking upon the starch granule. Grinding and cooking lessen the labor of the jaws and the salivary glands.*

Disintegration and solution constitute the digestive act. The conversion of starch into grape sugar is brought about by the salivary diastase (an albuminous body whose formula has not yet been determined), by a process of hydration; and grape sugar has the formula $C_6H_{12}O_6 \cdot H_2O$, *i.e.*, some water has been introduced into the molecule. The saliva is operative in the mouth and in the gullet, and also in the stomach until it becomes distinctly acid. The soluble grape-sugar passes through the wall of the stomach into the blood.

This removal of much of the starchy matter of the food leaves the stomach to act upon the other constituents. When food is hastily swallowed, our mastication is imperfect from haste, and

* Grain and bread-crusts developed the jaws and teeth. The former is now in vogue for infants, some fact which is generally overlooked.

stomach is hampered and embarrassed by quantities of unchanged starch, and its digestion is slow and painful.

When the stomach becomes distinctly acid, the conversion of starch is arrested. When the stomach has done its work and gastric digestion is complete the acidity seems pretty much exhausted, and the chyme (as the pulvaceous mass is termed) finds its way into the duodenum, where it becomes mixed with the bile and the pancreatic secretion. In this alkaline medium the pancreatic diastase comes into play, and the transformation of insoluble starch into soluble dextrine and grape-sugar re-commences. As the more or less disintegrated or denuded starch granules pass along the small intestine they become fully dissolved into grape-sugar which passes into the blood of the portal vein.

Such then is the digestion of starch. But before following up the history of grape sugar in the body, it will be well to consider the different forms of sugar. Sugar may be divided into crystalline or cane sugar ($C_{12}H_{22}O_{11}$) and glucose or grape sugar. The crystalline sugar is the object of the sugar-baker; and the faces of sugar boiling, treacle or molasses, are the non-crystallisable sugar. On the other hand non-crystallisable, or invert sugar, otherwise levulose, is the object of the manufacturer who supplies the brewer with substitutes for malt. Crystalline sugar is found in the sugar cane, the beet root, other roots, the maple and the sugar grass. Both are sweet to the taste; crystalline sugar being more intensely sweet, while the flavour of non-crystalline sugar is finer to most palates. As sugar is the food of the seedling plant, so sugar is found in all young shoots (notably in those of the briar and the grasses, as all rural children know).

"Although cane sugar requires no digestion to fit it for absorption it may be considered probably that it undergoes conversion into grape-sugar, certainly in part, if not wholly before leaving the alimentary canal. If cane sugar be introduced into one of the vessels of the general circulation, it passes off from the system without being utilised as a source of energy, still in the form of cane sugar in the urine."

"If, however, cane sugar be introduced into the alimentary canal before it is absorbed, say, for subsequent assimilation, sugar similarly passes off into the urine; but now in the form of grape sugar instead of cane sugar." (Payson.)

Whether cane or grape sugar—whether derived from starch or from sugar cane or beet—be food, it matters not—it passes into the portal vein, and then on to the liver. Here it is dehydrated, or

turned back into glycogen, or animal starch ($C_6H_{10}O_5$). In this insoluble form once more it can be stored for the body needs. Carbo-hydrates then furnish the glycogen or body-store of fuel in the liver. The liver stores up from each meal so much glycogen; and gives it off as required; otherwise life would be only one dreary meal. (Any disturbance in these glycogenic arrangements leads to the presence of sugar in the urine. These disturbances may extend from the mere presence of sugar in the water after a meal, when the dehydrating power of the liver cannot keep pace with the diastatic activity, up to the dread disease diabetes.)

The insoluble store of glycogen is given off as required in the form of soluble grape sugar; which is probably burnt up as lactic acid ($C_3H_5O_3$) in union with soda, as lactate of soda. (Muscles have their little stores of glycogen; the muscles of the embryo being especially rich in it.)

Such then is the history of the Carbo-hydrates, our great "fuel-food"; corresponding to the coal in the tender of the locomotive. The surplusage of grape sugar forms the fat, or adipose tissue of the body; a further reserve of "fuel-food." The store of glycogen in the liver is readily got at, and so we may liken it to the coal-scuttle by the fireside, while the fat deposit is the coal cellar. In some individuals it is spread over the body generally; while in others the omentum and parietes of the abdominal muscles are its chief seats.


In some animals, as the pig for instance, the fat is wide-spread and general, while in the camel the fat is stored in the hump, and in the codfish the liver is the seat of the storage of fat, as well as of glycogen.

This grape sugar is the natural body fuel, and where it is burnt up, the organism perishes of exhaustion; whether in starvation or acute disease. This matter will be further discussed when the dietetic treatment of exhaustion is given further on.

Albuminoids.—Albumen is a complex body notably containing nitrogen. This is the cardinal matter which fits it for the non-oxidisable framework of the body in which the combustible hydrocarbons are burnt. "Of all the elements of the animal body, nitrogen has the feeblest attraction for oxygen; and, what is still more remarkable, it deprives all combustible elements with which it combines, to a greater or less extent, of the power of combining with oxygen, that is, of undergoing combustion." (Liebig.)

By virtue of this nitrogen-factor then, the albuminous substances are prevented from the body-combustion extending to them. These then

framework of the locomotive they *rust* but do not *burn*. Their composition is as follows:



From this we can see that they differ from the hydro-carbons in the fact that they contain some sulphur as well as a distinct amount of nitrogen. The egg from which the living bird is developed contains albumen, some fat, and water. Probably the shell furnishes the lime. White of egg is typical albumen. Albumen is found in the muscles or flesh of animals; it is found in milk, as caseine, and in all seeds, notably the pulse tribe, whose seeds are rich in vegetable caseine. A certain proportion is found in cereals as the gluten of wheat, for instance.

It is as necessary for the seedling to build up its tissues as it is for a baby. The ferment of seeds which converts starch into grape sugar, is an albuminous body. Albumen in a fluid form as the white of egg, coagulates and forms a firm white solid on boiling. Yet caseine, as in cheese, softens under heat. Milk does not coagulate on boiling; but meat juice does. Consequently if an infusion of meat be raised to the boiling point the albumen is coagulated. Beef tea raised to the boil and strained loses (with its albumen), what food-value it possesses.

Flesh cooked before the rigor mortis sets in, is readily masticated; and the Abyssinian it is said cuts his steak from the living ox as he wants it. Once the rigor mortis has set in the flesh is hard and tough; consequently it is customary to keep it until this has passed away before cooking it. The effect of cooking upon muscle is to loosen the muscular bundles or fibrillæ from each other, so that they are readily torn asunder and crushed by the teeth. Insufficiently cooked meat is hard and resists the action of the teeth; over cooked meat is stringy. Perfectly cooked flesh is more savoury than either underdone or overdone meat. When taken into the mouth the muscular fibres are torn asunder and crushed by the act of mastication; so that when they enter the stomach the gastric juice can readily attack them. If meat be not chewed but bolted, the solvent juice can only act upon the exterior of the mass; while "lumps" offend the stomach and arrest the gastric secretion, as was found in the case of Alphonse St. Martin. Solution is at work under the influence of pepsine, aided by an acid. "The natural bundles of meat and vegetables fall to pieces; the muscular fibres split up into discs, and the protoplasm is dissolved from the vegetable cells." (M. Foster.)

The albuminous stroma of bread or liver is dissolved and the

starch granules set free; while the albuminous envelope of the fat cells (adipose tissue) also is dissolved and the fat is left free. The food is changed into a pulraceous mass, "such pieces as most resist the gastric juice being the last to leave the stomach." The digestive action of the stomach is exercised exclusively upon albuminoids; carbo-hydrates and fat being untouched. The solvent action of the gastric juice in breaking up muscular fibrillæ and vegetable matter, is greatly aided by the movements of the stomach, which roll its contents about. While digestion is proceeding normally, a sense of well-being and comfort is experienced; a matter which has been practically utilized by the Anglo-Saxon, who invites men to a good dinner before asking them to subscribe to any charitable object. So successful has this been found that it is said, "The road to an Englishman's pocket lies through his stomach."

On the other hand, difficult digestion has a malign and untoward effect upon the temper and disposition, and a stomach-ache is perceptible in many of the sentences in Tom Carlyle's writings. A sudden shock will at once arrest the muscular movements and the gastric secretion, producing a feeling as if the stomach and its contents had been changed into a heavy hard substance. Quietude is conducive to good digestion, and an active man should sit or lie down, after a good meal—just as his dog does! Animals fill their stomachs and then lie down—not having the ambition of man, or being interested in the absorbing acquisition of wealth.

As digestion proceeds the pyloric ring relaxes and the chyme finds its way into the duodenum. The acidity of the stomach seems to exhaust itself in the digestive act, and the chyme is readily rendered alkaline by the bile. In this alkaline condition albuminoids are readily acted upon by the trypsin of the pancreatic secretion which completes their solution.

Just as we saw that starch is dissolved by hydration, so is an albuminoid. An albuminoid previous to solution is termed a "protoid." When rendered soluble by the digestive act it is called a "peptone." It has been rendered soluble by hydration so that it may pass through the wall of the alimentary canal, and enter the portal vein. But, as a soluble peptone would as steadily slip out of the circulation as the kidneys as it finds its way in by the portal venules, it is concentrated and once more made a protoid in the portal blood. (When this does not take place properly the peptone appears in the urine as "false" albuminuria.)

When the albuminoid matters reach the liver, a certain amount

tion is elaborated into the serum-albumen of the *liquor sanguinis* for the nutrition of the tissues. But as the framework of the locomotive needs little repair, so the albuminous tissues of the animal body requires little repairing material from day to day. A sufficiency can be found in a dietary far below what is ordinarily eaten by man, and especially civilized men. For fifty years Lewis Cornaro lived in health and strength and vigour on fourteen ounces of food, mostly vegetable, and twelve ounces of light wine daily. I have a suspicion that the said Lewis Cornaro belonged to that stalwart and somewhat corpulent race of man of whom it may be said "that the less they eat and drink the better they are," and who frequently are glycosuric. But allowing for this, it can safely be said that most people gratify their palates by consuming more albuminoid matter daily than their bodies actually require. This surplusage is called the "*luxus consumption*." What becomes of this *luxus consumption*? It is burnt up, and oxidized in a descending series as tyrosin, leucin, kreatin, kreatinine, passing on to uric acid and urea. It is unnecessary to give the formulæ of these different substances, except uric acid ($C_4H_4N_4O_6$) and urea (NH_2CO). "The antecedents of the urea in blood are in all probability, partly the kreatin formed in the muscle and elsewhere, partly the leucin formed in the alimentary canal, as well as in various tissues." (M. Foster.)

This is a piece of physiology of which we have as yet but broken lights. It is enough for the present consideration that tissue waste and albuminoid excess are burnt up, (mainly in the liver) into the above-stained product, of retrograde albuminoid metamorphosis. And with these excrementitious matters the physician has much to do, especially as years advance. We all know the sensations of well-being, enjoyed by a good meal, under the action of a competent digestive excretions which have ever had a potent attraction for the Anglo-Saxon race, with whom *dam vicinus vicinus* has been largely concerned to do it. But when the digestion of albuminoids is not so good as perfectly, then the case is widely different. Dr. Lauder Brunton, F.R.S., has shown that peptones in the general circulation, act as a general poison; and when the liver is deranged toxic substances, called "toxæmæ" are formed which render the individual, "ill, depressed, and sad," that case.

A case of this kind, once let give me a very vivid account one day of the effects of a meal upon her. At first she had the wonted sense of well-being, but in about two hours after the meal this gave way to a feeling of oppression, a sensation of depression and sense of misery and

impending evil. This condition is not uncommon, the "liver stuffs" or mal-products of deranged hepatic action having the opposite effects of the normal products of digestion.

The question may now be put: "How is it that nearly all of us eat too much meat, or its equivalent and yet only a certain proportion suffer, as being bilious or gouty?" The question is a pertinent one. The liver possesses a good deal of "spare power" as the late Dr. Murchison happily phrased it, and is, usually, equal to dealing successfully with a quantity of albuminoids, far in excess of the body needs. The answer is then "So long as the liver possesses the power of converting this *luxus consumption* into soluble urea all is well!" Soluble urea passes out of the liquor sanguinis in the renal secretion without any difficulty. But not so uric acid.

Urea belongs to the warm-blooded mammalia with a fluid urine; while uric acid is the form of excrementitious nitrogenised matter in the warm-blooded bird and the cold-blooded reptile. In other words urea belongs to the animal with four chambers to the heart; while the tri-chambered heart is found with uric acid. So long then as the liver can convert the *luxus consumption* into urea so long can man, yes, and woman too, eat albuminoids in excess of their body wants; but when this power wanes, and products belonging to a lower stratum of creatures are formed, then it becomes time to pull up and diminish the nitrogenized elements in the dietary. And if this be not done, gout in some of its Protein forms will show itself. Uric acid is the poison of gout; and when it is formed in any quantity in the body then a condition of lithiasis or lithæmia supervenes.

As to the cause of this impairment in the functional power of the liver, it may be said that long continued over-work in dealing with excessive quantities of azotised food is the cause *par excellence*. It may not be the act of the individual but that of an ancestor. The father may indulge in excess, and the son suffer for it in the form of an inefficient liver. "The fathers have eaten sour grapes and the children's teeth are set on edge!" Verily, it is so!

There are, however, other matters of albuminoid descent and lineage, that must not escape our attention; these are the bile acids. Both contain nitrogen; and one contains sulphur, ample proofs of their origin! Glycocholic acid has the formula $(C_{24}H_{44}NO_8)$ and taurocholic acid $(C_{24}H_{44}NSO_8)$. Some persons suffer from "cholelithæmia" while others suffer from "lithæmia" when their liver is out of order. Further some persons who are "bilious" in early life, lose their biliousness and become "gouty" in middle age. Cholelithæmia

and lithæmia equally and alike are causally related to the albuminoid elements of our food. Indeed, the products of retrograde nitrogenised metamorphosis are fertile sources of disorder and disease. Dr. George Johnson, F.R.S., has pointed out that kidney disease is set up by hepatic mal-products passing out in the urine and in doing so irritating the kidneys; and has written this trenchant sentence—*“Renal degeneration is a consequence of the long continued elimination of products of faulty digestion through the kidneys.”*

There can be no doubt in the mind of anyone who has had the opportunity to observe and has availed himself thereof, that when lithates abound in the blood one of two things happen: either some remain in the body as gout; or the kidneys become innred by their output of lithates (the common origin of Bright’s disease), or may be more or less of both are found. Many persons pass turbid urine laden with lithates for years apparently none the worse; but all the same their doom is before them. Sooner or later mischief in the kidneys is set up. We pay a price for pleasing our palates!

The effects of food tell us that there are certain properties connected with animal food lacking in other dietetic matters. The herbivora are widely different in disposition and build from the carnivora. The beef of the English soldiers and sailors has played a part in their exploits, it has always been said. Liebig has written—“It is certain that three men, one of whom has had a full meal of beef and bread, the second cheese or saltfish, and the third potatoes, regard a difficulty from entirely different points of view.”—Dominie Sampson was another man after some of the contents of the Gipsy’s cauldron. Refractory boys of old were subdued by a dietary of bread and water.

Granting all this, it is impossible to disagree with Dr. Pavy in holding that the advantages of a meat dietary have been much exaggerated and overrated. He points to the fact of the oatmeal and milk of the Scotch peasantry. Certain reports of the prison dietaries are referred to by him. Dr. Guy states, “That we possess conclusive evidence of the sufficiency of a diet from which meat is wholly excluded; and even of a diet consisting wholly of vegetable matter.” Dr. Pavy is careful to state that he does not advocate a non-meat dietary, but is opposed to the prevalent view that a certain quantity of meat is necessary to vigour; and, the outcome thereof of the principle, that “it is impossible to have too much of a good thing,” viz., eating more meat than is either necessary or good for the system.

There is another aspect of meat to be considered, viz., the comparative digestibility of the various forms of flesh which must be dis-

cusSED at some length in a future chapter dealing with digestion. Dr. Pavy has given us a reason for the taste for meat, to the effect that the stomach is the seat of its digestion, and that meat stays long in the stomach before leaving it. Certainly in the case of Alexis St. Martin, a Canadian soldier in the U. S. A. army, whose stomach was laid open by a charge of duck shot, at the distance of one yard, at Michillimackinac in 1822, and who survived the wound, this was the case. For some months the patient suffered terribly from exfoliation of the ribs, etc., but thanks to an iron constitution he rode out the storm. In a year the injury had healed, except that the stomach wound did not close, and he had to wear a bandage over the wound to prevent his food from escaping out of the stomach. A little later a fold of the lining of the stomach developed into a valve which closed the orifice, preventing the escape of the contents of their viscous, but readily pushed inwards by the finger, and so admitting a view of what was going on in the stomach. Dr. Beaumont, his medical attendant, paid him the most scrupulous attention; and in return St. Martin permitted him opportunities for inspecting his stomach when in action. The case created an immense attention at the time, but is now disappearing from our literature. Dr. Beaumont found that beef, mutton and pork took three hours for their digestion in the stomach of St. Martin, and cheese three and a half; but for the latter a much longer time would be required with some stomachs. His broad results were that the flesh of animals is not quickly digested. His conclusions remind me of the hearty eater who after an ample fish dinner remarked that "he felt quite ready for a beefsteak."

Of albuminoid food, and especially the flesh of animals, "fish, flesh and fowl," it may be said that it is agreeable to the palate and grateful to the stomach; and further that it is excellent material on which the cook can exercise his skill, and consequently, that it is a tempting viand; that it is a stimulant in its character and is therefore specially suited for those who lead a life of great mental exertion. But that it is not suited for those who are engaged in a sedentary life; and that most of the maladies which fasten on the body as it approaches are causally linked with the presence of an excess of albuminoid waste in the blood.

Fats. The third great element of nutriment, fat, is composed of far more carbon and much less oxygen than the albuminoid food. It has the formula $(C_{55}H_{110}O_2)$. Fat can be made in the laboratory, but rather can be broken up into a fat and a glycerol, and the glycerol

fatty acids are three, stearic, palmitic, and oleic. Stearine is the firmest fat, readily going solid at a low temperature, while oleine is still quite fluid; and palmitic stands between the two. The matter is readily seen in a bottle of olive oil in cold weather. Animal fat contains a larger proportion of stearine, and mutton and beef suet quickly go hard. Broadly speaking, the digestibility of a fat is linked with the proportion of stearine it contains, and with the temperature at which the fat goes firm. Thus beef and mutton suet are less digestible than butter or lard; while the best cod-liver oil has its stearine removed by freezing.

Fat is derived from animals and vegetables, both as fats and oils. Whale oil is too rank for food, at least for ordinary men; though the Eskimo revel in the blubber of the whale and the seal. Cod-liver oil is not attractive to the palate, though some persons overcome their dislike to it, until they take it readily; and a small number even profess to like it. Palm oil when fresh is palatable, and is freely eaten by the negroes of New Guinea. The oil of the African ground nut is a very thin oil, used to thin down and to adulterate other oils. Olive oil is used in cooking, as frying fish, and for salads; but there are reasons to believe that rapeseed oil is largely sold as salad oil. Many seeds are rich in oil, as linseed; while some nuts abound in oil. The cocoa-nut and the Brazilian nut of commerce are instances in point. Why such quantities of a concentrated hydrocarbon should be requisite in the case of certain seeds in tropical climates it is not easy to see. Nor is the passionate liking for fat as seen in the Hindoo's much-prized ghee and the equatorial negro's fondness for palm oil as readily intelligible as is the Eskimo's preference for blubber, which is shared by the Yakoots of Northern Asia.

Fat is a very concentrated fuel-food, and is the form of storage of surplus food, chiefly derived from the carbo-hydrates. The hibernating animal retires helen with fat, and emerge from their hiding-places gaunt and emaciated like the pig buried under the chalk cliff of Dover, which Dr. Mantell has made famous. His sty was a cave in the solid rock, under a chalk cliff; so that when a large portion of the earth fell, he was buried alive. He was estimated to weigh about 140 lbs. when the accident occurred; of course no hope of his survival was entertained, but 140 days later the workmen removing the debris, and clearing the whine, and got him out alive, but only weighing 40 lbs. No doubt he must have perished but for the moisture which exuded from the rocky sides of his cave. During this long period of pining, he lived on his own fat. The Tyrol cream-cheese hunter

has found by experience that a piece of beef suet is the best and least cumbersome food he can carry with him on to the mountains. His choice is a wise one.

Fat is not affected by salivary nor yet by gastric digestion. It is only when gastric digestion is over, and fat comes into contact with bile and the pancreatic secretion, that any change is effected in it. The change is not molecular like the hydration of carbo-hydrates and albuminoids; but is merely an emulsification, *i.e.*, a division into minute particles. These fine particles are small enough to enter the mouths of the lacteals in the intestinal villi, which may be said to "eat" the fat globules. From thence the emulsionised fat passes on to the lymphatics. The history of fat in the body after this is obscure; we know that some of it is taken by the tissues, and the rest is burnt up as body fuel. There is no proof of fat being stored up in the body as fat. Indeed, Ebstein has advocated the substitution of fat for carbo-hydrates in the treatment of obesity; and that too with success.

The difficulty with fat lies with the stomach. A delicate fastidious stomach is offended by the presence of fat in it; and this reacts upon the palate. Some stomachs will tolerate cold butter, which promptly and pronouncedly take offence at warm fat. Rancid fat is objectionable to all civilized stomachs; and compels the most scrupulous attention to all culinary utensils. With some persons, fat turns rancid in the stomach by the formation of an acrid acid, — butyric — causing heart-burn. There is a growing dislike to fat, especially animal fat, at the present time. Many children will no more eat fat than will the cat. They will turn with loathing from the sweet pieces of fat on their plates, yet they will readily swallow the fishy cod-liver oil. Their choice is instinctive, and evidently linked with the fact that they can digest the one, but not the other. The fact remains; and Dr. King Chambers tells the story of a phthisical Scotch terrier who whined at the door of the cupboard in which his cod-liver oil was kept, if it was forgotten.

The late Prof. J. Hughes Bennett of Edinburgh first introduced the use of cod-liver oil into the treatment of phthisis. There is nothing magical about it, it is merely the most digestible form of fat. Still, it is a mighty matter in dealing with wasting disease. The tissues of stroma improve under it. The phthisical patient frequently finds in it the means of curing his disease. Taken an hour and a half after meals it quickly passes out of the stomach, and so gives that viscous little offence. Taken before meals it is an

immediately after, it eructates and disagrees. But taken at the time when the disintegrated food is passing out of the stomach, it quickly reaches the point when fat is dealt with in the alimentary canal. But in some cases, especially when the bowels are the seat of tubercular disease, the oil is voided *en masse* and unchanged. Here the pancreas is clearly unequal to dealing with it; and it is useless to persist with it.

From the fact that fat is emulsified in the intestines, chemists have been led to imitate nature, and form fat emulsions. In milk, fat exists in the form of an emulsion; and the addition of a certain amount of milk to the dietary provides fat for healthy tissue formation, in persons with the most tetchy stomachs. Or cream with a little flavouring like maraschino, may be taken about an hour after a meal. By so doing in many cases it does not become necessary to resort to cod-liver oil. Fat emulsions are more digestible, or rather assimilable than unemulsified fats and oils. There are many fat emulsions in the market, of varying value, and various flavours. Hydroleine and oleo-chyle are cases in point. But there is a difficulty in keeping them, as sooner or later the emulsion separates. One specimen I had for three years before it began to part; but for sundry reasons I have no intention of giving its owner a gratuitous advertisement by mentioning it here. In other cases the oil has been mixed with malt extract; and though this gives an emulsion of coarse oil-globules, in time the oil gets to the top. Then in the *Cremor Hordeatus* of Leoflund the emulsion form of fat is preserved; and *Oleo-byne* is an emulsified fat. These last contain fat with soluble carbo-hydrates, and are palatable as well as being capable of being kept for any length of time in their tins. Such preparations are indicated where the palate objects to cod-liver oil; indeed the fat in both cases is a better fat than the oil; while the form of emulsion is favourable to assimilation. They form indeed very complete foods for invalids and travellers. With a little farinaceous matter, as a biscuit, and a little milk, they are well adapted to gouty persons and those who suffer from Bright's disease.

Fat-eating among adults is fortunately on the increase at the present day; if the valuable hydro-carbon is in growing disfavour in the nursery. Intelligent persons are now eating fat because they know it is good for them. Whenever there is any tendency to tubercle, the fat-eater should learn to eat fat, just as a seafaring man learns to swim. As a physician to a Chest Hospital I have learned to dread the announcement that fat is no longer taken, es-

pecially if the individual is of strumous build, with a small narrow chest. In my opinion the existence of a considerable area of affected lung where the digestive powers keep up, is less fraught with evil and less prognostically significant than intractable wasting with very little disease in the lung.

The statement often made that "fat is bilious," as being a direct irritant to the liver, is incorrect.

Certain it is that when the dietary is improved in some cases the liver is upset by the new demand upon it; but this can always be met by a cholagogue and a purgative.

Salts. Certain inorganic principles are also essential to our health, mainly chlorides and phosphates. When certain vegetable salts are lacking in the dietary, the disease known as scurvy is induced. The early navigators of the oceans which lave the shores of the new world were terribly afflicted with scurvy, in consequence of their dietary of biscuit and salted meat. Any vegetable matter, especially "scurvy" grass (*cochlearia officinalis*) a member of the Cruciferae, at once brought relief. Land scurvy is occasionally seen in persons who abjure vegetables, as do some dyspeptics. A cruel old Dutch punishment was to feed prisoners on bread made without salt. Much obscurity still overlays the functions of these body-salts. "We know that the various saline bodies are essential to health, that when they are not present in proper proportions, nutrition is affected, as is shown by certain forms of scurvy; we are aware of the peculiar dependence of proteid qualities in the presence of salines; but beyond this we know very little." (M. Foster.) A good illustration of the value of the body-salts is furnished by watching the effects of arresting the night sweats of phthisis. As soon as the outpouring of the salts in the sweat is checked, the appetite returns and soon the pallor of the features gives way to the hue of returning health. The effect is often almost magical. Chloride of sodium, our common salt, is a palatable addition to our viands and especially to the flesh of animals. Probably from it is furnished the hydrochloric acid of the gastric juice; while the soda goes to the liver, as traceable in the base of the glycocholates and taurocholates of the bile. The phosphates too are equally important. Phosphate of lime is necessary for the bones of the body, and is furnished by the vegetable and animal food, and in milk. "Caseine is a nitrogenous phosphoric compound, conspicuous for the tenacity with which it holds its phosphoric phosphate of lime incorporated with it." (Payson.)

It is found indeed in all the structures of our body, and in

Lecithin, the food of the nervous system, is a phosphorescent fat. Then potash is requisite for the muscles, and red corpuscles of the blood. Why there should be phosphates and potash in the blood-corpuscles with chlorides and soda in the plasma around, we do not yet know. Indeed, this is part of the darkness undispelled as to the body-salts. "Although so closely allied in their chemical properties, potash and soda cannot be made to replace each other in the living system, and the same is likewise true of lime and magnesia." (Pavy.)

Whatever the peculiar part played by saline matters in the body, we know their importance, by the consequences of being without them. The phosphates, the chlorides, lime, soda and potash, are essential to health and life.

Water. Water undergoes no change in the body, nevertheless it is a most important matter. Solution is an essential part of digestion. Our tissues are bathed in fluid; our secretions flow. If hunger secures the taking of food, a still more imperative sensation provides for the necessary fluids. Thirst is the security for our supplying our tissues with a sufficiency of water. When cut off from food and drink it is thirst which drives the sufferer delirious, and shortly closes the scene by death. Hunger kills more slowly. Water is even more essential to life than food itself. Water too, bathes the tissues and washes away our waste and excrementitious matter. The vascular system, like the canals of Venice, provide the food and take away the waste and offal. Water has, too, other properties in the body.

"Water does not itself undergo any chemical alteration, and hence is not susceptible of liberating force - does not in other words, constitute a force-producing agent, but it contributes to chemical changes by supplying a necessary condition for its occurrence in other bodies." (Pavy.)

CHAPTER III.

METHODS OF PREPARING FOOD.

IN the preparation of food of yore, the palate has no doubt been the great guide. Still, a blind instinct seems also to have been at work. The cook has been led to prepare vegetables with meat; to unite grain and milk; to boil the highly nitrogenised beans with fat bacon; or peas and pork. Experience at work through countless ages has no doubt instructed Man, albeit darkly, what combinations of foods are requisite for health under certain conditions. Sauer-kraut was a wise provision of vegetable food during the long winter, when salted foods and cereals formed the chief dietary of the people. A Lenten fast of vegetables was a useful hygienic measure for clearing away the maladies incidental to such a dietary; as useful as vegetables to scurvy-stricken crews before the days of lime-julee. No wonder people long ago spoke of the anti-scorbutic properties of certain vegetables.

Before proceeding with the preparation of foods, it may be well to give a Lethaby Table of the comparative value of various edible articles as tissue-food and fuel-food, without any pledge as to the absolute accuracy of it. It is certainly useful, as giving a good broad idea of the value of various comestibles:

	CARBON.	STEROGEN.
Fresh Butter,	64.56	—
Dry Bacon,	59.84	.95
Dripping,	54.56	—
Green Bacon,	54.26	.76
Lard,	48.19	—
Suet,	47.40	—
Salt Butter,	45.85	—
Fat Pork,	41.13	1.06
Cocoon,	39.31	1.40
Cheddar Cheese,	33.44	3.06
Indian Meal,	30.16	1.20
Sugar,	29.55	—
Oatmeal,	28.31	.36
Rice,	27.32	.68
Seconds Flour,	27.00	.76
Split Peas,	26.99	.48

	CARBON.	NITROGEN.
Rye-Meal,	26.93	.86
Pearl-Barley,	26.60	.91
Barley-Meal,	25.63	.68
Ttende,	23.95	—
Bakers' Bread,	19.75	.88
Skim Cheese,	19.45	4.83
Mutton,	19.00	1.89
Beef,	18.54	1.84
Red Herrings,	14.35	2.17
Bullocks' Liver,	9.34	2.04
White Fish,	8.71	1.95
Potatoes,	7.69	.22
New Milk,	5.99	.44
Pusnups,	5.54	.12
Carrots,	5.08	.14
Skimmed Milk,	4.38	.43
Green Vegetables,	4.20	.14
Turnips,	2.63	.13

In considering the methods of preparing food, it may be well to commence with Meat.

Meat. — *Boiling.* One of the most ordinary methods of cooking meat is to boil it. Of old a piece of meat, a ball of oatmeal, and possibly some greens, (when they are to be got), were all put into a huge pot, which was then heated. The old Border rule was, “no broth no ball; no ball no beef.” That is, first came a basin of broth; then some of the ball; and after that when the edge had been taken off the appetite, the beef. If the broth was not supped, there was no ball; if the ball was not eaten there was no meat to be looked for. In those days, when the broth was of moment, the meat was placed in cold water—about a quart to the pound of meat. “It should be covered with water during the whole of the process of boiling—but not drowned in it—the less water, provided the meat be covered with it, the more savoury will be the broth.” From fifteen to twenty minutes for each pound weight is the (Kitchener) time. But freshly killed meat, *à la*, while the rigor mortis is on, takes a longer time than meat which has been kept till in culinary phrase it is “ripe.” Meat loses from ten to about 30 per cent. of weight in boiling.

When the “soup,” the fluid in which meat has been boiled is an object then the meat must be put in cold water and not brought to a boil; but, as the meat is the first consideration then the water must be boiling when the meat is put in it. “When it is taken from the boiling water, the albumen of the flesh is immediately coagulated on the surface and to a certain depth inwards, thus

forming a skin or shell, which no longer permits the juice of the meat to flow out, nor the water to penetrate into the mass. "The flesh continues juicy, and as well flavoured as it can possibly become. The greater part of the savoury constituents is retained in the meat." (Liebig.) Meat so prepared is very much more toothsome than meat placed in cold water.

In boiling fish some prefer to heat the water, and add the different matters as salt and vinegar; then to take the pan off the fire and put the fish into the boiling water allowing it to remain in the water the same time that is given for boiling it. One gentleman who has his fish so "boiled" said the plan worked up to an 18lb. salmon. It is asserted that the flesh is more palatable this way than after ordinary boiling.

Boiled meat puddings are very savoury. The ordinary beefsteak and kidney pudding is well known, but the contents of the paste may vary; in Suffolk pork is substituted for beef; in Oxfordshire liver and bacon are placed in the paste. The best thing to do with an old partridge is to let it hang till it is high and then make a boiled pudding with it, packing some pieces of fat and lean mutton around it.

Custom has settled the choice of joints for boiling and for roasting. In the case of a ham it is well to leave the ham in the fat till the water has gone cold; especially is this desirable where the ham is a dry one. When dried fish is to be boiled it is well to steep it in lime water first. This conserves the phosphoric acid as "bone earth" or phosphate of lime; otherwise the phosphorus would be lost, and with it much of the food value of the fish.

Roasting. Probably this was the earliest form of cooking meat. The savage could put a piece of meat on a stick and have it kept before the fire. Before the days of the "jack," dogs called "turnspits" were engaged to keep the meat turning before the fire. As well as turning, basting is required to keep the meat from burning. The time for roasting is a little more than for boiling. On the whole roast meat is more savoury than boiled meat. Roasting is a method of dealing with fresh meat; while stewed meat is cooked. True to his theory of coagulating the albumen of the meat, Liebig exposed to preserve the juices. Liebig advised that the joints should be placed near the fire, and then after the surface was browned to withdraw it a little. Usually however, the cover is put on, and has the sanction of Dr. Kirchmeyer. "The basting of roasting is a form of skin religion duty. A meat-cooker or a cook is bound to

as it saves coal by keeping both sides hot." The fat which exudes is saved as dripping, while the meat juice is sent up as gravy, which gives relish to the vegetables eaten with the meat. As to whether the meat should be raw or well done is a matter upon which tastes differ. When the roast meat is destined to appear on table in cold form only it is well to avoid cutting into it when hot. Liebig's idea of condensing the juice in the meat is correct enough in this respect.

Baking. The convenience of the oven is a great temptation to its use. It is said meat loses less weight in baking than in any other culinary treatment of it; but what it gains in weight it loses in flavour. Dr. King Chambers says, "No kitchen is complete without an open range. It is impossible to have a properly roasted joint by any other means, as I learned by visiting the private premises of a 'Patent Kitchener' manufacturer; finding there an old-fashioned fireplace in full operation. He cared too much for his diet to employ his own works."

An oven, of course, is indispensable to a meat pie. Meat pies bear the reputation of being indigestible, and probably they deserve it, but whether this is due wholly, or only partly to the paste, and meat so treated is not itself indigestible, is a matter not yet determined. The addition of eggs which get as hard as leather may be sound practice from a cook's point of view, but the physician and physiologist can not be expected to be in favour of it.

Broiling. This is a very savoury way of treating a slice of meat, be it a mutton chop, a beefsteak, or a veal cutlet (which is often fried). Broiling belongs too essentially to the cook's craft to be discussed here.

Frying. For slices of meat, frying is an abuse except for bacon or ham. It is, however, an admirable way of treating fish, though it may be questioned if a fried herring is equal to one properly broiled or brandered.

The flat fishes are specially adapted for frying, and a fried sole is savoury food. But if the cook's fat be not sufficiently hot it soaks into the fibre of the fish and few edibles are more indigestible than a badly fried sole. With looser-fibred fish this carelessness is less injurious.

Frying is a surer plan for heating up cold vegetables, as potatoes, peas, puddings, and cabbage, which has much to be said for it from the point of view of the palate and digestion.

Steaming. This is the most economical method of cooking; it is especially adapted to households where strict economy must be prac-

tisen. As Dr. Buckmaster observes, "All and every kind of meat will do for a stew," and "All vegetables may be cut into slices and cooked in a stew, except potatoes." All gristly meat and trippings with offal, as sheep's trotters and pig's feet, will do for a stew of course, as he adds, "the better the meat, the better the stew." An Irish stew consists of mutton, potatoes, and onions, and an admirable as well as economical dish it is. Indeed, such compounds of meat and vegetables in a stewpan and well cooked would be most acceptable to most households, but are comparatively neglected in favour of joints. The long action of heat would affect the starch-granules favourably and thoroughly detach the muscular fibrille from each other and so prepare the food for the action of the different digestive agents and processes. As ordinarily prepared a stew is deficient in fat. But care and knowledge are requisite for a good stew and also time. "Time" is what all skilled cooks beg for in cooking. Haste in the preparation is the ruin of stews. Curiosity, too, is a drawback: the lid of the stewpan should never be lifted until it has to be taken off. Then the savoury compound is sapid, fragrant and appetising. It is also most digestible.

"Half subtilized to chyme—the liquid food
Readiest obeys th' assimilating powers."

This was written long before science came to bear its testimony to the truth contained therein.

Soup. Soup is alike a means of gratifying the palate of the wealthy gourmand, and feeding the poor in times of famine. A light soup is very suitable food for the sick. Simply, soup is an infusion of meat, flavoured with a little salt and some condiment. At the rich man's table, soup is a very complex affair. It consists of stock (the simple soup)—gelatine, which goes fluid on heating, often, cream and vegetables of all kinds and sorts; sometimes it is thickened with broken biscuit, as is done in the Channel Islands, and a very nutritious food is soup so constituted.

In some cases, fish is the chief meat constituent. In other cases, legumes take the place of meat, wholly or partly, especially with vegetarians. Like stews, soups, looked at from their food value, are unsatisfactory as being too poor in fat. Whenever cream is added, the objection falls to the ground. Soup precedes other food, and takes the van in a meal, as furnishing material for the action of gastric juice. A little meat-juice to furnish the pepsin (for the old body) and a little dextrine for the lacteal material (for the

stomach brews its own peptic solvent, while the viscus is filled and the palate gratified. The French invariably take a glass of wine after the plate of soup, "*le coup d'après*." Indeed, many soups contain some wine. Old cooks over-season soups, which should not be too highly seasoned—in order not to deaden or dull the gustatory nerve-fibrils. Really, the soup should just be seasoned sufficiently to stimulate the palate. Nor is it good for the lining of the mouth if the soup be served up scalding hot. On the other hand, cold soup is very unpalatable. The meat of which the stock is made, ought to be pounded and put through a sieve, in order to restore the natural condition and properties, and give the meat infusion a true nutritive value. Then some broken biscuit, or some bread-rasplings, or boiled potato put through a sieve, should be added, to supply the farinaceous element or conversion into grape sugar. In some cases, fine sago is used. White soups made with milk, and especially when some cream is added, are very nutritive—as containing a distinct proportion of fat—and very palatable.

Raw Meat—Some physicians have advocated the use of raw meat, scraped or pounded, in certain cases. The advantages of such meat are not obvious to the physiological physician. Indeed, Dr. J. H. Bennett discountenances such practice in these words:—"To give meat raw, however finely chopped, is to forego all the advantages gained by judicious cooking, and to force on the digestive organs double work." (Nutrition in Health and Disease.) Many of the advocates of raw meat have had to deal with hysterical women—a class so totally opposed to rational beings in every way, that it is no matter for surprise that uncooked meat agrees with them. The habit of eating uncooked meat has subjected those who indulge therein to trichinosis; and other cutozoa are common in Eastern Europe, where this objectionable practice mainly obtains. Meat juice and cold drawn beef-tea have not, at least, this last objection to their use. In milk, we find an albuminous body, which does not require cooking, certainly. Considering that all proteids must be converted into peptones before they can pass from the alimentary canal into the blood, any advantage to be gained from raw meat juice is quite hypothetical; and involves a lack of acquaintance with the physiology of the digestive act on the part of those who resort to such raw meat.

In the cooking of meat, there is no clinical or even molecular change involved in the proteid material. The muscular fibrille are loosened by the action of heat, while the coagulation of the albumen

renders the fibres more brittle. Consequently, cooked meat is more easily masticated than raw meat. While the further disintegration of the flesh is facilitated by this change, and disintegration precedes solution in the digestive act.

Starch. Bread Making. Starch is the characteristic element in bread, "the staff of life." When wheat was first bruised in a quern, a species of mortar, or ground by the "two women sitting at the mill," all the elements of the seed were in the flour. Long before chemistry could tell them the folly of what they were doing, cooks, led by the eye, began to prefer white bread. In order to attain this end the miller devised means to remove the external coat of wheat as bran. Unfortunately in this external layer lies the bulk of the salts in wheat. By this fateful gratification of the eye an error wrong was inflicted upon the young and growing creatures. An erroneous taste dies hard, and the attempts to induce the proletarian to adopt whole meal bread have met with little, very little success as yet. Those who adopt whole meal bread have usually passed the growing age.

Flour, as now made, is far inferior to that of earlier days. The destiny of flour is the oven. Some of it may go another gate, but the bulk goes to the oven. Mixed with water, a little salt and yeast, flour is worked into dough, is set before the fire to rise; after which in loaves it is placed in the oven to be baked. Yeast is a fermenting agent (a form of lowly plant life—yeast fungus). "The presence of yeast in a substance containing sugar, or starch, which is convertible into sugar, and nitrogenised matter, induces certain chemical changes, comprehended under the term vinous alcoholic fermentation." These changes in the making of bread consist in the conversion of the sugar of the flour into *alcohol* and *carbonic gas*; the latter in its efforts to escape from the dough in which it is made, distends it, forming vesicular spaces in its interior, and so causing it to become porous or light. Much of the alcohol is dissipated in the process of baking.* During the baking, part of the starch is doubtfully converted into dextrine (A. H. L. Hess, *ib.*). They also see that some insoluble starch is converted into soluble matter in the process of baking. The bread is $\frac{1}{3}$ the ovenable weight of the three quarters, and the temperature of the oven is from 400° to 500° Fahr. If bread could be exposed to a higher temperature, it would

* This will be welcome news to the patient who is told that he might be led to exclude bread from his diet, because it is "alcoholic."

kept in the oven a longer time without burning, the conversion of the starch into dextrine would be more extensive and complete.

From its vesicular or spongy condition bread is readily masticated and mixed with saliva in the mouth. For delicate children, in whom the saliva is deficient in quantity, or in quality, some guiding principle led shrewd persons to use only the crust of the loaf, which contains more dextrine than the crumb. This was a wise choice. Then came the preparation of rusks and tops-and-bottoms. Finally came "baby foods," of all kinds, a matter to be discussed further on. The principle underlying all "baby foods" is the conversion of insoluble starch into soluble matters. Newly made bread is a poison to most dyspeptics. It is moist and readily goes into a lump in the mouth; while stale bread is drier and of firmer consistency, and does not readily lose its vesicular character in the mouth. Hot breakfast rolls are toothsome but not digestible.

The reason why boiled paste is indigestible with many persons is that it is not vesicular, but a solid mass. When flour has been mixed with fat, as lard or butter, as is done in making pastry, the starch and fat are so intimately mixed and incorporated that the saliva cannot get at the starch granules. Further, too, disintegration is not so easily performed as with bread, and between the two the stomach is handicapped with masses of unchanged starch and fat, upon neither of which has it any solvent action. No wonder then that it aches!

As the confectioners' wares consist largely of flour and fat, well kneaded into paste, they are shunned by the parents and guardians of children of tender years, from the ulterior consequences of eating them. Not only do these masses offend the stomach but they load the bowels; except when their irritant presence sets up diarrhea for their removal.

The children of the United States are reputed a dyspeptic race and certainly a number of representatives of the great Republic have found their way into my consulting room, suffering from indigestion. In consequence of personal acquaintance with the New World I am obliged to rely upon what I hear, and upon American cookery books. In regarding the instructions as to the making of "corn-cakes," I find "Bake quickly and steadily in a buttered skillet. Do not more than hour will usually suffice." This is not long for the action of heat upon starch. Then it is to change these cakes into eaten cakes. Very little time for the saliva to act upon the starch. In the name of common sense can we wonder if the

outraged stomach sometimes protests? Hampered with enormous quantities of unchanged starch it cannot perform its work without effort.*

If corn bread or cakes were eaten slowly much of the resultant dyspepsia would be avoided. The albumen of maize or Indian corn is not gluten. Consequently it will not alone make bread. The combination of maize flour with wheat flour is what is required for a readily digestible bread. All boiled paste, all boiled puddings like a suet dumpling, or a meat pudding-paste, should consist of a union of the two kinds of flour. All baked puddings should be made of the two, especially when eggs are used. Wheat flour and eggs form a compound not easily disintegrated, however palatable. Certainly paste should contain the largest practicable proportion of maize flour. If the medical profession were only generally familiar with the properties of gluten they could give a direction to the dietary of the nursery of a desirable kind.

In the North of England cakes of all kinds are in vogue, and the frying-pan and the "girdle," or griddle is in constant request as well as the oven. One form of cake is disappearing. When an addition was made to the family group, a cake was prepared by adding plenty of butter to some flour, throwing in a few currants, rolling the paste about a quarter of an inch stick, and placing this in the frying-pan if the oven was not available. While cooking it was divided into squares about the size of the palm of the hand. When cooked it was still a sodden mass, but toothsome. The little squares were then split and butter added freely to the warm cake. Old-fashioned doctors liked this cake, and drank it down with tea (said to have been "laced" with rum) to prepare them for the journey home. It is dying out, probably because the less potent digestive organs of the present generation cannot deal with it. Paste rolled out thin on a circle, with jam or fruit laid over one half, while the other half is folded over, and the edges closely pinched together, is a common cake, or pastry in the North. Then there are Eccles cakes, and Barbury cakes, combinations of flour, fat, currants and small pieces. The early emigrants to New England took their taste for cake over with them; and sundry rolling-pins were in the hold of "M^{rs} Flower." we may depend upon it. The bane of many a southern cook is their indigestibility on account of their forming so dense a mass in the mouth, which will not fall to pieces in the stomach. Some of the

* Of course this is not the sole cause of indigestion in the stomach.

cakes are tempting to the palate, especially in early years. A story is told of a little girl who wanted more cake. "She was told if she ate any more she would be ill, and the doctor must be sent for." This effective deterrent on all ordinary occasions failed in this emergency—"Just anoizzer piece, and then send for the doctor," she pleaded. Fat and flour sweetened form a good food from a chemical point of view, but the resultant product is unfortunately not digestible; nor are flour and eggs as in sponge cakes much better. The addition of maize flour to wheat flour gets rid of some of the difficulties of pastry; but not of all of them.

Starch of various kinds is in use for milk-puddings. For children, invalids, and dyspeptics such milk-puddings are admirable, unfortunately dyspeptics usually abhor them; a milk-pudding is usually allowed an hour in the oven so that the time for the action of heat upon the starch allowed is double that of cakes. The combination of starch and milk is capital food. Unfortunately the cook will add eggs, which increases the food value doubtless, but detracts considerably from the digestibility. The addition of butter is good. If the various forms of starch, as sago, rice, tapioca, etc., were subjected to heat before being added to the milk the conversion of starch would be greatly facilitated. Acting on this principle, I have taught my cook to put the farinaceous matter in a little pie dish in the oven, dry, and bake it for an hour or two, taking care not to scorch it by having the oven too hot. This does not impair either lightness or the flavour of the starchy matter. The conversion of starch is proved both by the iodine test, and by artificial digestion either with malt or a malt-extract. The most perfectly digestible milk-pudding is made by adding some ground malt to the baked starch, when the warm milk is poured on, and stirring the whole together, after that setting it in a warm place for ten minutes before putting in the oven. The diastase of the malt acts upon the cracked and changed starch granules, and the resultant product is fit for the weakest baby, invalid, or dyspeptic. Indeed, ground malt is a much better addition than cane sugar wherever such sweet milk puddings turn acid in a fastidious stomach. (There is however one point on which the cook must be put upon her guard, *i.e.*, not to add any ordinary sugar to a pudding containing malt. It is sweet enough with the maltose sugar. Especially is this caution to be observed in those cases where ordinary milk-puddings sweetened with sugar turn sour in the stomach.) Indeed, ground malt alone with a little milk makes a very nice pudding, and is improved by an egg.

The principle of employing farinaceous matter which has already been subjected to heat (so that a considerable conversion of starch has gone on before the human salivary diastase comes into play), is carried out in practice in the form of bread and butter pudding. Slices of stale bread buttered are placed in a dish, some currants are thrown in, and then some sweetened milk poured over all; and the dish is placed in the oven; or bread is crumbed and milk added; but this pudding lacks the fat of the first form. Or a milk-pudding can be made of broken captain's-biscuit (which is baked at a high temperature). The substitution of malt for cane sugar, as described above, is all that is required for an ideally digestible dish for delicate stomachs; the flavour of maltose being also finer than that of cane sugar.

When milk puddings are served at table as a rule they lack fat. If hot it is well to place a knob of butter in the middle of the mass of pudding on each plate; it will melt before the pudding is cool enough to eat, and mixes readily. Very fastidious personages would perhaps say that such a practice is "messy," and object to "making mortar," but these objections do not outweigh the real value of the addition of the butter. If cold, like crested rice, it is well to eat butter and sugar, rather than sugar alone, to the cooked farina.

The Potato.—The potato lies midway betwixt starch and vegetables. It is a vegetable rich in starch; and boiled potatoes mashed have been used as an adulterant—well, hardly that—to flour in preparing dough for bread. The object of cooking a potato is to transform it from a firm indigestible vegetable into a ball of flour. About the cooking of a potato opinions differ. Some boil, some steam, some pour off the first water; some pour off the water when boiling is completed, and leave the potatoes a little time in the pot. Before the potato disease in the Irish famine of 1846, potatoes were commonly boiled in their "jackets," or skins. When boiled enough to take the potato in both hands, and with a gentle pull off came each half of the peel, and the floury ball dropped into the dish. Hard potato is utterly indigestible, and potatoes cannot, like other vegetables, be used to stew with meat. Mashed potatoes are most comfortable. The cooked potato is beaten up and mixed with cold, or hot butter and milk, until a light white product is obtained. This is most palatable if well made; as such it can be eaten as a vegetable, or mashed potatoes and sausage are familiar and comforting in the house-lunches; or a wall of mashed potatoes is made.

meat, or a cottage pie. Mashed potato with salt and pepper and set before the fire to brown, with a glass of milk, forms an ideal lunch-con, according to King Chambers. The introduction of the potato revolutionised the fare of Europe in winter. Sauer kraut the Germans had; and pickled gherkins were common over the most of Europe, and in Southern climes some greens were possible; onions too they had; but, after all, their winter dietary was deficient in vegetable food, and consequently scurvy, and other scorbutic affections, were rife. The monks cultivated the wild carrot and turnip as luxuries; but these were only for the well-to-do. The potato, which keeps well with a little care, at once supplied the desired vegetable food, and the winter fare of man was no longer without the desirable elements.

Vegetables.—The vegetables in common use are roots and greens, while celery is a stem like rhubarb. In roots as the turnip, carrot, parsnips, beet, mangel-wurzel, also the onion, etc., there are quantities of starch and sugar, while in greens the leaves consist of vegetable parenchyma of low food value but mostly rich in salts. The most common are cabbage, cauliflowers, broccoli, Brussels-sprouts, spinach, lettuce, endive, salsify, and others. Celery is an excellent escent food. In speaking of the value of alkalies in our food and especially phosphates, Liebig says: "The sure guide of man and animals has taught us how to counteract the deficiency of yeast, fish and eggs, in alkalies by the addition of vegetables, potatoes, or salad. Kitchen vegetables, in this point of view, fill up many blanks. The quantities of salts of alkalies and alkaline earths, contained in many of these, excites astonishment. Celery contains from 16 to 20 per cent., common salad 23 to 24 per cent., and cabbage-heads (buds) 10 per cent., of the weight of the dry part in the form of incombustible salts." Of the value, then, of vegetables, there can be no doubt; nor of the sure-footed instinct which led man in the days of yore to boil vegetables in the pot with his animal matter. The cruciform, especially cabbage, however, do not go into this pot, but into one of their own on account of the offensive smell, "green-water" possesses.

The cooking of vegetables is a great art; more especially greens. "They are best," says Dr. Knebelson, "when in full season;" "turnip vegetables are not so good as mangel-wurzel and mangel-pe fruit." "Fresh gathered" is the motto for vegetable gastronomists. An orange or a banana in London or New York is not the same thing from the same fruit plucked ripe from the tree. Root, green, and salad are the

same. Potatoes should be dug out of the ground the last possible moment before cooking, to do justice to them. With roots and greens when stale it is well to put them into cold water, which will freshen them up a little. Vegetable matters are succulent and lose water rapidly, which impairs them in every way. They should be plunged in the pot when the water is boiling. They are a ticklish affair to cook. "Raw" vegetables are unpalatable; while if "vegetables are a minute or two too long over the fire, they lose all their beauty and flavour." Fresh vegetables require considerably less time for sufficient cooking than do stale vegetables. The advantage of life in the country is great in the matter of vegetables. If taken fresh, they, however, preserve well in tins.

After being carefully picked, greens should be washed in cold water and then drained, after that boiled with a little salt; when cooked they should be drained well and served hot.

Cabbages, as eaten in London, are an abomination. They should be treated as in Germany and the United States. When boiled, "drain well, chop and stir in a tablespoonful of butter, with pepper and salt." Such cabbage is also good cold; and heated up next day it is excellent. Like soup, cabbage is better the second day than the first. But served as they are in the South of England they are a very poor affair.

In the North of England, in early spring, it is customary to gather the leaves of a wild lettuce with the young shoots of the common nettle and chop them fine, then to mix some pearl barley with them and boil in a bag. This is called a "herb pudding" and it was no doubt a valuable food in old days for scorbutic affections. The "nettle" beer of the Midlands no doubt had a claim to be regarded as anti-scorbutic, especially when hearnhound was added.

Salads.—Uncooked vegetables are usually eaten as salad. The lettuce, the endive, the dandelion, mustard, cress, and watercress alone or mixed, are the main constituents of salad. Onions and onion form a savoury salad. Potato salad is not much valued in England. Some like a little sorrel in their salads; others omit it. Tomatoes, Celery, radishes and watercress are as often used as the lettuce. Some persons prefer to dress the salad on the day of eating it, others prefer it dressed before it comes to the table. As to the old Spanish proverb run to this effect, "To be a good salad there should be a miser for oil, a spendthrift for vinegar, a wise man for salt, and a madcap to stir and mix," it is not gathering much ground. No steel knife should be used. All

picking and washing them, "let them remain a while in the colander to drain; lastly, swing them gently in a clean napkin," then place them in the salad dish and prepare the dressing, and "pour it down the side of the salad dish—and don't stir it up till the mouths are ready for it." So says Dr. Kitchener, who adds—"We recommend the eaters to be mindful of the duty of mastication—without the due performance of which all undressed vegetables are troublesome company for the principal viscera, and some are even dangerously indigestible." Beetroot boiled in slices is commonly added and makes the dish look well. Then cold meat is often added. The Italians used cold chicken; the Dutch, cold boiled turbot or lobster, and in spring, young onions.

As to salad dressing, it can be bought, but it is well to make it, when and as wanted.

The following is Sydney Smith's recipe for it.

"Two boiled potatoes strained through a kitchen sieve,
Softness and smoothness to the salad give;
Of mordant mustard take a single spoon—
Distrust the condiment that bites too soon;
Yet deem it not, thou man of taste, a fault,
To add a double quantity of salt,
Four times the spoon with oil of Lucca crown,
And twice with vinegar procured from town,
True taste requires it, and your poet begs
The pounded yellow of two hard-boiled eggs.
Let onion's atoms lurk within the bowl
And scarce suspected animate the whole;
And lastly in the flavored compound toss
A magic spoonful of anchovy sauce!"

It will be seen that he is not at one with the Spanish proverb as to the proportions of oil and vinegar. He makes no mention of sugar, which is indispensable in a salad in which there is sorrel. Then the French have very compound salads, and add black pepper, chervil, and sometimes a scrap of cheese. Even the dissolving the salt in the oil rather than the vinegar is insisted upon by some gourmands. There can be no doubt that the addition of a few drops of Worcester Sauce is good. Carminatives are desirable in salads, for vegetables are "windy." It is needless to say that salads are unsuitable for dyspeptics and those who possess tender bowels.

As a wholesome article of diet, salads are too much neglected in many households, where they might be adopted with advantage.

Fruit. Fruits are part of the food of man, and in some areas a

very large factor in the dietary. They are very wholesome, though in some country districts, fruit is regarded as unwholesome and tending to set up diarrhoea. Certainly, if a child is debarred from access to fruit, it will take the earliest opportunity of gorging itself on any fruit which presents itself; and not be very particular or select about the quality either. If the fruit be unwholesome—either unripe or over-ripe, the best thing that can happen is for the alimentary canal to get rid of it as quickly as possible. But sound, ripe fruit is wholesome and good; and if it formed a part of every meal, children would be pretty safe against indulging in indifferent fruit. Cotton Mather said: "Doubtless God could make a better fruit than a strawberry, but doubtless he had not." Others might prefer a jargonelle pear, or a muscat grape, or a mango, while others would prefer the orange. All fruits contain levulose sugar, and usually some acid in varied proportions, ranging from the pineapple to the lime. They all contain alkaline salts, which are good for the blood. "This influence of the alkalis is shown in a decisive manner, in the effects produced on the salts of organic acid in the circulation. It has long been observed that, after eating juicy fruits, cherries, strawberries, apples, etc., the urine becomes alkaline. All these fruits, as well as the juice of edible roots, tubers, and green vegetables, contain these alkalis in the form of salts of the organic acids; usually as malates (all kernel-fruits, pineapples) citrates, (stone-fruits, currants, potatoes) tartrates (grapes)." (Liebig.)

The utility of such fruits in the uric acid diathesis is apparent and obvious. If gouty persons would learn to eat fruit and teach their children to follow the example set, it would save much resort to medicine in later days that might well be avoided.

The growth and decay of fruit is thus described by Pavy: "During its progress, it increases more or less rapidly in bulk and weight; and as it approaches maturity it loses its green color, becomes brown, yellow, or red, and no longer acts on the air like the leaves; on the contrary, absorbs oxygen and gives off carbonic acid. As the process advances, some of the proximate principles contained in the unripe fruit, particularly the vegetable acids and tannin, gradually disappear, apparently by oxidation, and thus the fruit becomes less astringent. At the same time, the starch it contains becomes soluble into sugar; and the insoluble pectose into pectin, and other soluble substances of allied composition, are liberated from their insoluble condition. The fruit in this way enters into a condition favorable for eating. Oxidation, however, still continues, and the

sugar and remaining acid become destroyed, giving rise to the loss of flavour which occurs after the full ripened state has been attained, and deterioration has set in." From this we can readily understand why unripe fruit, as well as over-ripe fruit, is not wholesome and salutary. "The vegetable acids and the pectin disappear in the body speedily after being eaten; while the alkaline base appears in the urine as a carbonate;" though if uric acid be present, as a urate.

Some fruits are preserved by drying them, like the fig, raisin and currant; while others are preserved by boiling them with cane sugar, as guava jelly, strawberry jam, etc., while in the South of France, candied fruits are a large article of manufacture. The addition of the cane sugar in the two latter cases, gives such fruit a tendency to turn acid in the stomach. Of old, fruits were bottled, and now they are tinned extensively; and when so preserved, keep perfectly good for years.

At one time, preserving was a household affair, and preserves were home-made. Now, preserving is a business. The introduction of apple and turnip into marmalade, sent the best firms to Spain for their manufactories. Then raspberry jam was adulterated with the pulp of gooseberries. Dried figs, of indifferent quality, with their seed, offer great temptation to the rascally manufacturer; and when mixed with plums that have seen their best day, and boiled with a cheap glucose (made by boiling starch with sulphuric acid), and finally flavoured with various essences, the resulted product is sold as "Family Jam." Such preserves are used for jam puddings, and are eaten on bread. They have a food value of their own; but are far inferior to butter, for which they are too commonly substituted.

When sugar is boiled with butter, the well-known toffee (a very excellent hydro-carbon), is furnished. The old-fashioned plan of making toffee extensively in cold weather had much to be said for it. It supplied an extra ration in fact, of admirable and concentrated fuel-food, to meet the external cold; while it also supplied fat for healthy tissue-growth. When delicate children object to fat, and their stomachs will not tolerate cod-liver oil, it is no bad practice to encourage them to take toffee. Of course, the mother smiles derisively at the idea; but the results are often very satisfactory. And such union of carbo-hydrates and fat might often be utilized with advantage with delicate children.

Such then, are the various forms of food in common use with man, and their methods of preparation. Their food-value and their digestibility vary widely—not only in the material itself, but also in the methods of preparation.

CHAPTER IV.

CONDIMENTS. SAUCES.

Condiments.—Condiments can be regarded from two distinct and different points of view, viz.: one, hygienic, and the other gastronomic. Salt is an excellent addition to our food, and in some parts of the world where it is scarce, as in Central Africa, the saying for a rich man is:—“He eats salt with his food.” Animals will often go long journeys in search of salt; and the buffalo made periodic excursions to the “salt-licks.” Man is rarely deprived of salt; but animals frequently are from ignorance or neglect of the subject. I have read somewhere (though I cannot lay my hand upon the statement), that the hamster which lays up large stores of provisions for winter, gathers amongst the rest some leaves of the gentian and the century, to give flavour to its food; the bitters no doubt acting upon the appetite. Certain agents, as carminatives, have a good effect as reducing flatulence, and in India some birds are said to pick fruits and seeds containing aromatic oils to meet some craving; just as the Hindoo adds curry to his rice. When the dietary consists largely of vegetables, carminatives are speedily indicated, particularly pepper. And vegetarians find that carminatives are essential, as Dr. Kitchener says, to comfort.

Mustard is mainly eaten with meat; except when added to a salad. Pepper, the Romans valued so highly that they exchanged gold for it, weight for weight. Indeed so much pepper formed part of the ransom of Rome when besieged by Alaric. Cayenne pepper is still more potent and pungent, and has to be used sparingly. Of old it was common for persons suffering from flatulence, to make the pills of Cayenne pepper and bread crumbs, and swallow them at once. Ginger is another excellent condiment, either added to a curry, or eaten in powder as is customary with melon. Essences have the same effect in relieving flatulence.

The right use of condiments and their true proportion and combination has formed a great part of the cook's art in all times. It formed what is called a “relish,” and the painter would not paint

eating a red herring with his frugal meal of potatoes—"to give them a relish." So an anchovy is added to a beefsteak.

"The spirit of each dish and ZEST of all

"Is what ingenious cooks the *Relish* call."

And the condiments should be added in the act of cooking. If pepper be wanting in a beefsteak-pie or pudding, to add it after serving is a poor substitute for the real article. Some persons prefer highly seasoned dishes; and as a broad rule it may be said the taste for condiments grows with years. Children rarely care for highly seasoned dishes. Also as the palate flags, condiments and sauces are more in request; and this is specially true of gourmands in hot climates. The curry of the East Indies finds its correlative in "the pepper pot" of the West Indies. Pavy says of them—"Through their aromatic and pungent qualities they also assist digestion; the *modus operandi* being by promoting the flow of secretion and increasing the muscular activity of the alimentary canal. In some cases they may be further useful by serving to correct injurious properties that may belong to an article of food." They are of various kinds. Pickles are eaten with cold meat, especially when fat. Some are added to other food as a flavouring agent, as the onion, or garlic. Many as vanilla, lemon, almond, ginger, and nutmeg are used with milk puddings. Not only do they gratify the palate, but they serve a useful purpose otherwise.

Sauces.—Sauces are combinations of condiments, more or less complex. Some, as anchovy sauce, are used to add to or bring out the flavour where this is latent or little, as in boiled white fish. Some sauces are made of various ingredients, as with roast duck for instance, it is common to make a sauce by stewing the giblets, chopped fine, and mixed with the gravy in the dripping pan, and to this some finely chopped chabot is added. Such is a savory sauce. To some sauces wine is regarded as a good addition. *De gustibus non disputandum*, but a little plain gravy is nicer with meat and vegetables than an elaborately made sauce including port wine, - to my mind! Sweet sauces are used to puddings; and ruin sauces was once in great favour.

Then there are prepared sauces. Catsup is an old favourite. So is Indian soy. Many are named are the sauces and relishes now manufactured and sold. Harvey's was one of the first. As an all-round sauce Tarrant Pease's Worcester Sauce, "made from an old recipe for a nobleman's kitchen," is as good as any. It is said its peculiar

flavour is due to a certain admixture of asafetida; the exact amount constituting the secret. Of old, the flavouring agent was added to the ragout; but now various sauces are placed upon the table, and each guest flavours for himself. There is no broad rule like that of mustard with beef and pork; while it is never eaten with mutton and scarcely with bacon or ham. No one dreams of eating mustard with boiled chicken, or forgetting it with roast goose.

Condiments are curiously omitted from the dietary of invalids. There is a general impression that the plainer the food for them the better. But this leads to a terrible monotony in the dietary. Salt is often forgotten in the meat broths of the acutely sick; and a little Worcester, or allied sauce often gives a zest to a palate which has been palled by plain beef tea. If condiments do not possess a true food-value, they are not without a value of their own; and are not matters to be neglected in dietetics. Of chloride of sodium or common salt, its utility is easily recognized; though its *role* in the animal economy is not quite as clear as could be wished. Boussingault made experiments upon oxen, by giving salt to some, while withholding it from others, which told unmistakably of its value. As to the food value of vinegar, it is not easy to find anything positive. No doubt it is changed and oxidised in the body; and that it is agreeable to the palate as an addition to many articles of food; but beyond that it has a further utility. "It seems not improbable that as vinegar powerfully excites the secretion of the salivary glands, it exerts a similar influence upon the stomach, augmenting the flow of gastric juice, and thereby increasing the digestive power of the organs." (Stille.) Of its evil effects when taken to excess—as is done by some persons for the reduction of obesity—there is no question.

Condiments are agreeable to the palate, and, in moderation, good for the digestive organs.

CHAPTER V.

BEVERAGES.

Water.—As well as food, animals require drink; indeed the necessity for the latter is the more imperative of the two. Our bodies largely consist of water; and our secretions daily entail a considerable quantity of fluid; while for excretion a certain bulk of water is essential. Our tissues indeed, are bathed in fluids which at once bring nutriment, and carry away waste matters. The good effects of watering places are largely due to the increase in the bulk of fluid taken there, irrespective of any special quality of the water. Especially is this seen in the case of ladies, who, as a rule, habitually take too small quantities of fluids—for obvious social reasons. Further there is an impression abroad, that it is undesirable to dilute too freely the gastric juice. On this matter Pavy says, "Whether, as the result of the influence of this notion upon the public mind, or not, mischief is frequently occasioned, especially among the higher ranks of society, by a too limited consumption of fluids. Instead of taking a draught of some innocent and simple beverage, it is at many tables the fashion to sip fluid—and this a more or less strongly alcoholic one—only from the wineglass. It is a mistaken notion to think that when we drink with a meal, we are diluting the gastric juice. The act of secretion is excited by the arrival of the meal in the stomach, and the gastric juice is not there at the time of ingestion. It happens, indeed, that the absorption of fluid takes place with great activity, and the liquid which is drunk during a meal, becoming absorbed, may be looked upon as proving advantageous by afterwards contributing to yield the gastric juice which is required." So much for the dilution of the gastric juice. Then there is the question of excretion. Urates, especially, are sparingly soluble, and require a certain amount of dilution for their removal. Country ladies certainly drink far too little fluid, and more water is unquestionably desirable in their case; and if the water contain some saline matter it will be all the better for it. But whether alkaline waters should be drunk at meals or not, is a matter on which opinions may differ. The digestion in the stomach goes on in an acid

medium; and too much alkali certainly seems contra-indicated. On the other hand a draught of such water half an hour before a meal is certainly desirable. To provide an alkaline bath for the system is an excellent practice, especially first thing in the morning, after getting out of bed. But it is also well to take a second draught before a late dinner. That is one aspect of the subject.

There is, however, another. Sidney Ringer, in discussing alkalis, speaks as follows: "We wish to draw attention in this paper to one important property of alkalis—namely, their power to increase the secretion of the gastric juice—itsself an *acid* secretion. We venture to think many facts warrant the following generalization: that alkalis applied to the orifices of glands with acid secretions increase their secreting power; while alkalies applied in a corresponding way to glands with alkaline secretions lessen or check the secretion." It is good, then, to take alkaline fluids before a meal in order to increase the flow of gastric juice as well as to wash away waste matters from the tissues.

How far draughts of iced water at meals are injurious, either to the stomach itself, or to the digestive process, is a mooted point. Iced fluids have ever been sought for in hot climates, and are certainly agreeable to the palate. About that there is no doubt or question. The Austrians love iced beer to their meals; nor do they suffer from indigestion therefrom; especially the female element. How far the dyspepsia of Americans is due to their free use of iced waters, is a matter on which some persons are insistent; but I am not in a position to give an authoritative opinion on the subject. There are certainly other factors in operation in their case.

The amount of fluid required daily, is stated to be about three pints, to meet the water-loss by the kidneys, the skin, and the lungs. The gastric juice requires fluid. The bile secreted, the pancreatic flow, and the saliva all involve a considerable bulk of water. Beyond the matter of dilution of secretions and flushing of excretions, there are the relations of water to the body temperature. The cooling effect of water passing from the fluid form to that of vapor is the means of keeping down the body temperature, which otherwise would rise to a point incompatible, not with capacity for exertion, but with life itself. Dogs which do not perspire profusely, live in hot climates. Where there is great exhalation of water, and the sun is active; whether it be in the tropics or the heat of midsummer in the north. The workers in founderies, and iron factories, and in the mines measure their drink by the gallon. But taking persons generally,

life, the amount of fluid taken depends very much upon what it is; whether it is palatable or otherwise,—and whether or not it be charged with alcohol.

Fluids are taken with our meals, and are essential to digestion, and the secretions involved in the digestive act. Beyond that, most fluids chosen for such purposes possess a certain stimulant action. Alcoholic beverages were once the drink of all classes. Now tea, coffee and cocoa have largely taken their place. But the actions of these various stimulant beverages—as stimulants—will engage our attention in the next chapter.

Beverages as food-accessories.—As to the effects of beverages taken with meals upon the digestive processes, Sir William Roberts, F. R. S., of Manchester, has recently made extensive observations. In his address on Therapeutics at the Cardiff meeting of the British Medical Association in July, 1885, he gave the following conclusions, as the result of his enquiry.

“In studying the influence of our food-accessories on digestion, it is necessary to distinguish sharply between their action on the chemical processes, and their action on glandular and muscular activity. These two actions are quite distinct, and generally opposed to each other; for while all the food-accessories were found to exercise a more or less retarding influence on the speed of the chemical process, some, if not all of them, exercise a stimulating influence on the glands which secrete the digestive juices, and on the muscular contractions of the stomach. It is also necessary to distinguish between the effects of the food-accessories on salivary digestion, and their effects on peptic digestion, inasmuch as wide divergencies were found to exist in this respect. The distilled spirits—brandy, whiskey, and gin,—were found to have but a trifling retarding effect on the digestive processes, whether salivary or peptic, in the proportions in which they are commonly used dietetically. Their obstructive effects only become apparent when used in quantities which approach intemperance. Taking this in conjunction with the stimulating action which they exercise on the glands which secrete the digestive juices, and on the muscular activity of the stomach, their effect in these moderate dietetic proportions must be regarded as distinctly promotive of digestion.

“Wine and malt liquors exhibited an action differing considerably from that of ardent spirits. Wines were found to be largely inimical to salivary digestion. Even very small quantities of cherry, claret, hock, or champagne, inhibited the action of saliva on

starch to a very high degree. This is due to the acidity which all wines possess. When this acidity was neutralized by the addition of an alkali, the inhibitory effects of wine on salivary digestion was entirely removed. It is a common practice to mix wines, especially sherry, claret, and hock, with soda, or other mineral, effervescent table waters. These waters all contain a small amount of alkaline carbonate, and it was found that, when wines were so mixed, they ceased to embarrass the salivary action. This practice may therefore be looked on as highly commendable by the great persons of weak digestion.

On peptic digestion, wines exhibited a retarding effect altogether out of proportion to the alcohol contained in them. Both the stronger and the lighter wines, except in very moderate proportions, checked the speed of the peptic digestion. In the customary dietetic use of wines with meals, there is, probably, a double action; on the one hand, a stimulating action on the secretion of gastric juice, and on the muscular contractions of the stomach; and on the other hand, a retarding effect on the speed of the chemical process. In the case of persons of weak digestion, wines should be taken sparingly, and the quantity so adjusted as to bring out their stimulating action without provoking the retarding effects which follow their more liberal use. Champagne was found to have a distinctly less retarding power than an equal volume of claret or hock. This I judged to be solely due to the mechanical effects of the effervescence, and liberation of gas, whereby a more efficient stirring up of the digesting mass would be effected. Effervescent wines, therefore, other things being equal, favour the speed of peptic digestion more than still wines.

The effects of tea, coffee, and cocoa exhibited some interesting diversity. It was found that tea had an inhibitory effect on salivary digestion; even in very minute proportions, it completely paralysed the action of saliva. On the other hand, coffee and cocoa had only a slight effect on salivary digestion. The inhibitory action of tea on saliva was found to be due to the large quantity of tannin contained in the tea-leaf. Some persons have supposed that by drinking tea for a very brief period—two or three minutes—the passage of tannin into the beverage could be avoided. This, however, is not the case. Tannin is one of the most soluble substances known, and is very soluble in sugar in hot water. One gentleman, of my acquaintance, in his terror of tannin, was in the habit of tearing the green tea leaves on a paper filter, and simply pouring on them a small quantity

In this way he thought to evade the presence of tannin in his tea. But, if you try the experiment, and allow the product, as it runs through the filter, to fall into a solution of perchloride of iron, you will find that an intense inky-black coloration is produced, showing that tannin has come through in abundance. You can no more have tea without tannin, than you can have wine without alcohol; and I found, experimentally, that tea infused for two minutes had almost exactly the same inhibitory effect on digestion, as tea infused for twenty or thirty minutes. If you wish to mitigate the effects of tea on salivary digestion, you should direct the patient not to sip the beverage with the meal, but to eat first and drink afterwards. In this way time is given to the saliva to perform its functions unhindered. Another device is to introduce a pinch of carbonate of soda into the teapot; this removes the deterrent effect of tea on salivary digestion; it is a practice occasionally followed in some households, under the idea that soda helps to extract the virtue of the tea-leaves. It was found that the addition of so small a proportion as one per cent. of the weight of the dry tea, greatly mitigated its injurious effect on starch-digestion, and that twice this quantity (two per cent.) almost entirely removed it. This latter proportion corresponds roughly to ten grains of bicarbonate of soda to an ounce of tea-leaf.

"The effects of tea, coffee, and cocoa on peptic digestion were found to be as nearly as possible alike for infusion of equal strength. All three exercised a retarding effect when their proportion in the digesting mixture rose above twenty per cent. These beverages should, therefore, be taken very moderately by persons of weak digestion. The good reputation of cocoa, in regard to digestion, seems to be wholly due to the fact that it is used in weaker infusions than tea or coffee. The directions for the preparation of this beverage printed on the packets of cocoa sold in the shops, indicate a strength of about two per cent.; whereas a medium tea is usually made of a strength of four or five per cent; and a medium coffee of a strength of five to seven per cent. The strong coffee which it is customary to hand around after dinner must have a powerful retarding effect on gastric digestion; and although this practice may be salutary to robust eaters, it is not to be recommended to those of feeble peptic power."

I have preferred to make this long quotation in order to let Dr. Roberts speak for himself, as any summary might give rise to some misconceptions. Whether his conclusions will be entirely confirmed by other observations or not, we may feel pretty confident about the

man's facts. The well-known effect of the tea-drinking on the solid meal, seems to find its explanation in the fact that the turning digestion and especially salivary digestion of starch be thwarted and a quantity of undigested starch is left in the stomach, no wonder if pain and still more discomfort be performed. The fool should be eaten - say with a little beer or water - and then the tea drunk afterwards, by those who are ill. It is not that of the ostrich itself. The founder of the "great tea" is certainly not a benefactor of his species; nor is the late introduction of tea from China - a sterile country, the King of the tea country distinct advances - and from there on, however the Anglos have spread himself, been an unmixed good. Tea and campong cake of dough without leaven or yeast may be a meal of strength and easily prepared, but nothing can be said for it from a digestive point of view. No wonder that indigestion is as common among the stock-riders of Australia as among those of New Mexico. Tea may be drunk as a refreshing stimulant beverage; but it is not desirable as a food-accessory, especially where the food is largely farinaceous.

As to what Sir W. Roberts says about the good effects of the mixing of effervescent water with wine, and the action of the liberated carbonic acid gas upon the digesting food, it quite falls in with my own experience personal and professional, in the matter of malt liquors. I had begun to observe that for invalids and convalescents all malt liquors should be a good condition, but we changed water with carbonic acid; and that if "flat" or "dead" they did not do, whereas when a fierce attack of scarlet fever left me in a position to take further observations on myself, with the result that my previous views were confirmed. The liberated carbonic acid was a positive benefit while the acetic acid of "flat" malt liquor was as distinctly injurious. We also from this can understand the value of effervescent wines over still wines in the sick room. Probably a little potassium within an effervescent water is the best drink for a dyspeptic or invalid, and is commonly adopted by such persons.

From what has just been said about the beneficial action of the stomach upon digesting food, the more we learn, the more we are away with the idea that fermentation of food is a bad thing. A notorious fact being that fermentation is a good thing. The gas of fermentation is quite a different thing from the hydrogen from carbonic acid gas, and is a good thing. It is what underlies King Chambers' system of dieting, and is a good thing.

cludes his discussion on alcohol—"However little a man's purse allows him to drink, let it be good."

That wines and malt liquors have an action upon the stomach, which is pleasant and grateful, is a fact most people know for themselves. That they stimulate the gastric flow and increase the muscular activity of the stomach, is probably the underlying cause of the sense of well-being which accompanies their use. Consequently in atonic dyspepsia,—whether primary, or secondary to a general condition of debility—some combination of alcohol and carbonic acid in dilute form is indicated. But, of course, this is not to be made a rule absolutely to apply to all cases. There is a well recognized form of indigestion which is intractable to all medicine and regimen, except the avoidance of fluids. A dry dietary is a swift cure.

CHAPTER VI.

STIMULANTS.

THIS chapter covers a vexed ground—a debatable land, where views are in fierce conflict. Strong feelings have been, to say the least, and beef-teen, vegetable principles (as the vegetarians call them) and alcohol have assailants and supporters who too often do not give each other fair play; and use declamation instead of argument. Indeed, sometimes the declamation is in inverse proportion to the argument, consequently I cannot expect to please all for what will be written in this chapter; certainly, not the extremists in either wing.

Stimulants divide themselves into: (1), animal and vegetable substances of which nitrogen is the conspicuous factor (as it is of all explosive compounds); and, (2), alcohol.

1. Animal principles. Beef-teen may stand as the typical representative of stimulants derived from animal food at the present time, and in the present state of public opinion. Yet it is strange a food—as seems so universally assumed. On the very threshold of this chapter it becomes necessary to say a popular correction. When the relatives of a sick person tell one in all earnest, of the amount of beef they have converted into beef-teen, and how the patient has been sustained thereby, it is with a difficulty that the muscles of expression are prevented from throwing a look of scorn over the features. "There is no pain like the pain of a new word," according to Bagehot; and to tell these good people that they are simply talking nonsense—that in fact they do not understand what they are talking about, would be to inflict upon them more pain than they would probably arouse a feeling of distrust, rather than to do any good else in their minds. In fact, it would scarcely be worth the trouble in most households. Still, all the same, to give a sick person a stone when he asks for a loaf, is not a fool, it is a stimulant! Grateful and no doubt, the patient and the stomach, possessing stimulating qualities, has its value. But all the same, as regards food, it is a stone in a jackass in a lion's skin.

What can really be said for it?

It is a decided stimulant. When flesh is boiled its albumen is coagulated. That is a simple fact. Beef tea, as ordinarily prepared, contains no albumen. The fluid so obtained is rich in extractives and salines. There are present the phosphates of the flesh in solution. The extractives are nitrogenized bodies—kreatin and kreatinine—the first with the formula $C_4H_7N_3O_2$ and the latter $C_4H_9N_3O_3$; which means kreatin is a hydrated form of kreatinine are found in meat juice and muscle extracts; but they belong rather to muscle as derivatives as histolytic rather than histogenetic matters. In other words, which can be understood of the people, they are not material of which muscle can be built up; but rather stand as effete muscular matter in the road to urea. Albumen is material of which and from which tissue can be formed. It is like the bricks which are going to form the walls of a new building. Kreatin and kreatinine are rather the old bricks and mortar of an old building, (which have served their day) coming down to be carted away as rubbish. The liquor sanguinis brings the one and removes the other; just as contractors' carts may be seen together when buildings are being pulled down, and rebuilt, the one laden with new bricks for new walls, the other filled with the *débris* of old walls. Now I trust the matter is unmistakable. These extractives are not food-material for tissue; but stimulants. "Kreatin and kreatinine are products of the vital process and constituents of the flesh of all vertebrata hitherto examined. Human flesh is particularly rich in kreatin." (Liebig.) It is easy from this to understand why savages ate the bodies of their enemies believing human flesh to endow them with courage. No doubt it did stimulate them and make them feel warlike; just as Dominic Sampson felt exhilarated by some of the contents of the gipsy's cauldron, saying, "Verily I feel mighty elevated and afraid of no evil which may befall me!" a very different mental condition from that in which he was when Meg Merrilies accosted him at the Kaum of Dornelough. Of the stimulant effects of meat extractives, of the utility of the meat salts in beef tea, there is and can be no question. But it is destitute (practically) of matter which can ever form tissue; and is equally without value as fuel-food.* As a food it is no impostor, neither more nor less. As some readers may feel indignant at this state-

*Certainly some oxidation goes on at the secondary products of retrograde tissue metamorphosis pass on to ureic acid and urea, but it is infinitesimal.

ment, and fling out that the imposture lies rather with the nation, let Sir William Roberts once more be heard:

"Beef tea and its congeners take rank as restoratives and stimulants, rather than as nutrients. They contain nonalbuminous matter in solution, and the small amount of gelatine contained in them cannot be of much account. There is a widespread misapprehension among the public in regard to the nutritive value of beef-tea. The notion prevails that the nourishing qualities of the meat pass into the decoction, and that the dry, hard remnant of meat fibre which remains undissolved is exhausted of its nutrient properties; and this latter is often given to the cat or dog, or even as I have known, thrown away as useless rubbish into the midden. A deplorable amount of waste arises from the prevalence of this erroneous notion in the households of many who can ill afford it. The proteid matter of meat is quite insoluble in boiling water, or in water heated above 160° Fahr. The ingredients which pass into solution are the soluble extractives and salts of the meat, and nothing more except a very trifling amount of gelatine. The meat remnants, on the other hand, contain the actual nutriment of the meat; and if this be beaten to a paste with a spoon, or pounded in a mortar, and duly flavoured with salt and other condiments, it constitutes not only a highly nourishing and agreeable, but also an exceedingly digestible form of food." Yes, Dr. Roberts, and there is another "deplorable waste" involved in this error and that is a "deplorable waste" of human life; hundreds of our fellow-creatures are daily dropping into their graves, unfed, unsuccoured, because beef tea is given as a food—given for what it is not!

Dogs may starve on lixiviated meat, and even turn from it with loathing; for this meat lacks the salts and extractives. But all the same, human beings cannot sustain life on salts and extractives without the other constituents of flesh. So there is another error, even to pieces, and a cherished conviction destroyed! Liebig himself states only too well that meat broths have no true food value, and that cold-drawn infusions of meat. Such infusions are food only in so far as they derive considerable value from the amount of albuminoid matter they contain.

I can see, sad dry readers, having been driven past the state of incredulity by the hard logic of facts, wringing their hands in anguish over the thought of departed relatives who have been literally starved to death on beef tea. We can feel, but we cannot help them. "Let the dead past bury its dead." But, oh, how we may sorrow for them! Let us be wiser in the future, and let

The mistaken views about the nutritive value of beef tea have been murderous. All the bloodshed caused by the warlike ambition of Napoleon is as nothing compared to the myriads of persons who have sunk into their graves from a misplaced confidence in the food value of beef tea. As a food, it is but as the mirage of water seen by the thirsty traveller in the desert: there is no real water. So with beef tea, it is not a food. A traveller once requested some chicken broth at an inn, when brought its quality was very indifferent. Complaining that there was nothing of the chicken about it, the waiter assured him he was mistaken. "It was made from water from a pool in which a chicken had paddled!" In so far as the fluid was "chicken broth" is beef tea a food!

What says King Chambers: "Make the cook understand that the virtue of beef tea is to contain all the contents and flavours of lean beef in a liquid form; and that its vices are to be sticky and strong and to set like a hard jelly when cold. When she understands this, let her take half a pound of fresh killed beef for every pint of beef tea required, and remove all fat, sinew, veins and bone.

Let it be cut into pieces under half an inch square and soak for twelve hours in one third of the water. Let it then be taken out and simmered for two hours in the remaining two-thirds of the water, the quantity lost by evaporation being replaced from time to time. The boiling liquor is then to be poured on the cold liquor in which the meat was soaked. The solid meat is to be dried, pounded in a mortar, freed from all stringy parts and mixed with the rest."

This last matter is the important matter, to restore the meat fibrine to the fluid. That it is which gives food value to the meat infusion. At the Victoria Park Hospital we have recently had the beef tea prepared on this plan which Dr. Chambers calls "whole beef tea" and though at first the patients do not like it so well, it is of course much better for them. The nitrogenized principles derived from meat are a descending series which commence with kreatin and kreatinine and pass into tyrosin and leucin by an oxidation towards the final products, uric acid and urea. As said before, all explosive bodies contain nitrogen, and these nitrogenized bodies do not supply force in themselves, but they liberate stored force, converting it into "kinetic energy." They are like the fuming matter of a cartridge which does not itself propel the bullet, but which explodes the gunpowder whose decomposition is the real propelling power. Consequently to give them alone to a sick person is to exhaust the stored up material—the "potential energy" whose oxidation produces

force. They should properly be accompanied by some so-called carbohydrate. To give them alone is as illogical and erroneous as it would be to thrust a bullet down a gun barrel and put on a cap, omitting the powder. These explosive nitrogenized bodies play an interesting and important part in the economy. A carnivorous animal could never catch an antelope in a clear run, but it can make a supreme effort and pounce upon its herbivorous prey. And this it can do by the nitrogenized elements derived from its food—flesh. Greyhounds are fed on meat, because they require to put forth immense energy for a brief period. Sporting dogs who cover long distances in a day—but at a less rate of speed, are fed on broth and oatmeal. And the same occurs in the feeding of horses. Trainers will tell you that the hunter and the draught horse require to be fed differently. The hunter is bred and fed for speed and carrying power; the draught horse for bulk and strength. In the hunter is wanted rapid liberation of energy within a comparatively short space of time; in the draught horse is wanted a more gradual liberation of energy and for a longer period. To bring out these qualities each strain must be fed appropriately. The hunter is fed on a concentrated and stimulating form of food—chiefly on the heaviest and most expensive oats—which if I may so express it, is the “beef” of the vegetable feeders; and unless he is so fed, he will not perform satisfactorily in the hunting-field. The draught horse is fed on a lower and less stimulating diet—on Indian corn and chopped hay—food which tends to increase bulk and weight, (Sir Wm. Roberts’ “Lectures on Dietetics and Dyspepsia”). Not only is this so, but when the ground is heavy the hunter gets some beans or peas extra—legumes which are specially rich in nitrogenized principles; while in ploughing time, especially in spring when the hardest agricultural work is performed, the farmer adds a liberal amount of clover hay to the ordinary food of his horses.

This rule applies to races. The conquering Anglo-Saxon, the master and too often the exterminator of aborigines whose lands he covets, is a meat-eating man (*patre et filii*). His carnivorous propensities are linked in some subtle manner with his conquering power.

B. Vegetable principles.—Closely allied in chemical composition as well as in physiological action to the animal principles, stand a certain group of vegetable principles. The green tea of China and to a less extent, maté, or Paraguayan tea, are brewed in infusions with hot water and drunk as food-necessaries. Sugar, which supplies at once fluid for digestive purposes, and a certain amount of

effect which has doubtless been the cause of their adoption. Theine with a formula $C_7H_8N_2O_2$ is the type of these principles. They are a considerable group, commencing with theine, caffeine and theobromine, and ending with brucine and strychnine,—a mighty tonic of toxic properties. Strychnine, with a formula of $C_{21}H_{22}N_2O_4$, is constantly used as a tonic of rapid action so as to come under the heading of diffusible stimulants. When the Resident Medical Officer of the Leeds Public Dispensary, where I could command any amount of drugs of every kind,—but where meat and drink were beyond my reach, I found in many cases of debility and convalescence from acute disease, that ammonia and nux vomica in combination, supplied the place of alcohol very fairly.

The chemical composition of these vegetable principles, and the similarity of their action was a matter which did not escape the observation of Liebig, who wrote—"Theine is related, in composition, to no organic nitrogenized base more closely than to kreatinine, that remarkable compound produced in the vital process, and occurring in the muscular system of animals; and to glycocoll, which we may suppose to exist in gelatine, coupled with another compound, as may be seen by the following formula:

Theine,	$C_7N_2H_8O_2$
Kreatinine,	$C_4N_4H_8O_2$
Glycocoll (anhydrous),	$C_2N_2H_2O_6$
Kreatin,	$C_4N_4H_7O_2$
Theobromine, (in Cacao),	$C_7N_2H_8O_2$

There are no drinks which in their complexity and in the nature of certain constituents have more resemblance with soup than tea and coffee; and it is very probable that the use of them as a part of food depends on the exciting and vivifying action which they have in common with soup.

There are some discrepancies betwixt these and more modern formulae, which may or may not depend upon the new notation; but which are of comparative unimportance as compared with the main fact of the strong similarity and chemical resemblance which exists between these substances. Infusions of meat and infusions of certain vegetable principles alike contain stimulant matters, whose properties rest upon the large proportion of nitrogen they in common possess.

These vegetable principles are used largely by races who do not take alcohol. The Arab and the Turk, forbidden by their religions

to taste alcohol, have found a substitute in coffee; while the numerous Chinaman has cultivated the Tea-plant. The total abstinence of Anglo-Saxon descent has adopted the drinks of these races. "It is true that thousands have lived without a knowledge of tea or coffee, and daily experience teaches us that under certain circumstances, they may be dispensed with without disadvantage to the merely animal vital functions. But it is an error, certainly to conclude from this that they may be altogether dispensed with, in reference to their effects; and it is a question whether, if we had not tea and coffee, the popular instinct would not seek for and discover the means of replacing them. Science, which accuses us of so much in this respect, will have in the first place, to ascertain whether it depends on sensual and sinful inclinations merely, that every people of the globe has appropriated some such means of acting on the nervous life, - from the shore of the Pacific, where the Indian retires from life for a few days, in order to enjoy the bliss of intoxication with koka, to the Arctic regions, where Kamschatkales and Koriakes prepare an intoxicating beverage from a poisonous mushroom." (Liebig.)

Not only is there a widespread instinct for "these means of acting on the nervous life," as Liebig happily phrases it; but the consequence of excessive indulgence is seen in the various nervous affections which we notoriously associate therewith.

The "tea-drinker's heart" is a well-recognized malady. People, especially of the female sex, who drink largely of tea, and still more when they do not eat a sufficient quantity of food, suffer from many nervous troubles as well as palpitation and neuralgia; and present all the phenomena of nerve-exhaustion. The nurse often drinks tea to sustain her when there is no appetite for food, until utter exhaustion is the untoward result; and a state bordering on the delirium tremens of alcoholic excess is revealed. The "tea-drunkard" is a well-known patient at all hospitals; and is not unknown in private practice. "The cup which cheers but does not inebriate" is not without a toxic influence, when taken to excess.

While tea is in almost universal use with the Anglo-Saxon, coffee is equally in vogue with the Germans. They have the same effect on a great extent. Coffee, however, has a distinct effect upon the heart; and caffeine with a formula of $C_8H_8N_4O_2$ is used as a cardiacine in cases of cardiac debility. Vogel has advocated, and has strong coffee with cream as a tonic and food in the debility which attends upon acute disease in children. No such use of coffee

yet found an advocate. Tea contains an aromatic oil as well as theine, and its effects upon the nervous system—for good or evil—may to some extent be due to this rather than the theine.

So much then for those stimulant bodies derived from the animal and vegetable worlds alike, whose properties depend upon the nitrogen contained in them.

Alcohol.—If the matter of beef tea is difficult of discussion in consequence of ignorance and prejudice, how much more difficult is it to handle alcohol—a matter bristling with difficulties, when the pure white light of truth can scarcely be seen clear of the coloured rays around it?

Alcohol has a formula C_2H_6O (better known, however, by the old formula $C_4H_{10}O_2$), and is a readily combustible substance—out of the body. A very light, volatile body it is, highly inflammable. All matters which are oxidizable in the body are “foods” (Hermann). Liebig held the opinion that “Alcohol stands only second to fat as a respiratory material” while adding “Alcohol and alcoholic drinks are from their price most costly materials of respiration. The same effect could be produced in the body by means of saccharine and farinaceous articles of food at one-fourth or one-fifth of the cost.” A fierce war, embittered by the presence of sentiment, has raged around the matter of the food-value of alcohol. Readily combustible outside the body, some have denied its combustibility within the body. Fifty years ago, the most advanced chemists believed in the “spontaneous combustion” of the bodies of drunkards. In that they were certainly mistaken. Then came the recoil of opinion. Total abstiners are ranked in battle array on the matter with other observers. These last, and their names are both numerous and weighty, hold that alcohol is largely burnt in the body by oxidation and is therefore a “fuel food.” Personally, after very considerable attention to the subject, I must say that I am among those who hold “that the chief portion of the alcohol injected undergoes consumption in the body.”

But the question of “alcohol as a food” can never be separated or divorced from that of “alcohol as a stimulant,” as a force-liberator. Liebig recognized this aspect of the subject, and wrote as follows:

“The use of spirits is not the cause, but an effect of poverty. It is an exception from the rule when a well fed man becomes a spirit drinker. On the other hand, when the laborer earns by his work less than is required to provide the amount of food which is indis-

penable in order to restore fully his working power, an inexorable law or necessity compels him to have recourse to work. He must work; but in consequence of insufficient food, a portion of his working power is daily wasting. Sports, and their action on the nerves, enable him to make up the deficit, *power at the expense of his body*, to consume to-day that quantity which ought naturally to have been employed a day later. He draws, so to speak, a bill on his health, which must be always renewed, because, for want of means, he cannot take it up; he consumes his capital instead of his interest; and the result is the inevitable bankruptcy of his body."

This matter of "physiological bankruptcy" made a deep impression upon me at the very threshold of my entrance upon medical practice, and has been my guide to the use of a careful stimulants in the treatment of disease. If much alcohol be given without other readily oxidizable food (while furnishing a certain amount of respiratory food in itself) its effect is to consume the body stores; in other words to produce "physiological bankruptcy." Especially is this matter to be borne in mind when there is danger of the system sinking from exhaustion. There is a deep-rooted fear among me that many a case of acute disease (and especially in fevers, where a certain course of so many days has been sent into the grave over-dug by nature" as H. C. Wood mildly puts it) by over-stimulation by "physiological bankruptcy." Other hydro-carbaceous food as well would have been preferable and the demands of energy evoked by doses of alcohol, were as disastrous as the excesses of wealth manifested by a reckless young square who mortgages his means for them by mortgages on his property: each day pay rendering him distinctly poorer.

This great cardinal fact that stimulants are "neutral agents" must never be forgotten in practice. If force be not available to meet some emergency then alcohol (supplied in some form or other) as well as liberating other force may be required, and the after-effect must not be left out of the calculation. In the case of conditions of acute exhaustion free resort to neutral stimulants is at hand; and may be, too, to a certain extent, in the case of chronic quantities to tide over the period of exhaustion. The mode of its operation must be borne in mind, and the fact that it is a "neutral agent."

As to the "food value" of alcohol, it is not to be compared to its action on the system, and is a very small part of its costliness. Further, it is not a food.

lent of a fair meal in alcohol, he would be dead drunk. If he took it comparatively quickly—as when men, for wagers, drink so much spirits in a limited time—the strong probability is, he would in a very short time sink into coma; which ere long would deepen into death. Its “food-value” in health is such that its consideration need not engage our attention for long.

But in ill health and especially in acute disease the question of its “food-value” as well as its value as a “stimulant,” may well engage our attention at some length. As well as a force-liberator it has some value as a “fuel-food.” Indeed, at times, dilute alcohol is all that the patient can take, or the stomach retain. Alcohol requires no digestion; let that fact be kept in view. By its ready diffusibility it quickly passes by osmosis from the stomach into the blood. When the digestive powers are seriously enfeebled this matter becomes very important. Alcohol provides some force, and borrows some from the system. How far can the borrowing go on? That becomes the question. It may be possible to provide the alcohol in unlimited quantities; but is the store of body force illimitable? That is where the shoe pinches. Whenever possible or practicable the alcohol should be accompanied by some soluble carbo-hydrates, which would supply the place of the borrowed force. Say with some Malt Extract diluted; that would be a capital combination; or in milk, excellent. Or with lemonade, home made, containing a distinct amount of soluble sugar. The neglect of this matter is fraught with disaster. This will be further considered when the treatment of exhaustion comes to be discussed.

The effect of alcohol is to produce increased action in the heart, while the carotid arteries are dilated, and so the brain is flushed with blood. This flooding of the posterior cerebral lobes produces a sense of well-being, *i.e.*, the patient feels better for the time; no doubt, too, the alcohol borne along in the blood acts directly upon the cerebral cells. Probably a like continued effect of increased blood supply and direct stimulation goes on in all parts of the nervous system—the organic or sympathetic as well as the cerebro-spinal. But this effect is temporary and evanescent. Such stimulation carries with it the after effect of exhaustion. The heart flags, the carotids contract below their normal calibre and the brain becomes bloodless. If this lead to sleep, as often happens, then the alcohol may have been advantageous in every way. But the temporary time of stimulation ought to be utilized to assimilate some easily digestible food. Otherwise, it may have been useless; ay,

worse than useless. King Chambers says well of the alcoholic drink be limited to that quantity "which will excite the appetite." He is at the time speaking of its use in fevers. But the principle is a good one to hold to under all circumstances.

Alcohol is a two-edged tool, not to be used without discretion; and, like any other tool, all the more skilfully used the more danger with it. Sometimes it is withheld when the call for it is urgent. At other times it is pushed recklessly, and alcoholic gastritis comes up; the stomach rejecting it quickly, while the tongue becomes irritable, and the bowels filled with flatus. This condition is produced when alcohol, commonly in the form of brandy, is given in considerable quantities, insufficiently diluted, and at frequent intervals. Here Nature is stepping in to limit that stimulation, which is threatening the existence of the organism when exhaustion is close at hand, and "the beating of the wings of the Angel of Death are distinctly audible overhead."

This condition of exhaustion is specially liable to come on when alcohol, alternately with beef tea, is alone given to the patient. Beef tea which (as ordinarily made) is no food; alcohol which is a poison, carrying with it a heavy borrowing power! No wonder the system is in danger of sinking from exhaustion. All that has been said here applies to acute conditions, as fevers, and also to conditions not exactly acute, as in dropsy with a failing heart; or in acute conditions implanted upon a chronic state, as in an acute bronchial attack in a person the subject of chronic bronchitis. Here we wish to sustain the powers, and for that end we use alcohol. But we must so use it in combination with soluble carbo-hydrates that we do not make it a means of sinking the ship. If it be given, along with, or alternately with, a meat broth containing some beef-tea, or milk, or diluted malt extract, then the power may be sustained, the strength be kept up or maintained. But when alcohol is administered, experience has told us in unmistakable manner, the end is not far off.

We are told "to use the good things of this world's enough of them;" and to do this with alcohol is a test of judgment. If one comes the requisite knowledge, then the judgment is a true judgment, and might. In times of acute depression, alcohol is of great value. But then the end must be kept well in view. A course must not be indefinite; must not be continued for a long proportion as it can be combined with other food, and must not be continued. But when it alone can be taken, the end is not far off.

measurable distance. So much for alcohol as at once a food and a stimulant—a "force liberator."

When the subject of malt liquors comes before us, then no one can dispute, or think of disputing the food value of the maltose contained therein. In the brewing process a certain quantity of maltose is broken up into alcohol and carbonic acid; while a distinct proportion remains unchanged. Consequently the well-fed navvy who prefers to do the 1st hour's work of his laborious day on a pint of good sound ale, has a hard solid argument on his side. Having expounded my own views at some length and as judiciously weighed the subject as in my power lies, some reference may be made to what other men have thought and written on this difficult and complex subject. Brinton in "Food and its Digestion" after going over the pros and cons, wrote—after giving the obvious drawbacks of alcoholic indulgence and how it blunts alike the mental edge and the delicacy of touch and movement:

"On the other hand, however, it is necessary to remember how often the whole question must practically be approached from a totally different aspect: how often alcohol constitutes not the single feather which distracts the sleepy savage, but the bed of down which restores the exhausted man. It may disturb a balance exquisitely adjusted; and yet, in the main, counterpoise a scale heavily laden with disadvantages. If alcohol exhilarates, imparts comfort and energy, counteracts fatigue, hunger and unrest, then it does in effect, increase the capacity for work of those who take it under such circumstances; and affords, in so far, a direct benefit and advantage."

Pavy says of it as a food: "From a review of the evidences as it at present stands, it may reasonably be inferred that there is sufficient before us to justify the conclusion that the main portion of the alcohol ingested, becomes destroyed within the system, and, if this be the case, it may fairly be assumed that the destruction is attended with oxidation and a corresponding liberation of force; unless, indeed, it should undergo metamorphosis into a principle to be temporarily retained, but nevertheless ultimately applied to force-production."

King Chambers sums up his consideration of alcohol so: "(A). Let it be taken never as a stimulant or preparative for work, but as a defence against the injury done by work, whether of mind or body. For example: it is best taken with the evening meal, or after toil.

(B). Let the increase in the desire for and power of digesting food be the guide and limit to the consumption of all alcoholic liquids.

(C). Let the forms be such as contain no fusel oil.

(D). Let all with an hereditary tendency to functional disease of the nervous system, to take them together, even though as yet they are not good to drink. And hereditary tendencies must be classed a priori with congenital weakness, which remarkably runs in families. "One person's an heirloom had best be kept to us into an age," says the old "tasting strong drink."

Indeed, there can be no question about the desirability of holding alcohol from children under any circumstances but those of dire necessity. It is an injury to health to subject to the stimulant action of alcohol. Children are in many respects like infants. And of these Sir William Roberts writes: "There are certain inferior races who appear to be altogether intolerant of alcohol. That it does not suit their type of nutrition, or they lack the secretion which is necessary to its beneficial use. The Indians of North America are said to be excited almost to madness by a small dose of alcohol, inasmuch that the Colonial authorities forbade, under heavy penalties, the giving or selling of alcoholic liquors to these tribes. The Ainu of Yezo, a subject race inhabiting the Northern island of Japan, appear to be wholly wanting in self-control in the use of alcoholic stimulants, for which they exhibit an irrepressible passion. Whenever they have the opportunity, both the men and women drink themselves to a state of insensibility. The defective reaction of these races towards alcohol may be compared to a peculiarly defective reaction in certain individuals and certain families among ourselves."*

On the other hand, just as strong men usually tolerate alcohol well, so apparently do the stronger races of mankind. King Chambers writes: "All nations that have led the van in the march of civilization have been addicted to drink, and, and addicted to drunkenness. The Jews, the Greeks, the Romans, the Germans, the Swedes, the Danes (not to mention the English) are noted for this; and are amply attested by their own native literature to have been distinguished above their contemporaries in this way. But even among some reactionary races, famous as we are for having been so, there

*Colon Kemmis, of Sidney, informs me that this is equally manifested by the Aborigines of Australia, who are well protected against alcohol as are the Bechuanalanders. The Crown

but they and their faiths are dying out, and the coloring they have given to civilization is even now fainter than that left by the robust races a thousand years before they were heard of." While Maudsley has pointed out that drunkards in society are like the waste-heaps of manufacturing industries—they are so many witnesses of the intellectual activity going on around them. The stronger the race, the greater their enjoyment of that exaltation which alcohol produces. As Burns says:—

John Barleycorn was a hero bold,
Of noble enterprise;
For if you do but taste his blood,
"T will make your courage rise,
"T will make a man forget his woe;
"T will heighten all his joy;
"T will make the widow's heart to sing,
Though the tear were in her eye."

This effect it is which constitutes the danger of alcohol; and if the delight in such excitement cannot be kept within strict bounds, the downward career of alcoholic excess ending in drunkenness and degradation, is apt to be set on foot.

In concluding his contribution to "The Alcohol Question," which was discussed in the "Contemporary Review," some few years ago, Dr. Sam. Wilks said:—"In judging of the use of alcohol by the community at large, we must be guided in the same way as we are by other habits of mankind. We see persons enjoying themselves in various ways, eating and drinking all kinds of food and beverages, occupying themselves with amusements of every description, and yet none of these would be allowed in Utopia. They get through the world, although indulging in certain habits, and declare themselves well—where, then, is the appeal against their procedure? If I, personally, am consulted as to the propriety of ordering alcohol in any individual case, if there be no experience to guide me, I am impeded by the principles I have enunciated. I believe alcohol soothes a worried nervous system, and by preventing wear and tear, actually supports the frame; but, discarding the notion of its stimulating properties, I denounce its use in delicate children and in women who feel 'low.' I also strongly prohibit its use in the early morning; in fact, those who then wish for it, have already imbibed too much. I always suspect people who require 'something' about eleven in the morning. Indeed, the man or woman

who has an acute consciousness of the hour of eleven—*“I am physically and morally lost.”*

No doubt there is much truth as well as force in the foregoing, and there are two matters connected with alcohol, well worthy of being in mind:—

1. Never have alcohol in the brain when it has work to do;
2. A little alcohol betwixt a man and a past trouble, is conceivable; but it is not well to put a little alcohol in front of a coming trouble.

I may fitly conclude this chapter by a quotation from the late Dr. Murchison, whose book on Fevers is so favourably known everywhere:—

“What then are the conditions of the animal economy in which alcohol may be of positive use? That there are such conditions I believe cannot be denied by anyone who has honestly studied the subject; but they are not the conditions of perfect health. It is especially when the circulation is weak or sluggish, that a daily allowance of alcohol may do good. Thus

“1. Alcohol is useful in the course of most acute diseases, when the organs of circulation begin to fail, as they are apt to do. A moderate quantity usually suffices. The large quantity, e.g., one or two bottles of brandy in twenty-four hours, still sometimes administered, may do harm by inducing congestion of various internal organs.

“2. In convalescence from acute diseases, or from other weakening ailments, when the circulation remains feeble and the temperature is often subnormal, alcohol is also useful in promoting the circulation and assisting digestion.

“3. In persons of advanced life the circulation is also often feeble, and a moderate allowance of alcohol often appears to be beneficial.”

“All other conditions of the system marked by weakness of the muscular wall of the heart, whether permanent or transient, are usually benefited by alcohol.”

Alcohol is “a good servant, but a bad master.”

The charge brought against the medical profession, that much alcoholic excess takes its origin in medical advice, is not to be preferred. It is undoubtedly often made the excuse for intemperance. I certainly think the greatest caution should be exercised in the case of neurotic women and their children, where there is a craving for alcohol, and the habit is most easily acquired. We must never forget its possible abuse.

CHAPTER VII.

FLUID FOOD.

THE type of "Fluid Food" is Milk—the food Nature provides for all young mammals. It is a complete and perfect food, containing fat (in an emulsionised form,) albumen, (in the form of casein,) carbo-hydrates, (as milk-sugar,) with salts, (phosphates and others,) all properly diluted. On this food life can be sustained for an indefinite period. In sickness, milk is really our sheet-anchor. But there is one matter to be borne in mind, and that is this—milk curdles as the first step in its digestion. The light curd usually readily re-dissolves in the alimentary canal; but sometimes it does not. Especially is this firm curdling liable to occur where there is acidity present in the digestive tract. The hard, firm curd is an irritant to the whole intestinal surface; and in conditions of ulceration, either tuberculous or that set up in enteric fever, hard pieces of curd may rupture, or perforate the weakened bowel. The pieces of curd may be seen in the stools. When such is the case the milk must be combined with means which will prevent this hard curd forming. At times it is well to stir some biscuit powder into the milk, which mechanically prevents too firm a curd; or where obvious acidity is present, some light carbonate of magnesia, which acts in two ways. First of all it chemically neutralizes the acidity; next the carbonic acid gas given off tends to disintegrate the curd and break it up. Milk is commonly given with lime water; but this is often too feebly alkaline, and magnesia or prepared chalk must be used. Usually as much as will lie on a sixpence (or its U. S. A. equivalent) to the half pint of milk is enough; or it may be given with a mineral water as Seltzer water, especially where the thirst is considerable. At other times it is well to combine it with such waters as Vichy, Vals, Marienbad; and in some conditions of lithiasis such combination may be freely drank.

At other times some malt extract well diluted may be added to the milk with advantage. This last is indicated in those cases where the patient feels the long fast betwixt going to bed and rising, or either betwixt the usual hours of bedtime and of rising; and

something is desirable in the small hours of the morning. A tablespoonful of malt extract to half a pint of warm milk in a hot water jug (with a lid) put on a piece of wood, or other non-conductor of heat, and covered with a tea cosy (the bigger and the better), will be found a pleasant, palatable, and nutritious drink. The tea cosy will retain the heat.

Then at other times milk is boiled with a little ground rice flavoured with cinnamon; and this is an excellent food when there is any looseness of the bowels.

Skim-milk is an excellent beverage in pyrexial states, where fresh milk is too heavy. Buttermilk with its lactic acid is a good beverage where there is much thirst and in diabetes.

Milk is sometimes too cloying and is preferred in the form of whey. Whey is a pleasant stimulant drink-food. It is made by adding one or two wineglassfuls of sherry to half a pint of hot milk; the curd is strained out by a sieve. It is not in fashion at present; but its day will come again. The milk-sugar in it gives it a certain food value.

Then there are certain pleasant drinks with a low food value, good in fevers, as apple-water, and tamarind water. They are pleasant to the taste. A favorite drink is lemonade. The rind should be thinly pared (to get out the essential oil) and the lemon be cut into slices, put into a jug and hot water be poured in. Then some sugar should be added or if this be apt to turn sour in the stomach some malt extract may be used instead of cane sugar. Such lemonade is grateful and nutritive; especially when made with malt extract or with Mellin's Food.*

Then as a "drink-food," oatmeal water is capital where persons are exposed to a high temperature as stokers, iron-workers and the like.

Barley water is a pleasant nutritive drink, and a capital vehicle for bitartrate of potash. Potus Imperialis sweetened is a capital beverage, good in a cold with a deposit of pink lithates.

Rice water is a drink in vogue in India. It is a nutritive drink, and thereby hangs a tale. When Clive was besieged in Arcot, provisions ran very short. There were a few Englishmen and a number of Sepoys. This is Macaulay's account of what happened.

It is invidious to mention names in connection with food, but Dr. Mellin has been such a pioneer, that his food may be spoken of as the type of a most useful class of foods and as such is spoken of frequently. For his food Mr. Mellin is unknown to me.

“The Sepoys came to Clive, not to complain of their scanty fare, but to propose that all the grain should be given to the Europeans, who required more nourishment than the natives of Asia. The thin gruel, they said, which was strained away from the rice would suffice for themselves. History contains no more touching instance of military fidelity, or of the influence of a commanding mind.” Literature contains no more striking instance of a man, accurate by habit, tripping egregiously. Those who understand the matter thoroughly assure me that the men who were satisfied with “the thin gruel” got much the best of the bargain.

Rice water may be flavoured by any vegetable juice as that of the currant, raspberry, apple, and mulberry. Where there is great thirst in pyrexia, such drinks supply all that is required. In phthisis however, it is desirable to increase the food value of the drink, and this can be done by the addition of a malt extract, or soluble malt preparation like Mellin's.

Rice which has been malted forms a capital addition to meat-broths, as the starch has largely been rendered soluble.

Dextro-saccharin, (?) or commercial grape sugar, prepared for the brewer, can also be utilised to give food value to drinks—whether meat-broths, or made with vegetable material, as apple-water.

Meat Broths.—Beef, veal, mutton, rabbit, and chicken, are the materials mainly employed for the preparation of meat tea. In the last chapter it is to be hoped that the brains have been knocked out of the superstition that beef tea, or any other meat tea, as ordinarily made, is a food. The addition of the albumen of the meat-fibre is required to give it a true food value. If that were done, the amount or bulk of water to the pound of meat might be largely increased. Then too, a certain amount of salt should be added. In invalid cookery there is a decided neglect of salt. Liebig advocated the use of meat extracts. He deplored the waste that went on in the large herds of cattle on the plains of South America and Australia, where thousands of oxen were annually slaughtered for their hides and tallow; and suggested the preparation of concentrated meat-extracts from the flesh. But Liebig himself was the first to recognize that such meat-extracts were a “means of invigoration,” rather than a food. There are present the pleasant salts of muscular tissue and the stimulant kreatin, which tempts the savage to eat his vanquished adversary—but nothing that is true food.

Beef, or meat teas, as ordinarily made, and Liebig's meat ex-

tracts stand on the same platform; they are not good. But they are excellent vehicles for soluble carbo-hydrate, and the combination is a food—a pleasant, nutritive, digestible food.

And at this point I feel inclined to be dogmatic, even more usual; unnecessarily so, perhaps, the hostile critic will say; but must be done. “Whole beef tea,” that is, beef tea to which the meat fibre has been restored after pounding, is no doubt a good food. Very good, no doubt, in convalescence from acute disease, where wasted muscular tissue has to be repaired. But in such patients as I have constantly under care at the hospitals, as physical patients either of chronic character, or with hectic in softening tubercle, or chronic bronchitis, or pleuritic effusions, and in many cases, valvular lesions or weakened walls—cases seen by me only of persons outside hospitals—it is open to question whether so very much albuminous matter is either good or desirable. The beef tea might be very much weaker and be flavoured with salt, and some soluble carbo-hydrate added. If the reader will at this point make personal trial of what is being said—an excellent practice with anything—food or medicine—that has to pass the palatal—it will be well; and will make the matter perfectly clear. Add a teaspoonful of any baked flour, of which Ridge’s food is a type, to a canful of ordinary beef tea, and then add some salt; first look at the resulting soup so produced—that is an appeal to one special sense, the eye; then taste—an appeal to the gustatory nerve. As to its palatableness it is unnecessary here to speak. But for the reader who is content to read and not to taste and try, he or she may take my word for it. Well-baked flour is largely changed into soluble dextrine; and beef tea containing some such addition is a capital food. If the flour be made from unbolted flour, that is, where the bran has not been removed, then some albuminoid matter is present, as well as the salts of the grain. Such will make an ideal fluid-food.

The steam-cooked cereals, of which there are so many varieties produced by different makers in the U. S. A., are admirably added in addition to meat broths. Plain biscuits powdered and added. The Channel Islanders always add broken biscuits to their soups, and thus give them a high food value. Soups are made much better for such addition, no matter whether made of animal or of flesh, or of vegetable matter. Sago is often added, and is better to heat in the oven for an hour before being added to the soup, is all the more digestible. The same may be said of the arrow-root. The starch-transformation into soluble dextrine.

about. Of course where cereals have been cooked by steam—much better than boiling water—this starch-transformation is more complete. Whether the soup be white or brown, cooked farina can be added with advantage as raising its food value.

Meat juice is of course a better food than a meat infusion, as it contains some albuminous matter. Mr. Mann-Valentine has identified his name with such meat juice. Others have followed in his footsteps. Murdoch has mixed meat juice with the juices of vegetables; a capital combination—in theory. But such preparations are not acceptable to the genuine Briton with his insular prejudices.

Then there are in the market “lactated foods.” That is, combinations of cereal matters with milk, desiccated. These are all good. And variety is as acceptable to persons who are put upon a “regulated dietary” as for other more fortunate persons, who can eat what they please.

The sting of a “regulated dietary” lies in its monotony. The palate palls at the restricted variety of food, and the person abstains from food as much as possible—hungers, in fact. Austin Flint and Lauder Brunton are quite right in their criticisms on the effects of a “restricted dietary.” Still, the dietary may be judicious and appropriate, and yet be varied—*i.e.*, if either doctor or patient know enough upon the subject. That is where the shoe pinches! Neither, commonly, are sufficiently familiar with food in all its aspects, to get over the difficulty. It is want of knowledge on culinary matters, that makes the dietary so restricted. And as our acquaintance with food increases, and familiarity with cookery books becomes part of medical teaching, the difficulty will be got over; and the invalid and dyspeptic be proportionately grateful.

The scientific aspect of food must be united in the bonds of holy matrimony with a practical knowledge of the cook's art, before a man can discourse learnedly of food. And another matter is—never take for granted that any cooked compound will necessarily be nice—either to the eye or palate. Taste and try! That is the only solution. I have been a judge at a food show, and ate some of all the exhibits; the palate being martyred to a great extent. I have tried most baby-foods. I have held deep consultations with cooks and housewives; I have experimented with foods to ascertain the food-value in relation to cost for some articles. (“Food for the Economical”) in *Good Words*; I have eaten every variety of tinned and preserved fruits; in fact, my personal experience extends

over the whole known field—except a cod liver oil emulsion. This is a *terra incognita* to my gustatory nerve fibrils.

Malt Extracts.—These are quite a modern form of food. Hoff's first idea was a preparation of malt which should contain all the qualities of beer, without the intoxicating alcohol. The maltster malts the grain, by which a certain starch transformation is produced; then the brewer mashes the malt with hot water, by which further starch-transformation is effected; and his "wort" is a sweet fluid, rich in maltose. The brewer then adds his yeast to the wort and then the maltose (or grape-sugar) is fermented into alcohol and carbonic acid. The malt extract manufacturer stops at the wort; and instead of adding yeast, evaporates the wort in a vacuum-pan, into a fluid like molasses. A malt-extract is, however, something more than either molasses, honey, or invert sugar. It contains, as well as carbo-hydrates, some soluble albuminoids and salts. In fact, it forms a very excellent food. It is not, however, as complete a food as milk, as it lacks fat. Some advertisements of malted foods talk of their being "substitutes for milk." That they are not;—and cannot be! But a malt-extract is a lovely food all the same. And when a stomach is in active revolt, a teaspoonful of malt-extract every hour is an aliment which offends it neither by bulk nor qualities.

Malt-extracts have been lauded for their diastasic power—their capacity to digest starch; but the diastasic power of a malt-extract is far below that of ordinary malt. Ground malt has a great future before it. As the digestive organs are enfeebled by the advances of civilization, predigested starch must come more and more to the front. And ground malt added to baked flour, or baked farin in any form, before the hot milk is poured on, makes a most digestible dish. This milk pudding should, when mixed, be placed on a hot plate, or anywhere else where its heat will be maintained, and then the diastase of the malt acts promptly upon the farina. Such a milk pudding is very fluid. It requires no cane sugar to sweeten it. This is a great matter with stomachs in which cane sugar or sugar swiftly turns acid. An egg may be added, or not, according to circumstances. This is a far more efficient method of predigesting starch than mixing a malt-extract with a farina, and then before serving in the nursery.

On the other hand, a malt extract can be made palatable for all ages by diluting it with a little water, or making it of the consistency of a syrup, and then adding an acid or lemon.

Gruels.—The admixture of farina with water or milk forms a gruel. Oatmeal is the form of farina usually employed in the making of gruel. Boiled in water with a knob of sugar dissolved, and a piece of butter added, oatmeal gruel is reckoned a sovereign remedy for a cold. Oatmeal gruel made with milk is splendid food. As the porridge of Scotland, and the North of England, this is the staple of breakfast with the whole of the working population; which, in thews and sinews as well as bones, can stand comparison with any race on the earth's surface. An excellent sustaining drink for the hay or harvest field, for foundries, stoke-holes, etc., is supplied by adding a handful of oatmeal to a gallon of water.

Boiled arrowroot was once in vogue for invalids, but is not fashionable at the present day. It was very good fuel-food; and sweetened with sugar and flavoured with wine was not at all unpalatable. But it is inferior in every way to a "treacle posset." This last consists of a pint of milk, a tablespoonful of treacle, a teaspoonful of powdered ginger, with as much rum as the drinker fancies. Not only is this good for a cold, but it has a high food-value. Such a posset last thing on getting into bed, would cheer many a starveling, and ill-nourished person.

Eggs.—The egg is certainly a fluid food until its albumen is consolidated by heat. An egg, beaten up with coffee, is a drink relished by many. Others prefer it with sherry. Some like it with milk and brandy. With a pinch of pepper and salt, and a little vinegar, an egg forms "a prairie oyster." The white of an egg added to home-made lemonade, enables it to be frothed up. This is a pleasant and nutritive drink. Under other circumstances the yolk is the part preferred. It can be taken with wine, or milk and brandy. The yolk of an egg is a constituent factor of the "rum and milk," so famous in the treatment of phthisis. Half a pint of milk, the newer and fresher the better, the yolk of an egg, a teaspoonful of sugar, a suspicion of nutmeg and a spoonful (the size varies) of rum, all beaten together and taken the first thing in the morning, has been credited with the cure of many cases of consumption. Taken early, it will often prevent the exhausting sweats which accompany the morning dose. For this purpose it should be taken about six or so, and after it, the patient often sleeps protected against hydrosis. Or it may be taken before dressing, and often enables an invalid to get over that tiresome process without a sense of exhaustion.

Syrups.—Then there are syrups of various kinds. The Germans

are fond of strawberry syrup, and raspberry syrup, the latter being the most popular: especially to breakfast, in lieu of tea or coffee. They are very palatable, and decidedly nutritive.

Syrups of all kinds are in vogue at present, too, in our mineral waters. They were in fashion in the past, with our country people, to be drunk for a cold, or when coming in from exposure to cold. The middle-aged persons cannot remember the economy of such a practice was made in every household; and very wise it was, as every sensible person thinks.

There are many circumstances where fluid food is indicated. What is most wanted is a knowledge of how to give food in this form: infusionist; and on this section of the subject, in my opinion, no apprehension prevails at present, which may well be dispelled with advantage to our sick persons.

Koumiss. A fluid food now becoming rather fashionable, is fermented mares' milk. Early travellers in Turkestan used the fermented mares' milk as an intoxicant and a diuretic among the Kirghiz. Its intoxicating properties are such as render it useless in the treatment of woman's milk, contrasting with the so-called diuretic effect of woman's milk. As a cure for phthisis, it has no established reputation in any part of Europe. Dr. Carrick, of St. Petersburg, had a number of cases brought to England and exhibited at South Kensington, in 1884. He had studied the matter thoroughly, and written a special monograph on "Koumiss" (published by Blackwood and Sons). He said: "Koumiss made from mares' milk as on the Steppes. It is a pleasant drink with the curd in small pieces on the surface. But it is not served to me." "Koumiss is equivalent to pure cream, and is as good as milk like English, but so a person cannot get it in the mountains of the arid Steppes, where the evaporation of the milk is so rapid that a person can with ease drink from three to five gallons of Koumiss in a day. Under these circumstances, the use of Koumiss are often very stilling." "From this it may be seen that it is comparatively useless to try Koumiss in the treatment of phthisis, certainly in pyrexial states it may be fed, as pure cream, and is especially suited to limited conditions of the disease, such as the intercurrent gastric disturbance of phthisis, and the rise of temperature. In some cases it produces a decided effect in

CHAPTER VIII.

PRESERVED AND CANNED FOODS.

THE qualities and properties of foods are not altered by the means resorted to for their preservation, at least to any material extent, as regards their food value. Still preserved foods require some attention in a work of this kind.

Probably the earliest method of preserving food was a blending of salting and drying. Not only was the pig so converted into bacon and ham, but the fattest oxen were selected in October and salted—the rest being turned loose to find their winter food as best they could. Mutton hams, too, were a delicacy. On such salted provisions the bulk of families in the country lived during the winter months. The pig, which could be fed on corn and the slops of the household, supplied the sole fresh meat. If the pig was not at hand (and in Scotland of yore the pig was as objectionable as an eel) and some fresh food was craved after, the cattle were driven into a pound, and bled, and this blood was mixed with barley, made into a pudding, baked, and eaten. When the pig was slaughtered, the blood was made into a “black-pudding” which kept well for some weeks. Sausages were made of the fat and lean meat chopped together and flavored with sage and other condiments. Bacon is a food of the greatest value; the lean is apt to be hard, but the fat is of the most excellent character. It is easily digestible and agreeable to the palate. It may be fried, as the rashers; and rashers of bacon with fried or poached eggs is a stock breakfast dish over the length and breadth of England; or it may be boiled and eaten cold with pickles. Beyond that, too fat bacon is used to lard fowls, liver, venison, etc., to add to dry meat as the hare and rabbit (which should always be stewed with some pieces of fat bacon), and to veal in a pie. However, as a matter of fact, it is if anything too little used as an adjunct to other food. The inside fat of the pig is “rendered” into lard, which is used in the making of pastry. The hams are a favourite food, and a slice of cold boiled ham will usually sit easily upon the most fastidious stomach. In speaking of “chronic gastritis,” Austin Flint, after recommending milk and

farinaceous food, quotes Jaccoud who was in favour of pig as a small food. "He advises that the patient be encouraged to eat certain kinds of food which are supposed to excite more than other kinds, the secretion of gastric juice, and he mentions *saucisson* as particularly eligible for that purpose." A thin slice of the meat of boiled ham cut across the fibre, well masticated, will no doubt be tolerated by a very irritable stomach.

Sausages are a very toothsome form of food. As ordinarily made they are known in London as "skin and mystery;" and the revelations which come out at times about sausage manufactories are certainly startling. Still on the whole they are fairly trustworthy, and their main adulteration is the bread crumb added. Ordinary English sausages are a good mixed food.

Then there are foreign sausages as Bologna, Strasburg, Gotha, Leberwurst, and the toothsome Gans Leberwurst made of the goose's liver. They are all savoury and many of them rich in fat. They are suitable for persons with defective teeth, as the meat has all been finely divided before being passed into the skin.

All sausages which contain the flesh of the pig, as well as bacon and ham, are liable to contain the *Trichina Spiralis*; which is only destroyed by proper cooking. Trichinosis is specially fatal where ham is eaten but little cooked, if cooked at all in some cases.

Potted meats are all savoury, from bleater paste up to *pâté de foie gras*. They are all liable to such manipulations as to allow one to say of them, "and things are not what they seem." *Pâté de foie gras* often contains but a suspicion of goose's liver, for instance. All potted foods, whether fish, flesh, or fowl, have of course been thoroughly disintegrated, and as such are suitable for dyspeptics. They do not usually form the *pièce de résistance* of a meal, but are savoury accessories. Little tiny sandwiches of stale bread, cut thin and the butter well rubbed in, and then some potted meat added, will often tempt a dainty appetite. Potted meats are usually highly spiced; a mock *pâté de foie gras* made with larded calves' liver seasoned is a very nice food. All these potted foods are "relishes" rather than material articles of diet.

Pickled foods are preserved by vinegar or brine. All kinds of fish are pickled in brine, especially herrings. They are usually soaked in water before the salt can be got out of them. Pickled pork is a great food with lumber-men and others who are out in the stores of provisions. In the U. S. A. a winter diet of pickled food with plenty of fresh air, exercise, and the fat pickled down the

to be a grand preventive measure in threatened phthisis. Mussels and oysters are pickled in vinegar and are very pleasant forms of food. Dried and smoked meats are very palatable forms of food. Fish readily lend themselves to this process: kippered salmon are famous, and so are "kippers" a form of split herring dried. "Finlon haddies" are well known, and equally so is the Yarmouth bloater. The dried haddock is unobjectionable; but the other forms are apt to upset the stomach, and can only be safely indulged in by persons who can trust their digestive powers. Kipperd salmon broken up small (after cooking) and mixed with mashed potatoes, makes a capital fish pudding.

Dried fish of a common order were once the animal food of Europe in Lent. They require much soaking and have a low food value.

Anchovies are both dried and pickled as well as potted. An anchovy on a beefsteak imparts a pleasant flavour to it. Anchovies are usually related to hot buttered toast, and hence are not desirable for the dyspeptic. Caviare, the roe of the sturgeon, is not acceptable to all palates, but it has a high food value of albuminous character.

Tongues are often dried, and when soaked and boiled are good and palatable food. Cut thin across the grain, tongue is like ham as regards its tolerance by the stomach. The Russian tongues possess quite a game flavour.

Vegetables. Sometimes vegetables are preserved by drying, as seen in the vegetable matter prepared for Julienne Soup. They are sometimes also compressed as stores for fleets and armies. They require soaking and then probably differ little from their kind in a fresh state. Many vegetable matters are pickled as relishes to food; notably the cabbage, the small onion, eschalots, gherkins, cauliflower, etc., all of which are trying to the digestive organs. Vegetables lend themselves readily to the tinning or canning process. Asparagus so treated has long been held in repute. The *petits pois* of France are miniature peas, not to be compared for a moment with the large marrowfat peas of American production. These last only require the addition of a little sugar when heating, to enable them to compete on fairly equal terms with fresh peas. Lima beans are also tinned and keep well. Succotash is a compound of the highly nitrogenised legume with maize and rich in fat. Maize in this as "Sugar Corn" is a beautiful digestible dish, whether eaten with sugar and milk, or with butter, pepper and salt. The baked beans of New England hold a high position as a food, no matter from

what point of view they are looked at. Eggs are preserved in a form not fat bacon bigger than an egg. The meat is cooked and the fat intermingled with the meat, and the high temperature at which they are baked (and its effect upon the starch), that they need no cooking in the stomach; while the fat is in a form which gives no trouble to the stomach. For food value, for cost, they stand very high. And they will come into vogue ere long with persons who are fastidious about fat.

Vegetables are easily stored, and also are easily preserved, as potatoes, carrots, turnips, beet, etc.

All *seeds* can be preserved dry, as those of the cereals, the legumes, and as such form a large part of the food of the world.

Fruits are variously preserved. As prepared in sugar by boiling they have been spoken of before. They are also pickled, and were at one time at least, largely pickled, in order to secure a good green colour. Once a pickled cucumber is cut with a knife into green pickles and fruits, to see what the pickle was deposited on the steel. Onions are largely pickled, and are usually eaten to clear the palate before drinking the wine. They are rich in oil. Red fruits are apt to be stained, and beetroot juice to improve their colour.

Tinned or Canned Foods. Tinned fruits are made in great favour. The apple, the pear, the plum, the grape, the cherry, the peach, and apricot, all find a market. They are additions to the dinner table in winter as well as in summer, are or cooked. Tinned cranberries are for sale, but the astringent native cranberries of Great Britain.

Apple rings are also gaining favour with the consumer.

It may be said as a rule that vegetables preserve better than fish and flesh.

Tinned salmon, though of good food value, is not so good as fresh salmon. Tinned lobsters vary very much in quality, according to the length of time the fish has been in the tin. A lobster is not to be despised when it is fresh. And tinned salmon will pot, or be cooked, with mashed potatoes. Other fish, such as mackerel, herring, fish, pilchards, and herrings and anchovies, are also preserved.

The sardine preserved in oil is well known, and is a good fish food, and mixed with its oil, and a little tomato sauce, is decidedly agreeable to the palate. No other fish is so

small fish get mixed with the sardines, but this scarcely constitutes adulteration.

There is one tinned fish, however, whose flavour is little affected by the tinning process, and that is the shrimp or prawn. The Barataria shrimp is delicious out of the tins.

Why the crab is not tinned—not the small soft-shelled crab which does tin—but the ordinary crab, is unknown to me. Probably there is some technical difficulty; also the eel, which would seem *a priori* to be well adapted for tinning.*

In Germany eels are pickled and sold in small barrels.

Tinned meats have established themselves over the face of the earth. Of course these tins of various sizes are most convenient for stores. A small quantity can be used at once—a great matter. Whether beef, mutton, or pork, as a rule, it may fairly be said the meat is overdone and stringy. Minced meat is fairly good and nice, and mixes well with mashed potatoes. Tinned tongues are really very good indeed; the flesh of the tongue seeming to lend itself to the preserving process very readily. Little sheep's tongues are decided delicacies, and so are tinned sheep's tails. The hare and rabbit tin moderately well. Turkey, goose, duck, fowl, and game, come out not amiss. Brawn is fair, and boar's head decidedly good. Pig's feet in tins eat capitally.

Venison I have not had a sufficient experience of, *i.e.*, in tins, to venture on an opinion.

N.B. One matter about tinned meats is never to be forgotten, viz.: the contents of a tin should be consumed as quickly as possible after being opened. No food improves by keeping after being cooked, and this is specially true of tinned meats. They not only become unpalatable but even positively unwholesome; by using them up at once the most is made of them.

Milk is now largely preserved in tins, both Swiss and other milk. That such tinned milk is a great boon for town populations, and specially for young children, there can be no doubt. Even in the country where milk is not always as easily procured as some may think, the tin of milk comes in very handy.

Cream is more difficult to manage. Partially desiccated—as Devonshire Cream, it is well-known all over England. Cream is a

* The same may be said of the mushroom, which I have never encountered in a tin—except the small French buttons, which are indifferent food from any point of view.

highly nutritive body, very rich in fat in an emulsioned form. Some time ago I wrote to the *New York Herald* suggesting the churning of cream. An enterprising man in Iowa sent me many samples; some tins turning out well, but many unfit for food. Butter is a thing could never be relied upon, and so was commercially a failure. A Dutch firm seems to have been more successful by an Examination of the Inventories (1885).

The products of milk, however, as butter and cheese, keep admirably. When butter has to be kept, some salt is added, and how long salted butter will keep is unknown. It is a most digestible fat, though of course, the object of churning is to get the fat out of its emulsionised form. It is largely used eaten to bread, and is invaluable to the cook for her dishes and her pastry. Melted butter is eaten to fish, and adds highly to its nutritive value. Butter is dear comparatively; but it is an item of household expense that should never be grudged.

It is a great mistake to cut "nursery" bread-and-butter thick; if the butter be sparingly applied there is not enough of it; if spread thick it revolts the stomach with delicate children of feeble digestive capacity. The bread should be cut thin and the butter rubbed well in, and then any child can eat it readily—yes, and digest it, too! Bread-and-butter puddings are capital food.

Cheese is a form of food requiring some consideration. It has the reputation of being most indigestible:

"Cheese is but a peevish elf,
Digesting all things but itself."

This applies to the highly flavoured cheese eaten as the last course at dinner. A nip of ripe Gorgonzola, or Rochefort, or Camembert, or Neufchâtel, or even Stilton, Cheddar, Cheshire, or double Gloucester, after dinner provides a fatty acid which aids the emulsification of fat in the alimentary canal. But this is scarcely the consideration of cheese as a food. Cheese is largely eaten as a condiment, and relish, by persons with whom economy is a matter of necessity. It is either eaten as it is, or cooked. At one time every farmer made his own cheese, for home consumption at least. Now cheese factories make, as a rule, so much better cheese than can be made that cheese-making is no longer general with farmers.

A skim milk cheese was a tough affair, but a highly concentrated form of albuminous food. A cheese made of fresh milk contains a large proportion of fat, while "ancient cheese" contains casein.

Casein softens under heat, and toasted cheese, or Welsh rarebit are sapid and savoury morsels, soon falling away if kept hot: indeed it is well to wait a little for them rather than to keep them waiting. They are reputed to be indigestible, and certainly are so with some persons. Other stomachs, again, tolerate cheese: and there are born "cheese-eaters" as well as "cheese-haters," to the one cheese is a poison, to the other it is most acceptable. To the latter even when dyspeptics, cheese need not be forbidden. A great matter is its disintegration. Finely grated in soup or even boiled macaroni it is unobjectionable. Macaronic cheese done in the oven, or before the fire requires stronger digestive powers. If hominy be well boiled, and then grated cheese with milk mixed with it, and the whole put into the oven to brown, a delicious digestible dish is the result. Crushed cereals, shredded maize, samp and broken biscuit so treated with milk and cheese, are admirable foods of high food value, savoury, and finally, digestible.

Cold cauliflower and vegetable marrow served up with cheese *au gratin* are very good: while Staffordshire folk fry cheese with bacon, and for this end prefer a crumby cheese—not otherwise in favour for culinary ends.

CHAPTER IX.

PREPARED FOODS.

BY "prepared foods" here is meant the foods which are *made* for babies and invalids. To Liebig, among so much more owing to food, we owe the introduction of foods specially prepared for infants. The food of the newly-born is milk; but many attempts have been made to substitute for it, mainly or in part, other foods: the motive being in some cases economy, in other cases the difficulty of procuring milk in sufficient quantities. Sometimes the mother could not suckle the child, being ill, and sometimes dead.

By what instinct persons were led to choose the crust and seed of the loaf to boil for babies' food, can now only be a matter of speculation. After thorough boiling the water was poured off more or less completely, and some milk added. The outside of the loaf is the part most thoroughly exposed to the baking process, and therefore in it the starch is most completely converted into soluble dextrine. Such carbo-hydrate matter with some fresh cow's milk formed a very complete food. But such "pobbs" with only a little sugar, is a poor food; and the marvel is that such lusty infants ever were raised on "pobbs" as undoubtedly have been the case in the agricultural districts of England and elsewhere, where milk is not easily procured. How, and by what means the growing organism has secured materials for tissue-growth out of this unprovisional food, is a mystery, when looked at from a scientific point of view. There came rusks and "tops and bottoms," and however, made with less flour, and lacking the salts and albuminoids of the bread. Then came the introduction of malted preparations.

Something has been said before about the preparation of malt extracts, and this need not be repeated here. The wort is allowed to run down to the consistency of honey, and is a sticky and therefore a difficult food to handle; albeit a beautiful food. As a means of procuring any diastase activity, a dry preparation like Maltin is a much better way preferable. There are dozens of such foods on the market, and many very good foods they are, too. They are especially valuable in the

to milk for adults as invalids and dyspeptics, and to a little milk and water for babies.

Combinations of baked flour and malt are now sold by the hundred as babies' and invalids' foods. They vary somewhat in composition. Some contain more malt than others; some contain some desiccated milk. No doubt these last are the most complete foods, as from the milk a certain amount of fat is present. They are also most palatable.

Pure malt in fine powder must soon be a regular article in the market. It is economical as compared to the malt-extracts, fluid or dried, and contains much more diastase activity. Added to baked flour in quantities varying from one-third malt flour to two-thirds of baked flour, up to half and half, and then hot milk either poured on—(for a pudding) or added for a drinkable food, malt can be made most useful. Ground malt, too, contains the husk of barley and is rich in phosphates. Maltose, dextrine, soluble albuminoids, and phosphates, all are there in a very palatable form. Only fat is lacking. And that can be got too, in the *cremor hordeatus*, and in *oleobyne*; both of which contain milk-fat in considerable proportions.

The ordinary prepared foods in powder are legion. There is one good thing about this, and that is, there is choice and variety. One form is unpalatable to a baby or invalid, while another pleases the palate. The palate grows tired of one food, then try another for a change. Monotony is the curse of all dieting, and should be avoided as far as possible.

A medical man should not be wedded to one food, but should be more Catholic-minded. This is a matter far too little considered, and yet it lies across the very threshold of resort to prepared foods. Change about from one to another just as you expect your cook to cater for yourself.

Say for breakfast the invalid has some milk and Mellin's food; at eleven o'clock some whole beef tea with baked flour in it; at luncheon a milk pudding, made with broken captain's biscuit, or other biscuit; at tea-time some chicken broth with shredded maize; and at supper a digestive biscuit and butter with a glass of sherry. By attention to these items—but very important items—the invalid's dietary could be made acceptable instead of revolting, as it too often is.

When I read of Dr. David Vandall fasting when ill with typhoid fever, I knew Mrs. Vandall's capacities as a housewife too well to suppose for one moment that monotony in food was the cause of his

fasting; and that his conduct would cause a great deal of suffering to many an invalid hunger, partially at least by reason of the want of variety. Variety indeed is quite as acceptable to the dyspeptic as to anyone else; and babies no more have to be taught about food just as much as their elders. Because the baby cannot tell us what it thinks, it is not therefore to be assumed that it has not its thoughts, and these it communicates by its rejection of one food and its delight in another. A very young baby can reason. I remember once seeing a youthful mother nursing her first baby, and putting the mouth piece of her feeding bottle in the baby's mouth and wondering why it would not suck. The baby got nothing but air when it did suck, and so abandoned the effort; it did not want air; it wanted milk! On directing her how to have the teat filled with milk before putting it in the baby's mouth, it sucked away famously, and developed into a lusty child promising to grow up into an astute Yorkshireman. So with the prepared foods, one baby likes a good share of malt in the "food," while another prefers a certain food in which there is little or no malt; while another again delights in a food in which there is a certain admixture of leguminous flour. Tastes they have, of course, and preferences—like other and older folks!

And of course the same individuality of taste and palate belongs to dyspeptics and invalids. Because one form of prepared food is distasteful, it does not follow that all are. If there existed a little more inventiveness in nurses there would not be so much complaint of monotony of food on the part of sick persons. When the food consists solely of milk and seltzer water, beef tea and calves' foot jelly, is it any wonder that the palate is palled? And yet this is the complete range of invalid cookery in many households. Not long ago I was in company with three general practitioners, two of the good average specimens and one decidedly far above the average, and they could go no further. It was only too clear that they were not in a position to afford much suggestion as to the dietaries of their patients. They answered "slops" to my leading question, but their acquaintance with "slops" in concrete forms was very limited. Yet there was no reason to suppose them specially ignorant of their fellows.

Every medical man ought to try the various prepared foods in various ways, so as to personally understand what they are like about when speaking of various foods. Other workers in the "dumpling" on the subject. I wish my acquaintance with them

prepared foods of the United States was anything like as extensive as my knowledge of the prepared foods of Great Britain, for then I could speak freely on the different forms. To mention those I happen to know, would exclude many of which I have never heard, which may be equally good.

All sweet foods containing malt go best with milk; while plain baked flour suits best meat-infusions. Lentil preparations have a strong flavor whatever mixed with. Pea flour is best with meat-infusion as the "brose-meal" of Scotland. The highly nitrogenised character of all leguminous flours renders them suitable rather for the hale than the sick. And much the same may be said about oatmeal, as it does not agree as a rule with a sick stomach. Its richness in fat has something to do with this, and a fatty acid is apt to be brewed therefrom in many stomachs. It is the fat in oats and maize which makes them objectionable with many. Rye is a pleasant addition to wheaten flour to the fancy of many, as seen in the pumpernickel of Germany, and the brown loaf of the North of England, especially in the farmhouses, where brown bread and skim milk cheese are large factors in the dietary.

Prepared foods, which contain a certain proportion of desiccated milk, have a high food value, and require only water (with a pinch of salt) for their culinary treatment.

There is one form of prepared food which is, so far, unique, and that is the *cremor hordeatus* of Loefflund, which is made from the best cream from selected herds on the Bavarian Alps, deprived of a certain amount of water and preserved by malt extract. This highly palatable food is excellent in its food properties. It contains some albuminoids, a considerable quantity of fat, and some carbo-hydrates. It can be eaten alone, and in certain cases where the stomach is irritable and contracted, so that only small quantities of food can be tolerated, a piece the size of a hazel nut may be taken every hour. For an invalid in bed this is quite sufficient to keep up the strength until the stomach is well enough to hold a larger buck at once. Or it can be used to coffee or tea, and is especially indicated in those cases where cane sugar gives rise to acidity.

Of like character in the main, is the new food called "oleobyne," which as its name indicates is a combination of fat and milk. It also is very palatable, with a high food value. Its fat is in an emulsified form, and therefore very digestible. Such foods are adapted to persons whose tissues require more fat and yet who dislike cod liver oil: a very large community. For persons of phthisical ten-

PREPARED FOODS.

deny these foods are suitable, while the fact is, they are. Properly prepared, they should be thrown much upon a meat diet, and a good deal of them should make these foods part of their stock. They are especially indicated for persons with indolent Bright's disease, and for those highly charged with nitrogen. Another name for one of these foods is "kepler," where the oil is incorporated with milk, and the milk is

Then there are cod liver oil emulsions to be considered. They, of course, contain only fat, and that in the form of oil, and are not digestible, but otherwise, probably, the least digestible form of fat. Good cod liver is frozen to get out the stearine, and as much of the palmitine as possible, the remaining fluid being very pure oil. Such fat is often digested when other fats, as animal fat, and tallow, are rejected. Nothing is easier than to make an emulsion with oil, fat, suet, or oil. But the difficulty is to make an emulsion that will keep an indefinite time. If a pancreatic preparation is used, the tendency is, for the ferment to act upon the fat, and split some of it up into fatty acids, matters very irritant to the stomach, and not especially welcome to the palate. There are many such preparations to be bought; and when they are ready, a preparation can be bought as required in a fairly new and fresh state.

There has been a prejudice, and a strong one too, against prepared foods. First announced as "baby-foods," and used for infants, the public did not seem to notice the "and babies," and in many instances follows the "babies." Advancing science, however, breaking down this prejudice, is helping society. A more general recognition of the value of so-called artificial sweeteners, than cane or beet sugar, on the part of the medical profession, is desirable; and when systematic lectures on food are given, and given, every young medical man will understand that pre-digested farinaceous matters, and their glucose, are not for the weakly digestion. Grape sugar is the food of the baby, and is supplied to a certain extent every one. Undigested starch is not in the stomach and prevents its getting on with its own work. Prepared foods with pre-digested starch are not a menace to the digestive powers, but under the most favorable conditions, and the still greater study of the subject. The various kinds that have been subjected to analysis, and are found to be foods of farinaceous origin, must, except in the case of the soluble preparations, be rejected in the kitchen. Soluble preparations are not for the muscular fibre, but for the nervous system, and for the brain. Sidrabile quantities of soluble preparations are used in the

By varying the soluble meat and the carbo-hydrates, variety can be got, if not a very wide range. Then where the stomach resents fat floating about in it when at work, the fat can be taken when the gastric activity is over, as cream, or emulsionised fat artificially prepared. Such a scheme of dietary is not very attractive; but then we all know that we must cut our coat according to our cloth—not according to our liking.

CHAPTER X.

ARTIFICIAL DIGESTIVE AGENTS. ARTIFICIALLY DIGESTED FOODS.

NECESSITY is the mother of invention, says the old saw. And failure of the digestive powers has led to the adoption of artificial digestive agents. First the chemist seized upon the diastase of the seedling to help our human saliva. Then he appropriated the digestive agent in the pig's stomach and utilized it for the needs of man. Hence pepsin in its various forms. Finally on discovering the functions of the pancreas, that viscous was enlisted as furnishing a most useful recruit to the ranks of the artificial digestive agents.

Of the action of diastase upon starch, enough has been said, both as to its action and how to make use of it in practice. No more requires to be said. The diastase, or rather the matter containing it, can be added to the starchy food: 1, before being taken into the mouth; or, 2, while being taken into the mouth; or, 3, immediately after the food is swallowed so as to act before the stomach becomes too strongly acid for its operation.

Pepsin is an agent which acts in an acid medium, and concentrates its digestive qualities upon albuminous materials, so to say. It is put up in a fluid form, in the powder, in pills, and in tablets. It is impossible to express any opinion upon the respective merits of the different forms. The matter is to secure a good and potent action. The pills and tablets have the advantage of being taken and swallowed. The palate is not offended, that's one matter. There, there is another. There are many persons whose digestion requires the help, who are yet well enough to visit and dine out, and it is under these circumstances that some artificial digestive agent is necessary. A bottle of fluid is out of the question, because the pills or tablets can easily be opened and the contents taken at the table without attracting attention. For the pills and tablets are sold in a plain wrapper, and are not even patented. Indeed by so doing the manufacturer is enabled to dine in company with one who is not a doctor, and the physician, or careful selection of the medicine, is

to digestive aids:—both exceedingly unpleasant for nervous persons, who are very liable to be dyspeptic.

Where a meal is taken alone, it is a matter of comparative indifference in what form the pepsin is used.

Then come pancreatic preparations of all kinds and varieties. So far, we have been most familiar with the fluid preparations recommended by Sir William Roberts, F.R.S., of Manchester. The pancreatic fluid has a large range of digestive activity. It transforms starch into grape sugar; it digests albuminoids (in an alkaline medium); it curdles milk, and it emulsionises fat. The difficulty about it is, however, that it is inoperative in an acid medium. The acid stomach lies betwixt the mouth and the duodenum—the area where the pancreatic secretion exercises its action. Dr. Roberts met this difficulty by placing his *Liquor Pancreaticus* under the protection of an alkaline guard, in the form of 10 or 20 grains of bicarbonate of soda. But the soda has a filthy taste, and renders the draught most offensive to many palates. It has seemed to me, instead of soda and water, it is well to give the pancreatic ferment in a glass of milk with as much prepared chalk, or light magnesia as would lie on a sixpence—about the size of a male thumb-nail. This is a far less outrage on the palate. It is a preferable form of alkali to my mind. By waiting until the acidity of the stomach is exhausted ere giving this draught, less alkali is required, and the result is more certain. The acidity of the stomach wanes as gastric digestion reaches its completion; and there's little acid left to neutralize about an hour and a half or two hours after a meal. This is the time to select for the artificial pancreatic digestive agent to help out the natural pancreatic secretion.

The same principle of action must guide us in the use of pancreatic agents whatever their form; and other than fluid preparations are already being manufactured.

Combinations of the several digestive agents on the "shot-gun" principle are largely sold. Unphysiologists, as they appear they certainly are often useful. "If the one misses, another hits." That is the principle. If the contents of the stomach be still acid, then the pepsin comes in useful. If the food be passing the pylorus, and coming into contact with the alkaline chlorides of the bile, then the pancreatic products are ready for action.

Such then are in brief, the artificial digestive agents.

Now as to predigested foods. Enough has been said about predigested starch.

Some description of them becomes essential, as there are cases where they are required. But whether great gastric irritability existing is an indication for their use is a moot matter. I am not aware that it is pleaded that peptones sit more easily upon the stomach than proteids, though it may eventually turn out that such is the case. Here it is rather a question of bulk than of anything else. Small quantities at once of whole beef tea, with some baked flour, malt extract, or milk with or without some soluble carbohydrate, should be tried. Especially in gastric catarrh is this desirable. If anything approaching solid food be taken then, as it is rolled over in the stomach, it becomes coated with mucus, and so forms an utterly indigestible mass. Here peptonised milk may become useful as no longer undergoing or requiring the curdling which might transform ordinary milk into something of a solid. In gastric ulcer predigested food would pass readily through the stomach and occasion little movement, and would do away with the pain caused by the muscular contractions dragging on the base of the ulcer. The presence of the acid of the gastric juice irritates the raw surface, and vomiting relieves the stomach at once of both acid and movement, and so immediate ease follows. The circumstances under which resort to predigested food is indicated, can only be broadly spoken of here, and must in each case form the subject of private judgment upon the part of the medical attendant. After such cautions and qualifications, some account may be given of the means of preparing them: following upon the lines laid down by Sir William Roberts.

THE PREPARATION OF PEPTONISED FOODS.

In peptonising, or partially digesting food by means of Liquor Pancreaticus, it is important to remember that the liquor must not be added to food of any kind at a higher temperature than 140° Fahr. This temperature can be estimated with sufficient accuracy, should no suitable thermometer be at hand, by tasting. If too hot to sip without scalding the mouth, it would entirely destroy the activity of the liquor pancreaticus, and must be allowed to cool to that point before such addition is made. This rule will hold good with other pancreatic preparations.

Peptonised Milk.—A pint of milk is diluted with a quarter of a pint of water, and heated to a temperature of about 146° Fahr., (60° C.). Or the diluted milk may be divided into two equal portions, one of which may be heated to the boiling point, and then added to

the cold moiety; the mixture will then be of the required temperature. (We are not told what is the temperature of the cold portion, but we may assume it is below 80° Fahr.) Two or three teaspoonfuls of liquor pancreaticus, together with ten or twenty grains of bicarbonate of soda (about half a small teaspoonful) are then mixed therewith. The mixture is then poured into a covered jug, and the jug is placed in a warm situation under a cosy to keep up the heat. At the end of an hour, or an hour and a half, the product is raised to the boiling point for two or three minutes in order to arrest any further digestion. It can then be used like any ordinary milk.

The object of diluting the milk is to prevent the curdling which would otherwise occur, and greatly delay the peptonising process.* The addition of bicarbonate of soda prevents coagulation during the final boiling, and also hastens the process. The purpose of the final boiling is to put a stop to the ferment-action when this has reached the desired degree, and thereby to prevent certain ulterior changes which would render the product less palatable. The degree to which the peptonising change has advanced is best ascertained by the development of a bitter flavour. The point aimed at is to carry the changes so far that the bitter taste is distinctly perceptible, but is not unpleasantly pronounced. The extent of the peptonising action can be regulated either by increasing or decreasing the dose of liquor pancreaticus or by increasing or decreasing the time during which it is allowed or permitted to operate. By skimming the milk beforehand, and restoring the cream after the final boiling, the product is rendered more palatable and more milk-like in appearance.

Peptonised Gruel.—Gruel may be prepared from any of the numerous farinaceous articles which are in common use, — wheat flour, oatmeal, arrowroot, sago, pearl barley, pea or lentil flour,†

The gruel should be well boiled and made thick and strong. It is then poured into a covered jug and allowed to cool to a temperature of about 110° Fahr. Liquor pancreaticus is then added in the proportion of a tablespoonful to a pint of gruel, and the mixture is kept warm under a cosy as before. At the end of an hour or so, when the product is boiled and finally strained, the action of the pancreatic extract on gruel is twofold; the starch of the cereal is

* May not the matter go far to explain the necessity of diluting the milk with an alkaline water as compared to the ordinary water?

† In accordance with what has been already said, the gruel should not have already been exposed to a high temperature, as this would have destroyed the milk.

verted into sugar, and the albuminoid matters are peptonised. The conversion of the starch causes the gruel, however thick it may have been, at starting, to become quite thin and watery. Peptonized gruel is not generally, by itself, acceptable food for invalids, but in conjunction with peptonised milk (peptonised milk-gruel) or as a basis for peptonised soups, jellies, and blanc-manges it is likely to prove valuable.

Peptonised Milk Gruel.—This is the preparation with which Dr. Roberts has had the most experience, and with which he has obtained the most satisfactory results. It may be regarded as an artificially digested bread and milk, and as forming by itself a complete and highly nutritious food for weak digestions. It is very readily made, and does not require the thermometer. First, a good thick gruel is prepared from any farinaceous material. The gruel while still boiling hot, is added to an equal quantity of cold milk. The mixture will have a temperature of about 140° Fahr. To each pint of this mixture two or three teaspoonfuls of liquor pancreaticus and twenty grains of bicarbonate of soda, (half a small teaspoonful) are added. It is then kept warm in a covered jug under a cosey for a couple of hours, and then boiled for a few minutes, and strained. The bitterness of the digested milk is almost completely covered in the peptonised milk-gruel,—and invalids take this compound, if not with relish, without the least objection.

Peptonized Soups, Jellies and Blanc-manges.—Dr. Roberts has sought to give variety to peptonised dishes by preparing soups, jellies, and blanc-manges, containing peptonised elements. These contain a large amount of digested starch and albuminoid matter, while possessing excellent flavour, and which the most delicate palate could not accuse of having been tampered with. Soups were prepared in two ways. The first way was to add what cooks call "stock" to an equal quantity of peptonised gruel, or peptonised milk-gruel. A second and better way was to use peptonised gruel, which is quite thin and watery, instead of simple water for the purpose of extracting shins of beef and other materials employed for the preparation of soups. Jellies were prepared simply by adding the due quantity of gelatine or isinglass to hot peptonized gruel, and flavouring the mixture according to taste. Blanc-manges were made by treating peptonised milk in the same way, and then adding cream. In preparing all these dishes it is absolutely necessary to complete the operation of peptonising the gruel or milk, even to the final boiling before adding the stiffening ingredient. For if liquor

pancreaticus be allowed to act on the gastric contents, the food undergoes a process of digestion, and its power of nourishing is utterly abolished.

Peptonised Beef Tea.—Half a pound of lean beef is mixed with a pint of water and twenty grains of soda. This is simmered for an hour and a half. When cooled down to 140° Fahr., a tablespoonful of the *Liquor pancreaticus* is added. The mixture is then kept warm under a cover for two hours, and occasionally shaken. At the end of this time the proportions are decanted, and boiled for five minutes. Beef tea prepared in this way is rich in peptones. It contains about 4 per cent. of organic residue, of which more than three-fourths consists of peptones, so that its relative value in regard to nitrogenous materials is about equivalent to that of milk. When seasoned with salt it is scarcely distinguishable in taste from ordinary beef tea.

Another way.—One pound of the young beef, lean beef is mixed with one pint of water, and simmered for an hour and a half. The resulting beef tea is then decanted off into a covered jug. The undissolved beef residue is beaten with a spoon into a pulp, and added to the beef tea into the covered jug (where beef tea). When the whole is cooled down to 140° Fahr., (or when it is cool enough to be tolerated by the mouth) a tablespoonful of the *Liquor pancreaticus* is added, and the whole well stirred together. The covered jug is then kept warm under a cover for two hours, at the end of this time the contents of the jug are boiled briskly for two or three minutes, and finally strained, and served ready for use. The extreme solubility of digested products, which is so manifest in buminoids, detracts from their acceptability to the patient, and when they appear thin and watery, they miss the taste of substance and solidity, which is characteristic of food in its natural state. But for the weak invalid without appetite, this source of nourishment is generally an objection, and it is taken with more relish when it is presented in a form which they can drink like water. When an artificial food has been made, then the soups and broths are not to be neglected.

The Use of Liquor Pancreaticus as an Addition to Food shortly before it is Eaten.—This is a very useful mode of using the pancreas, for the contents of the stomach are not digested before the food, or coffee, or cocoa, or fruit, or any other food, is taken, or when it is taken with the *Liquor pancreaticus*. A teaspoonful or two of the *Liquor pancreaticus* is added to the

up with the warm food as soon as it comes up to table. And such is the activity of the preparation that, even as the invalid is engaged in eating—if he eat leisurely as an invalid should—a change comes over the contents of the cup or basin: the gruel becomes thinner, the milk alters a shade in color, or perhaps curdles softly, and the pieces of bread soften. The transformation thus begun goes on for a time in the stomach, and one may believe that, before the gastric acid puts a stop to the process, the work of digestion is already far advanced.

The food should be cool enough to sip before the Liquor Pancreaticus is added, else the ferment is killed.

Peptonised Enemata.—Such peptonised milk-gruel as has been described, is admirably adapted for enemata. Here the palate has not to be considered, and monotony is not distasteful. Peptonised beef tea, with some soluble carbo-hydrate may be used at times. But the fat in the milk makes the milk-gruel more desirable, in theory at least: for how far fat can be taken up by the large bowel is a matter on which the writer knows nothing personally, nor, on enquiry, can he find anything to favor the view of fat-absorption by the large intestine.

There lies no objection to peptonised preparations for enemata on account of their taste, and an enema of peptonised beef tea, with any of the prepared foods consisting of soluble carbo-hydrates, would be easily prepared in any household, however humble.

There are other preparations than fluid pancreatic preparations, and the following are from the instructions furnished by Messrs. Fairchild. This form has its pancreatic preparation in the form of a powder—*Extractum Pancreatis*. It is sold in glass tubes, each containing five grains of the Extractum, and fifteen grains of bi-carbonate of soda. (They are sold in boxes of one dozen). The contents of one tube will digest one pint of milk in half an hour.

The procedure is simple:—"Into a quart bottle pour a pint of milk, a quarter of a pint of water, and one peptonising powder. Let this stand for thirty minutes in water as hot as the hand can bear, and then boil for two or three minutes. It is then ready for use, and should be kept in a cool place."

Instead of testing the temperature by sipping, Dr. Roberts' plan—Messrs. Fairchild prefer the hand: either is a good and ready test. From the evidence Messrs. Fairchild have been able to collect, they are strongly of the opinion that peptonised food is more readily borne and tolerated by the stomach than food not predigested. They

say of peptonised milk. "It is retained in the stomach, especially of the digestive tract, when even the most powerful is rejected." Time will tell if this surgery has been successful. As to the peptonising of meat their directions are: "Take one pound of whole beef tea and half a pint of water, then add 10 grains of Pancreatic Extract, and twenty grains of bicarbonate of soda, and place in a warm place not over 140° F., for three hours, and then add 10 grains to kill any remaining activity in Pancreatic Extract." So it is seen that acid is essential to the action of trypsin as acid is to the action of pepsin.

The use of peptonised foods became very extensive, so much so that Sir William Roberts introduced them to the notice of the public, but the demand is now much less. Why is this? Probably one matter is that those practitioners who love to dabble in the very latest novelty, had peptonised food prepared in many cases where ordinary food would have served equally well, and so gave the friends and nurses of many invalids a great deal of unnecessary trouble. Then other novelties cropped up and hustled peptonised foods out of the memories and minds of the said novel-loving practitioners. That is part of the explanation. Another part is that the resultant products were not palatable, because the instructions were not rigidly carried out. A fine instrument in the hands of a careless fellow is soon put out of working order, and the finer the instrument, the sooner disaster is induced. The method of peptonising food is a matter of precision, 1st, the right temperature—high enough to permit of the ferment readily acting, and yet not too high to kill it. If too high no peptonising is possible, if too low the digestive agent is inoperative, and no peptonising follows. 2nd, The neglect of raising the fluid to the boil at the proper time. "The hotter the oven the shorter the time" is a maxim to be observed. If the fluid was kept warm under the cover the time to raise it to the boil comes earlier than it does, when, some cooking occurring, the time to raise it to the boil is delayed, then, the time to bring complete digestion is brought out, and so the resultant product is offensive to the palate.

It is easy to see how carelessness, or perhaps too much haste, led to how the want of careful exactitude led to some of the failures. If these were only used when it is understood that they are to be used, it would follow that proper care would be exercised. As a result it follows upon use, and dabbling in, and dabbling in, and dabbling in, curiosity or keep somebody out of a subject, and keep it as a valuable measure into disrepute, and out of use.

In ordinary cases, predigested carbo-hydrates are enough with whole-meat broths, and milk with seltzer water, in small quantities at once, alternating with malt-extract for variety. If such dietary is insufficient, and life is quivering in the balance, then some responsible person must be put on to do the peptonising, with a full sense and consciousness of the grave responsibility of his or her action. If this were done, then the practitioner could rely upon his weapon as being well forged and trustworthy. But to set a meddlesome fool to work peptonising food, is to inaugurate a solemn farce, and if the case be really a grave and serious one, such procedure is little less than manslaughter, and deserving of the most severe censure. If a medical man does not possess sufficient acumen to discriminate his agents, he is not fit for the post he has assumed. And if he cannot exact military obedience from his subordinates, he should dismiss them from the service, as a General does an incompetent soldier; and if he cannot do that, he should dismiss himself from the generalship of the case—and his resignation would be the best plan for all concerned.

With proper care and precautions, peptonised foods are a most valuable aid in many cases, especially where there is any morbid condition of the stomach present. They require little or nothing of natural digestion, and sit easily on the stomach; but then such food requires to be made with as much care as is requisite in putting together the works of a watch—if it has to go well.

An attempt to provide a peptonised preparation which only requires some hot fluid, and so dispense with household peptonising, has been made in "Soluble Food."

With this last remark, which indeed applies broadly to all culinary preparations, but specially to peptonised foods, Part I. of this work fitly concludes. A knowledge of what food is, of what it consists, chemically, of its destiny in the body, of the effects of cooking, and the different portions of the digestive act, must constitute the basis or foundation of all dietetics, and must be thoroughly mastered in detail before the nutrition of a sick person can be safely approached. It is the want of such knowledge which has hitherto hampered the profession in dealing with the matter, leaving each practitioner single-handed to follow his own experience—an experience often leading him in a manner which reminds me of a clergyman's answer when asked if a certain parishioner followed his conscience. "Oh yes," was the reply, "he follows his conscience very much in the way he may be said to follow his conscience, but he follows

it in front of him!" A man, to some extent, is ignorant of his own mind as he goes, tentatively, on certain lines, and he does not know how to turn how? The fashions which have swayed the gastronomic room, demonstrate that no guiding principle has been in the matter. Veal broth and eel broth had yielded to the milk and seltzer water, over which beef tea usurped a position which it still holds. But the facts are slowly and imperceptibly changing, and common sense, illumined by scientific knowledge, is attacking its fortress, and, before long, we may fairly hope, the pretensions to be a sustaining food will be abandoned, and a more and sounder position taken up, viz.: that it is (as originally prepared) a stimulant vehicle, pleasant to the palate for some of its carbohydrates — no mean position by any means for it to hold. Little knowledge in the making, and "knowledge grows, not wisdom lingers," as is evinced in a piece of instruction in a medical journal, which came under my notice, while penning this chapter. Swedish fish there is advocated as a food for the diabetic, and especially oysters. Now the great bulk of the oyster (like some other sea-fish), consists of its liver, which is crammed with glycogen, or animal starch. How such a curious and pernicious error can have found its way into a respectable medical journal, is one of those things which Lord Dundreary decided "No fellow can understand." Were dietetics are properly taught in our medical schools, and every medical student is made to understand the chemical composition and utility of each article of our dietary, as carefully as he is taught the action of each unit of the Pharmacopœia, he will handle his food as skilfully as he wields each medical agent. And if that which he can do this, he will achieve results when many agents are used by others as well as himself.

Every sick person is a dyspeptic for the time being, and it is the dyspepsia of acute disease, or of chronic disease, or of trouble connected with the stomach, or of the bowels, or of the liver, has only to encounter "impaired digestion," and "indigestion," patients at all! And a knowledge of such a nature is a constant acquaintance with *Materia Medica* and all the rest of it, is a neglect of one or the other, than a neglect of the stomach and bowels, and rungs.

PART II.

CHAPTER XI.

FOOD IN INFANCY.

HAVING discussed the subject of Food, in its entirety, it now becomes possible to enter upon the practical application of this knowledge.

When the babe is newly born it was, and in some cases may still be, customary to put something into its mouth; commonly some sugar and butter.

The child had had a long fast, it was pleaded, and the welcome to the little stranger could best take the form of some nourishment. The motive was good, at any rate. "At birth the stomach and bowels, not having been used for the purpose of digestion, contain a quantity of mucous secretion—meconium—which requires to be removed before they can enter upon their function. To effect this object Nature has rendered the first portion of the mother's milk purposely watery and laxative; and on the part of the infant nothing further is required than to allow it to follow its natural instinct and suck it in." So says Dr. Andrew Combe in his famous treatise on "Digestion and Dietetics." Laxatives, other than this, are not required. He has some more remarks which seem to me to be worth quotation.

"At birth the stomach is feeble and as yet unaccustomed to food, its cravings are consequently easily satisfied, and frequently renewed. A healthy infant seeks the breast with avidity, but sucks little at a time. It leaves an interval for thoroughly digesting the little which it has swallowed; after which its appetite revives, and a fresh supply is demanded in language which no mother can mis-interpret. During the first month appetite ought to be the mother's guide in offering the breast, and if she know how to read the expression of her infant aright, she will want no other. At that early age there ought to be no fixed time for giving nourishment. The stomach cannot

be thus satisfied. In one child digestion may be so slow that the interval be consequently too short; in another it may be so rapid that the interval too long. But the active ear of the child is the guide, which need never be mistaken, and none else ought to be so trusting."

Overfeeding is, however, the danger *par excellence* of lactation; most of the troubles of early infancy are due to this, as constipation, irritation and diarrhoea. The carminatives, aromatics and antacids, so frequently given to infants are required mainly as a consequence of errors in feeding. We must not put out of sight, however, the fact that any derangement of the maternal health, any improper food on her part, even a mental emotion may so affect the milk as to alter its qualities. This is a well-known fact with mothers. Further, if a child is constipated, the pleasantest method of correcting the bowels is to give some mineral laxative to the mother; and of all these the phosphate of soda suggests itself on physiological grounds. The dose to be given to the mother which is sufficient to influence the child is usually not so large as to affect her health. The mother's milk is usually quite sufficient for the first six months of existence. Nor need the mother be specially fed for the purpose of sufficient lactation.

Of old, if the mother was removed by death or from any reason rendered unequal to suckling her babe, a wet-nurse was procured. For the purpose, a mother in a humble position of life was usually selected. But "the health and usefulness of country nurses are often utterly ruined by their transplantation into the families of rich and luxurious employers," said Combe.

The "wet-nurse" is now almost a thing of the past, and her modern substitute is the "feeding-bottle." But a new care is required with the one as with the other. Perfect cleanliness is requisite; scalding with hot water containing some soap, or boiling, after each use of the bottle; the bottle, the tube, and the nipple to be placed in a clean glass. The milk is usually furnished by the owner of the cow, in a matter of indifference how that cow is fed. If given but coarse, luscious, lush grass, and other food which is too rich for a cow, the milk of cows so fed is apt to be rich, but it is not so plentiful. In most dairy-yards a cow is kept for the purpose of furnishing milk for babies. Sometimes the milk is given to a baby, for no apparent, or only a slight reason, and then the owner changes the milk, and it is in a moment changed from rich to poor.

In other cases the child milk is procured from a cow which has a special reputation for the purity of her milk, and is given to the

peculiarly wholesome from the herbage which forms the dietary of Swiss cows. Tinned milk usually contains some added sugar, and consequently only requires a certain dilution with water (about one third) to simulate human milk very closely.

This leads us to a subject which has been threshed out again and again. The curd of human milk is lighter and softer than that of cows' milk. The milk of various animals is used as a substitute for human milk. (According to Dr. Thomas Dolan, considerable differences, sufficient to give the subject a practical value, occur in the milk of different women). On this matter Dr. Pavy says: "The milk of the cow, according to above analysis, (Payen's) the most closely approximates to that of women, but it is rather more highly charged with each kind of solid constituent. Next follows the milk of the goat, which, taken altogether, is again rather richer. That of the sheep is characterized by its marked richness in nitrogenous matter, and butter. The milk of the ass and mare present striking differences from the rest. The peculiarity consists of the small amounts of nitrogenous matter and butter, and the large amount of lactine or sugar."

There are variations in the composition of the milk in different females of the same species of animal, just as there are in different women; hence the variations in the analyses and estimates made by different authorities. The milk of the Alderney cow is the richest of all cow's milk; and FHeretier found the milk of a *brunette* richer than that of a *blonde*,—a conclusion borne out by the further observations of Dr. Dolan of Halifax, while Dr. Hill Hassall found that the morning milk of a cow is not so rich in cream and casein as is that given in the evening. Looking at the analyses of milk, it would seem that a small addition of water to cow's milk, brings it down to human milk; while some contend for a small addition of sugar. Nor need necessarily the sugar be cane sugar; a little maltose sugar is easily procurable, as in Mellin's Food, for instance. The advantage of maltose sugar, in whatever form, to the milk is that maltose sugar rather undergoes lactic acid fermentation, while cane sugar undergoes acetous fermentation, and acetic acid is far more irritant than lactic acid, whether free or in combination with a base.

For some months milk is all a child requires; and, if it can get a sufficiency of milk, nothing else is desirable. But amongst the poor the amount of milk is apt to fall short of the infant's needs, and then some substitute must be found. Up to a recent time the

crust and sole of the wheaten loaf were broken up, and the resultant product, "pobbs," mixed with milk. This compound was not objectionable; but when a little sugar was added to the "pobbs," then partial starvation was the result, as we have stated before. The addition of a little butter to the "pobbs" is a remedy; but even that did not provide the satisfaction of the "pobbs" (rides) which are so desirable. Specially prepared "Infants' Foods" have much the same food-character; but at last the subject attracted the attention of the chemist who started to prepare special "Infants' Foods;" of which generally it may be stated that they are good. This last can scarcely be said of "lactated" foods, that is, foods containing a proportion of milk. Of the importance of fat to infants no one can entertain a doubt, after witnessing the effects of its deficiency upon an ill-nourished child. It may be said of almost all the prepared Infants' Foods, they are intended as additions to milk, not as substitutes for it. Until the teeth are out, and the effort of making their way through the gum, arouse the salivary glands into activity, the infant has no means of dealing with starch, before it reaches the pancreatic secretion; and raw unaltered starch is not acceptable to the infant stomach. Numerous disorders of the alimentary canal can fairly be laid to the charge of unaltered starch.

The length of time a child is kept to the breast is a matter of months. There may be reasons for weaning the infant earlier; there may be reasons (in the maternal mind) for continuing to suckle it. But considerably before this time is reached a certain amount of food other than the mother's milk is usually given to it. Porridge, or in nursery language "pobbs," is well enough to begin with, but its preparation is somewhat troublesome. The specially prepared foods have been discussed before and need not be discussed again here. They consist of admixtures of baked flour, or malted flour, or malted barley as a ruler; some containing dried milk and a little fat, and a certain proportion of ground legumes (e.g. green peas). Wheat should be preferred, and the circumstances may determine which would seem more suitable than another, most of the decisions of our knowledge, being for the private physician to decide in each case. Too long continuance with any one food will pall on the palate. A variety of foods is a good thing, as to an adult—that is of "artificial" food. He who is a matter thoroughly entertained in the subject of infant nutrition, will read Easton Smith, on "The Diseases of Children."

There is one drawback to this system of feeding, and that is, the

mastication, and but little digestion, which may as well be considered here as at any other point—though not coming into operation till a later period. It is a digression, but one of such practical importance that it may be as well entertained now instead of being postponed. It is the relation of cooked and prepared foods to the development of the teeth, and the prevalence of dental caries. The jaw is imperfectly developed and the teeth are crowded. A small mouth and a light, pointed chin are now our ideals of beauty. “Muckle-mouthed Meg Murray” the ancestress of Sir Walter Scott, would find her chances less now even than they were when Scott of Harden almost preferred death to marrying her. There is, too, an imperfect development of the teeth; to all appearance they are quite right, but their dentine is of inferior quality, and when caries sets in, this inferior dentine is readily eaten up by the tiny organisms which are at work in dental caries. A bigger jaw and teeth of superior structural quality would be an advantage. The causes in operation in this imperfect development of the masticatory apparatus is now attributed to a combination of two casual factors.

1. The fact that all food is now so prepared that there is little need of that labour of the jaws which was involved in eating a piece of hard crust, whether bread or cheese. There is, in fact, comparative disuse of the masticatory organs and that is leading, by an inevitable law to degradation of the organs; the jaw is ill-developed, and the teeth of inferior quality.

2. The further fact is the higher education of the young. A child is taken early and trained as highly as its powers will permit. The brain and the dental apparatus alike are fed by branches of the common carotid artery; to encourage the flow in one branch is to deplete another; when the blood-current is flowing freely to the brain its flow is diverted from other branches. Disuse of the jaws starves the areas supplied by the maxillary arteries; and then when the demand of education comes these arteries seem specially to feed this depletion by derivation to the encephalic vessels. The brain is fed at the expense of the maxillary area, in fact.

The lack of masticatory efforts is the starting point of this morbid departure. This is a matter not to be overlooked in the feeding of children, and a hard crust (which is highly digestible) is a good thing for a child—when it has got the teeth with which to chew or gnaw it.

When a child is weaned it has paracasein milk (or ought to have this last) in liberal quantities. Now it gets milk, when this is feast-

ble, with more or less of the myriads of people, of the same kind, in the market; and if it is in danger of being overfed, it is equally in danger of being run an equal risk of being overfed in the same way. For a considerable time the infant is so taken up with the necessity of learning to wield a spoon on its own account and share the food of its mother, or of hominy—of its older brothers and sisters. Oatmeal is the staple breakfast food of the North of England and Scotland. In the more Southern areas, bread boiled in milk is the staple food. The dietary of the American Continent is rich to meet the needs of the various preparations suitable for the nursery breakfast.

Then for dinner comes the milk-pudding, prepared with meat-broth, thickened with some farinaceous matters, which has been made the acquaintance of the fire before being added to the meat-infusion; or perhaps by some potatoe finely broken up with some gravy or meat-juice. Too much of this matter is apt to produce flatulence, colic, and diarrhoea. Furthermore, when there is any looseness of the bowels, this will scarcely be counteracted by medicinal measures, so long as meat-infusions form a part of the diet. Neglect of this fact, too little known, has had its effect upon infant mortality. On the contrary, when the child is constipated, it may be well to give some meat-infusion; as well as the fig, prunes, or manna, commonly given.

Too much food brings about mal-assimilation, just as much as deficiency of food. "Among the wealthier classes, imperfect nutrition most generally arises from excess in quantity, or a too stimulating quality of food; but among the lower classes from deficiency in quantity or quality, added to scantiness of clothing, want of cleanliness and imperfect ventilation." (Combe.)

In his treatise on "Scrophula and Consumption," the late Sir James Clark pointed out forcibly, how imperfect nutrition, or mismanagement of diet in early days, lay at the root, or very near it, of that imperfect tissue-nutrition which, as we shall see hereafter, is the essential condition of these morbid states.

The milk-pudding in the nursery used to precede meat in the form; and this matter becomes of more importance as the child begins to eat meat. The starchy matters of the food, before they are salivary digestion in the mouth, the gastric juice (which is secreted as it becomes acid), ought, on scientific and physiological grounds, to precede those matters which are digested by the pancreatic juice. The nursery rule would be a good one, for many years, if it were not for larger growth. Such milk-puddings are made with oatmeal, or

hominy, semolina, vermicelli, shredded maize, sump, sago, tapioca, rice and bread and butter, as their ordinary constituents. For delicate children these different farinaceous materials need some predigestion by heat. But for healthy children it perhaps is enough to educate them to take their food slowly, so as to bring the starch into sufficiently long contact with the salivary diastase to ensure its conversion into soluble matter. In avoiding Scylla let us not run upon Charybdis.

The dietary of the nursery is ample in quantity if restricted in materials, and this is good. The appetite of healthy young creatures rarely needs stimulation, or temptation by variety beyond what has been pointed out above. Plain, wholesome food is good for children. "The same kind of food ought to be continued till after the appearance of the canine, or dog-teeth. When these have fairly protruded, a portion of soft-boiled egg may be given as an introduction to the use of a more completely animal diet. In general, solid animal food ought not to be allowed in any quantity till all the teeth have appeared and the digestive powers be fully adequate to assimilation." (Combe.) This authority holds the opinion that too much animal food leads to (1) imperfect tissue-nutrition, and delicacy of constitution on the one hand; and (2) to mental irritability and peevishness of disposition on the other; and that this combination impairs the resisting power of the organism. Consequently, such improperly fed children present lessened resistance, both in catching and succumbing to the ordinary maladies of children, as the exanthemata, whooping-cough, etc. The stronger the food, the stronger the child, is erroneous reasoning; and strong meat is not good for babes. Milk and farinaceous foods are the foods of the nursery, *par excellence*.

As to wines and alcoholic drinks, they have no business in the nursery under any circumstances. When a child is so ill as to require such stimulants, it is well to take it out of the nursery to a room of its own for the time. To give an ill-nourished child wine, is an act of weakness in intelligence or morals. If given to please a foolish mother, it is a sin—a crime. If because the medical man does not know better, then it is high time that he be referred back to his studies—there is some lee-way to be pulled up in his case.

Of far more importance is it to insist upon a sufficiency of exercise. The invention of the perambulator has not been an unalloyed good for our little folks: it may be a ready means of getting fresh air, but riding in a perambulator is not "exercise." The waxing limbs

want unrestricted movement for their development. A short of fatigue should be taken every day.

Fat is essential to healthy tissue, and how is fat to be introduced into the juvenile organism?

This is a very important matter at the present time, of growing importance. I was once in Berlin engaged to write a book. The present distaste for fat in the nursery was not only a matter of opinion, of firmness on the part of nurse, and mother, but it was a matter of fact, had to be abandoned. There is a waxing indifference to fat, which, in a visible, tangible form, is simply repulsive to many mothers, and as such is loathed. It must be "smuggled" into the nursery by the matrons put in. The hunch of bread with a stout smear of butter over the upper surface—nursery bread and butter—for nursing children is utterly unsuitable. The bread and butter should be cut thin, and the butter spread lightly over the surface ("company" bread and butter), and then this should be doubled upon itself; such is the rational bread and butter for the nursery with delicate children. Then a good piece of butter should be added to each "help" of milk-pudding, and so mixed therewith as to be lost to the eye. Bacon-fat is especially in demand in the nursery. The solid fat is most digestible, and easily tolerated by the juvenile stomach. Bread crumbs may be mixed with it, or some floury potatoe. The latter with some bacon-fat and a light-boiled egg poured over it, is a very pleasant form of food. Then the more solid fat of the bacon is also readily taken by most children; and a bit of "papa's bacon" at breakfast is a welcome treat to most youngsters. Hominy feed with fat pork is a form of food in favour in the United States. The suet pudding with the suet finely minced is an excellent method of introducing fat into youthful stomachs. If a little treacle be mixed therewith, the pudding is all the better—both as to the palatability, and as to its own food-value. Such puddings should be eaten with sugar and cream (or good milk) and with sauce. Cold rice pudding is best eaten with butter and sugar.³

However it is practically managed, a sufficient quantity of fat must be supplied in order to build up the young system. The mother's milk should never be spared. The father or mother who is a "skin-and-bone" man, who does not allow the children a sufficient amount of fat, is a failure.

³ Where cost is a matter of moment, the suet may be replaced by mutton suet, prepared by cracking the bones of a mutton carcass, and straining the fat. This dish is much more excellent.

pay the surgeon's bills of a later day for operations on joints, and scrofulous glands: to say nothing of bills for the treatment of marasmus, tabes mesenterica, hydrocephalus and phthisis: together with the bills for board and lodging at seaside health resorts. There is such a thing as "penny wise and pound foolish:" and this is a case in point. They may not grudge the cost, but recoil from the unpleasantness of insistence with their children— but, depend upon it, this is infinitely less trying than that which they will endure when disease has fastened upon their hapless off-spring—when regret is unavailing. Unfortunately, too, it must be admitted that it is just the children who reject fat who most require it. There is something more than mere fad or fancy, in the child's rejection of a piece of sweet fat on its plate, and its ready acceptance of a spoonful of fishy oil. It is impossible, indeed, to insist too earnestly upon the supreme importance of an abundance of fat in the food of the nursery. It is all very well to talk of jams and preserves. They cannot take the place of butter. They are palatable adjuncts to butter: but they never can be substituted for it! If a child can be induced to eat bread and butter by spreading some jam over it—good and well. Jam, plus bread and butter, is a very different matter from bread and jam.

When a child is old enough to handle a knife and fork, *i.e.*, between four and five ordinarily, it is allowed small quantities of meat. But milk and puddings and sweets are preferred till much later years by healthy children. When children get too much meat, under the mistaken idea that this will make them strong—that the meat of the ox will make the child as strong as an ox—as great a mistake as that of calf-lymph in vaccination being followed by the growth of horns—the end aimed at is usually defeated. The child does not grow strong. It is indeed, ill-nourished and liable to all kinds of trouble, from colic to boils and eczema. In Yorkshire, where the father likes to see his little son eat "a bit of meat," it is quite common to find gouty skin affections, with the urine depositing a sediment, in children: a condition at once relieved by a suitable dietary. Children who, from a mistaken view, are given too much meat and encouraged to eat it, and who do not thrive thereon, are sometimes recommended so much wine. It is needless to say that the remedy does not lie in this direction, but in one diametrically opposite.

Probably children are allowed too little fruit. A liberal amount of sound fruit is not only good in itself but it is a pretty certain

protection, or insurance against the children getting sick from eating fruit, no matter what its quality—at the first occasion of sickness before, “one of the most pernicious habits, in which children are indulged, is that of almost incessant eating.” (Mason.) It is to be avoided, and it is to be taught from the facility with which, for a time, a selfish child will do anything nice* procures peace. Even from infancy, therefore, children should be gradually accustomed to eat only when hunger, and not caprice, is really required. After two years of age, an interval of an hour between meals will rarely be more than enough; and to give children, fruit or bread, in the meantime, is just subtracting from the positive power of the stomach. Like almost every other organ of the body, the stomach requires a period of repose after the labour of digestion, and accordingly, in the healthy state, the sensation of appetite never returns till it has for some time been empty. To give food sooner, therefore, is analogous to making a weary traveller or workman without a halt.” (Combe.) To egg on, or tempt the palate to abuse the stomach by offering scraps of seed-cake, bun, and sweets of all sorts, betwixt meals, is a cruel kindness. Old-fashioned people, who used plain words, used to talk of a “surfeit” when too much had been eaten. We are too polite nowadays, to use such a blunt word; but the thing for which it stands denotes is unchanged. A sudden rise of temperature, often accompanied by vomiting and purging, is common with children after some undue “indulgence,” the euphemistic modern substitute for “surfeit.” It is nothing more than a natural expurgatory process, accompanied by a pyrexia, (the pyrexia), to get rid of the surplus matter.

“Children should have four meals a day; but meat only at one, or at most, two; the latter when only a small portion of food is allowed. When in health, they should have no wine or beer, except as a festive treat; no coffee, strong tea, or other exciting drink.” (Chambers.)

One rule may be trusted;—a healthy child requires no temptation through its palate. It is quite ready for its meals when the time comes. Nor, as a rule, do children need any encouragement to persist in eating. They will generally eat till they feel full. That is a child for that! Its pleasures as yet are few, and eating is its chief pleasure, *per se*, of childhood.

As regards the dietary of schools, it is now fairly liberal, though in some schools the diet-table is undoubtedly too liberal, even as to the intellectual dietary provided. Some mothers, however, still regard a liberal table as a *sin qua non*. Where children grow up in such

food in liberal quantities is required. But as to the vexed question of plenty of work and little food, and little work and much food—in other words, betwixt the well-fed dunces of rich parentage and the half-starved, bright child of poor parentage, I am not called upon to decide. Work, play, and food in nice proportion, will be found to give satisfactory results.

M. Metz, of Mattray, the philanthropist, found good food, and even wine, necessary with weak-willed boys, to enable them to form a resolution and stick to it. While of old, bread and water was the dietary of refractory children until they could obey discipline: and was effective.

Carter's paribus. I would always back a good breakfast—from a boy to a game-cockerel. A good meal to begin the day is a good foundation!

When children are ill, they require to be fed accordingly. A child will often abstain from fluids to an extent that would be very painful to an adult, without apparent inconvenience: but when ill, it requires more fluid. A child is rarely indisposed without more or less pyrexia, and then more fluid is necessary. Sometimes fluids are purposely withheld, on the impression that they would dilute unduly the gastric juice: this is a gratuitous cruelty. No child will drink cold spring water—unless when overheated by effort. In illness, its thirst is the measure of its needs. Home-made lemonade is usually enough of meat and drink for a child in a febrile state, with the addition of a little milk and lime-water or seltzer water, or the equivalent of the latter. When tinned milk is used, whether to a sick child or one in health, it is well to remember the rule laid down before about tinned foods, viz. that they should be used up as quickly as possible after being opened. Tinned things readily become unfit for food after the tin is once opened.

When a child is convalescent, usually its appetite is keen, and it takes food eagerly, and even greedily. All sorts of milk puddings, rice-milk, plain broths with baked flour or broken biscuit, and stewed fruit with cream, are good and proper. At this time a child is very liable to take more than it can properly digest, and acute indigestion often delays convalescence. Its guardians are glad to see it eat, and it relishes its food. But to quote a plain nursery axiom sometimes, "Its eyes are bigger than its belly;" which, being translated, means, that it takes more than it can deal with. Then a "surfeit" follows. Some children are gross feeders; some are dainty feeders. Those latter need tempting by various vands, served up with scrupu-

lous cleanliness. Little neat sandwiches made with potted meat, or a thin slice from the thickest part of a tongue, or of cold roast beef, and nicely arranged on a plate, will often tempt a fastidious child, and a halting appetite. One golden rule there is in feeding an invalid or sick person, child or adult, and that is, never give them too much on their plate at once; this at once impairs the rest. Rather it should be on the short side, so that the inclination is for more. Further, little appetising dishes are often devised. "Often a child will eat a small bird, as a lark or a snipe, when he turns with disgust from beef and mutton." (Eustace Smith.) Indeed, with delicate children, their tastes must be consulted, just as in an adult. "Grown-up people are permitted to give their likes and dislikes free play, and judicious wives and cooks study the tastes of their lord and master, and see that he is duly furnished with his favorite dishes; but the poor victims of nursery abstract principles, are often forced to consume things which they abhor." (Farquharson, "School Hygiene.")

On the other hand, there is "bulimia;" which is not merely the gluttony of John Reed in "Jane Eyre." "John Reed was a schoolboy of fourteen years old, large, and stout for his age, with a dingy and unwholesome skin; thick lineaments in a spacious visage, heavy limbs and large extremities. He gorged himself habitually at table, which made him bilious, and gave him a dim and bleared eye and flabby cheeks." Such is Charlotte Brontë's typical sketch of a juvenile glutton. Such a boy was studied at home in a most foolish fashion; but no doubt, boxes and hampers containing rich cakes and other toothsome comestibles, were regularly sent to him at school, till a most unwholesome product—physically as well as psychically—was the result. There is "borborygmi" not infrequently met with, after some of the manias of childhood, and in my own experience certainly, after measles. There would seem to be some low inflammatory condition of the mucous lining of the alimentary canal, some sub-mucous enteritis set up, which intensifies the normal sense of hunger. The child begs piteously for food during the intervals betwixt meals, its pangs are counteracted, with the result that no portion of the food (or very little) is digested; the child remains practically unfed; its sensation of hunger is not only not relieved, but it waxes stronger. In such cases, moderate amounts of food at proper intervals are essential to recovery; the bulimia may be relieved by some sedative, for the pain and the sensation is really pain.

In inherited syphilis, the babe is not only cachectic, but its mother cannot feed it properly, as her own milk is thin and watery, and comparatively innutritive. Here Eustace Smith, and others, have found malted foods, after the manner of Mellin's, very satisfactory.

Then there are chronic conditions of tissue-degeneration, as scrofula and rickets, one of glands and the other of bones, not rarely found together, which require their own appropriate food. In scrofula, starchy foods, even if rendered soluble, are insufficient; especially wheaten flour, which has been deprived of the bran, rich in phosphates. Here "meat, fresh eggs and vegetables" are desirable. (Eustace Smith.) We have seen before, what Liebig had to say about the large quantities of salts, especially phosphate of soda, which exist in vegetables. Much the same may be said of rickets; where phosphorus for stouter bones is urgently demanded; and where the system is unequal to breaking up the phosphate of lime. If a laxative is indicated, then phosphate of soda may be given, which is not only a laxative, acting, too, upon the liver, but also a true tissue-food. Indeed, if phosphate of soda took the place of castor oil in the nursery, it would be well for infant economies.

In the feeding of infants, a great matter is to hit the happy medium betwixt too much and too little. The results of over-feeding," as of "under-feeding" are much the same; there is imperfect nutrition in each case.

CHAPTER XII.

IN YOUTH AND ADOLESCENCE.

For our present purpose infancy may be taken to extend to the commencement of the second dentition; usually about the 12th or 13th year.

When the milk teeth are being shed, and the permanent ones are taking their place, a like change goes on in the palate, and excellent and rapid meats are more craved after. Not only that, but there is more extensive growth going on, requiring a plentiful supply of tissue-building material. The change of taste comes on gradually, and it is not till after puberty that the full craving for the flesh of animals is ordinarily developed. King Chan saw, while still maintaining the need for plain wholesome food, that "bored monotony should be avoided." He also comments "tastes change" about food. "It is astonishing," he goes on, "how many of some of these acquired idiosyncrasies become, and indeed after full manhood is reached they may be concealed, but are never quite overcome. Yet few of the minor thorns in the road lead are so vexatious to oneself and others. I shall not soon forget the annoyance of taking a young man to a Greenwich dinner, and finding that he never ate anything which swam in the waters." It is curious indeed, that he wished that that young man was cast among desert isles, with only denizens of the sea for choice, where he would have no choice. Even admitting that the dietary of the young is to be simple, he thinks it a good plan in view of the temptations of the world, if young persons would learn to eat what is set before them, so far as they are able!

the toast with water to economize butter. An old North Country yeoman used to tease me when a schoolboy by telling me—"There are only two creatures that will pay for starving—school children and wintering sheep." (In both cases the pay was forthcoming if the creatures were handed over alive—no matter in what condition.) All that is changed now.

Dr. Farquharson strongly condemns the habit of eating confectionery and sweets betwixt meals, which he terms "gastronomic nipping;" in allusion to alcoholic "nipping" or drinking betwixt meals. He also protests against the system of hampers from home. King Chambers delivers himself thus—"The gorging themselves with pastry and sweet stuff at the confectioner's, as practised habitually by schoolboys and often by girls when they get the chance, lays the foundation not only for indigestion in after years, which is its least evil, but also for a habit of indulgence which is a curse through life." It is indeed a survival of the practice of supplementing the scanty rations in the old semi-starvation days, when some such extra food was really required. But at the present day, it has no valid foundation: some cheap sugar or sweets for the ill-fed children of squalid neighborhoods is another matter.

Dr. Farquharson, who was medical officer to Rugby school, has had both the opportunity and the inclination to study the school regimen. He says authoritatively—"Little children, probably, do not require meat once a day, but growing lads working hard both with brain and muscles, require a liberal allowance of nitrogenous food, and will hardly be satisfied with the old-fashioned breakfast of tea and bread and butter. To these must be added something of a more nourishing character, and actual flesh may be frequently varied by fish, bacon, or eggs, which need not be in large quantity, but good of its kind and well prepared." (Like King Chambers he is dead against a stiff lesson before breakfast.) As to dinner, he observes, "It is not of course an easy matter to provide much variety under the ordinary circumstances of school life; but the perpetual roast joint may sometimes be boiled, or baked, or stewed, meat pies and Irish stews and various hashes may be given from time to time; and the opportunity should not be neglected of proving how good a thing cold meat really is, in spite of the social obloquy to which it has been for so long subjected." At this point it may be said that the great objection to stews and hashes lies in the suspicions of the children and their parents, that they are made of inferior meat. As to cold meat, in hot weather cold meat with potatoes (old ones

mashed and new ones boiled) and other vegetables (broccoli, for instance, or greens, forms an excellent material), and the doctor goes on, "We have already spoken of the great advantages of vegetable soups and stewed fruit, and of green salads which keep up the balance, so important to be maintained between nitrogenous and non-nitrogenous food." He notes that beriberi, or infantile scurvy is apt to develop in schools where the vegetable element is defective in the dietary. "The major forms of the scurvy disease are rarely met with nowadays, but imitative types are so much more frequently than is generally supposed; and when boys grow pale and flabby, when they readily bruise when struck, and when their gums are spongy even to the smallest degree, their parents should be aroused, and we must direct the treatment accordingly, and this will not be by drugs alone, but by ordering the consumption of such things as will supply the vegetable needs which keep up a healthy quality of the blood." (*School Hygiene*.)

When boys are but day scholars and only dine at school, they share the family fare: and the watchful maternal eye sees that they get both sufficiency and variety.

Tea is a light meal, whatever its composition, and is followed by bread and milk, or perhaps bread and cheese and some light beer as supper.

"Then something may be said of athletic training. Absurd old whims of the prize-ring still linger as traditions. Meat, but innocent of the fire is preferred to cooked meat, contrary to all physiological teaching. Green vegetables are forbidden." Past puddings and sweets are eschewed rigorously; despite the fact that Hindu wrestlers train on sweets, as Sir Joseph Fayrer, M.D., K.C.S.I., informs me,— "while strong ale or port take the place of other beverages." "A schoolboy who gets plenty of sleep, in good air, is well-fed and not overworked, and plays at games vigorously all the year round, may be said to be in a perpetual state of training, and hardly requires to adopt any particular system; but he is apt to have picked up from books or ignorant people some fantastic notions about diet, and these should be dispelled as quickly as possible." (*Farquharson*.) He admits, "It may be necessary to enter into a diet regimen on professional athletes who tend to overstrain their lives." In the diet tables of boating men which K. C. S. I. copies from MacLaren's "*Training in Theory and in Practice*" there is nothing remarkable, let alone mysterious:

CAMBRIDGE SYSTEM.

SUMMER RACES.

A DAY'S TRAINING.

Rise at 7 A.M. Exercise.	Run 100 or 200 yards as fast as possible.	The old system of running a mile or so before breakfast is fast going out, except in the case of men who want to get a good deal of flesh off. Underdone.
Breakfast at 8.30.	Meat, beef or mutton. Toast, dry. Tea, two cups; or towards the end of training, a cup and a half only. Watercresses occasionally.	
Exercise (forenoon).	None.	
Dinner about 2 P.M.	Meat, beef or mutton. Bread. Vegetables—potatoes, greens. Beer, one pint. Dessert—oranges or biscuits or figs. Wine—two glasses.	Some colleges have baked apples, or jellies, or rice puddings.
Exercise.	About 5.30 start for the river, and row to the starting post and back.	
Supper about 8 or 9.	Meat, cold. Bread. Vegetables—lettuce—or watercresses. Beer, one pint.	
Bed at 10.		

This means meat three times daily, two glasses of wine (quality unknown, probably sherry or port) and two pints of ale—probably not a thin or poor ale.

WINTER RACES.

A Day's Training.—The only difference is a luncheon of cold meat at one P.M., "with half a pint of beer, or biscuit with a glass of sherry, perhaps a yolk of an egg in the sherry"—instead of supper; and the rowing exercise at two P.M., instead of half-past five.

The dietary is that of a well-fed person, who has to earn for himself so as to earn the best wages he can get, and who is therefore far short of his work: there being thus a steady and constant hard and steadily for many hours, while the effort is not a supreme effort for a limited time, coming in only once in a while. The highly nitrogenised dietary, as we have seen before, is not essential to the rapid liberation of force, but it is essential to the pounce upon the deer, which would soon have to be left far less behind in a fair race.

Pursuing the subject (with healthy young persons in consideration at present), something may be said of the diet of girls. The regimen of girls is purer than that of boys. Boys' "bread-and-butter misses" remain much the same. The girls are involved fragility: and for a boarding-school girl to come to school appearance-checked was to call forth the derision of her schoolmates, the crude satire of the young and allusions to "dainties." After speaking of the evils of overfeeding and the unwholesome food so introduced, Combe wrote—"The opposite error of *under-feeding* is the enemy of nourishing diet for the young, as, from a mistaken view, much more prevalent than it ought to be, particularly among boarding-schools, where the system of diet is often less than the due sustenance and growth; and where consequently the outward expression of impaired health, if not actual disease, is a marked feature in the aspect of most of the pupils." Such feeding-schools is the common school management in this, and other wise countries, have the best authority for considering it as a rare exception for a girl to return home in full health after spending two or three years at an English boarding-school. It is true much of this is due to confinement, neglect of cheerful exercise, want of fresh sleeping apartments, and other depressing influences, so that that of an insufficient dietary must often be added to the list of causes. It meets with double force from the impaired digestion, and indigestion fails to ensue where the laws of health are so much violated."

Dr. Barlow, on "Physical Education," in the "Practical Diet and Practical Medicine" thinks that of "excess of diet," "the effects are more deleterious on the whole than a moderate deficiency, and that the effects of excess and repletion are less prejudicial and more easily corrected than those of inanition." To an imperfectly developed system, the digestive power from visceral overgrowth is not increased, but is due, partly, to lack of outdoor exercise, and partly to

viscera in order to secure a graceful figure. To appearance, many sacrifices are made by girls. In order to acquire pallor and get rid of the hue of health some girls take an excess of vinegar, and attain their end by destroying their digestion. Others again in order to acquire the desired hue eat raw rice, until by setting up confirmed constipation they ruin their sanguiferous machinery. It is difficult for a schoolmistress to watch and detect the vagaries of her charges, no doubt, or discover the amount of slate pencil, chalk, etc., devoured in secrecy; and with this aspect of the subject we are not at present concerned. Then the dietary of school girls is less highly nitrogenised than is that of boys; and this is desirable so long as the physical exercise is so limited as it at present is.

Breakfast usually consists of bread and milk boiled, or porridge followed by toast dry, or buttered, with rashers of bacon or meat in some form. Then at midday dinner there is meat with vegetables and milk puddings of all kinds and varieties—excellent food, as Dr. Beaumont found with Alexis St. Martin,—with stewed fruit. A sufficiency of this latter is desirable where there lurks a tendency to constipation. Tea is a light meal, and supper consists of bread and milk. The dietary is apt to be deficient in fat, and butter should be freely used. Cod liver oil is often requisite where the assimilation of fat is defective, or the palate abhors it. So much for schoolgirls.

The broad rules of the dietary for an active outdoor existence indeed are followed in a vague way. With more physical exercise would come a taste as well as a demand for more meat in the dietary. So long, however, as the exercise taken is restricted, so long it is well to have a dietary arranged accordingly therewith.

And in connection with this subject of feeding young people, two matters may be considered of much importance in early life.

Of all the food taken into the animal body, carbo-hydrates, albuminoids and fats, each and all can be found in the vegetable world. Only metabolism and oxidation—the one for tissue-building, the other for the production of heat and force—go on in the animal world. The plant builds up, giving off oxygen; the animal oxidises. The plant has no temperature, it is alleged (but where molecular activity is going on there must be production of heat),* while animals vary; the lower possessing some heat while the higher animals

* To say nothing of tropical plants dying of cold when transplanted, and so of Arctic flora in warm countries.

have a distinct body temperature of some 100° Fahr., and a slow combustion within the organism. The plant is not conscious; the warm-blooded animal is active. These two matters—food and motility, belong to the animal world. Consequently the plant possesses for these ends a circulation and a nervous system.

Carbon and hydrogen unite with oxygen, but the blood corpuscles contain an oxygen carrier for this combustion. The red blood corpuscles bear the carbonic acid to the lungs giving it off there, and take in oxygen. The venous blood laden with carbonic acid is dark in hue. The arterial blood charged with oxygen is bright red in hue. What is it which carries on this chemical interchange? It is the hamoglobin of the red blood corpuscle. Hamoglobin is a very complex substance with a formula something like this $C_{12}H_8N_4O_2So_4Fe_{11}$ with 3 or 4 per cent. of water of crystallization. This complex body has to be built up within the animal economy, being essentially an animal substance. The plant has no requirement for a blood-corpuscle. Its circulating fluid corresponds to the liquor sanguinis. Consequently hamoglobin cannot be taken from without. Nor can it be introduced into the system by drinking blood. It consists of a union of hamatin ($C_{12}H_8N_4FO_2$) with a proteid, and that proteid must be converted into a peptone ere it can pass into the blood.

Anæmia is a dearth of red blood-corpuscles, which is not curable merely by furnishing iron to the system; though of course iron is part of the cure. At puberty, especially in girls, there is a distinct tendency to anæmia. Sometimes there is even chlorosis or green sickness, with a positive breaking down of the blood-corpuscles, and discoloration of the skin from hamoglobin: just as a leaf so changes its hue to green, as the effused blood is absorbed. For the cure of anæmia—and also for its prevention—good food, containing a fair amount of meat, and fresh air, is as essential as iron itself.

So much then, for one malady linked with the later years of growth, which requires its dietetic as much as its medicinal management.

Animals possess a nervous system; also a matter not to be found in the vegetable world. This nervous system has its own work to do; its food seems a phosphorized fat, known as "lecithin," with the formula $C_{42}H_{100}N_2PO_4$. It, too, is a complex body, built up within the organism. When the brain is overworked or injured, its functional activity is impaired, as seen in "overstrain." As a matter of fact, however, the young man who has "overstrained" his brain

a person with a naturally feeble brain, which has been found unequal to surmounting the usual educational tests. Recently, when compulsory education has caught in its net all our waifs and strays, or social Arabs, it has been found, as a matter of fact, that "over-study," leading to brain collapse, is quite common. When the poor little things can get enough to eat and feed their brains, then these last are quite equal to what is demanded of them. When the dietary of girls' schools was notoriously insufficient, a break-down was far from an infrequent event. Rest from work, change, and a more liberal table, brought matters right in time.

We may suppose a failure in the supply of lecithin as the material factor of this brain exhaustion. As phosphorus is a constituent of lecithin, this element should be supplied in any form which can furnish it.* Our assimilative processes seem ordinarily to possess the power to break up phosphate of lime, or phosphate of soda, and so provide the free phosphorus required for the nutrition of the nervous system. But under the unfavorable circumstances of under-nutrition, the power seems to be more or less completely lost, and then it becomes necessary to furnish phosphorus in forms less stable and more easily broken up. The hypophosphites are available for this end: and may advantageously be added to the dietary when the demand upon the brain is great and the assimilative powers are feeble, or food is supplied to them in insufficient quantity.

Iron then, for anæmia, phosphorus for brain exhaustion:—but both with a sufficiency of food.

Indeed, in youth and growth, the maladies are mostly (except the contagious diseases and the results of exposure to cold) the outcome of insufficient nutrition. As was pointed out before, the consequences of over-feeding are much the same as those of under-feeding: because, in each case it is a question of defective assimilation. To swallow food—and to digest it, are not one and the same. It is what is digested,—not merely what is eaten—which has to do with nutrition.

The appetite of youth needs not the stimulus of food tempting to the palate. Plain foods are quite enough in most cases. With invalids some temptation may be desirable, or even requisite. But of that anon. A sufficiency of plain food, meat puddings, etc., should be provided: and when this is done, eating betwixt meals at

* "Ohne Phosphor kein Gedanke."

confectioners is not necessary, and is certainly not for the benefit of the people of either sex.

Until growth is completed, alcoholic drinks should be rejected, and only given when there is a very clear indication for their use, and this is not likely to occur in a condition of health.

CHAPTER XIII.

FOOD IN ADULT LIFE.

WHEN the period of growth is completed, the dietary chosen varies with the individual, the occupation followed, the opportunity for acquiring food, and of course according to the appetite and palate.

Some broad rules, however, may be laid down. In the first place persons of square abdomen and large viscera prefer a good substantial meal at not too frequent intervals, while persons of thin flank usually find small quantities at once, and frequently repeated, suit them best. It is just the difference seen in steam-engines, or rather locomotives. The express engine, running long distances without stopping, has a large capacious tender to hold the requisite quantity of fuel. Locomotives which run short distances, as yard-engines for instance, have small tenders; which, however, will accommodate what fuel is required in their case. If the engine with the small tender were to attempt to do a long distance its fuel would run out; in fact its tender would not hold enough fuel to enable it to do the distance. So it is with human beings. A wife of slight physique (with a small tender) tries to accommodate her meal-times to the ways of her substantially built lord, and fails signally. She must have one at least, or more meals betwixt his solid breakfast and his late dinner; she could not hold enough to carry her over the long interval, if she tried her best. She makes the attempt to do as he does, and utterly fails. She is not built on the same fundamental plan, and cannot do as he does.

Men differ too; the man of small viscera requires his meals at comparatively short intervals, because he can take in but a small quantity at once. Some savages who have to eat as they can procure food, eat enormous quantities at once, when they get the chance, and then fast accordingly. They break their fast when they get the opportunity. Our word for the morning meal is "break fast" because it breaks the fast of the night. This matter of eating, and the large and small tender, is far too little considered, where a number of persons are living together. If it received due attention

much discomfort. If nothing more, would be avoided. "Many females and delicate persons injure their powers of digestion by laying their exercise till the system is too much exhausted to carry by it." (Coombe.) When I was in country practice, I noticed that certain farmers' wives were always in after attending market. They were all of the "small tender" type. They breakfasted early, bustled about at market, took no midday meal (as usual), and so had a racking headache before they got home, and were "just eating" to use their own language. Too long delay destroyed the appetite when the opportunity for food came. Many persons cannot eat their breakfast for this reason; the long fast betwixt supper and breakfast is too much for them; such persons need something in the middle of the night. Some milk, with or without malt in it, in a jug with a lid (so as not to acquire any taint of the room), and if the weather is cold, kept warm under a "tea-coscy," is a ready and handy form of food for such persons. Or some beef tea and baked flour may be used instead. "Chronic invalids and persons of a delicate habit of body, are also familiar with the fact of the animal heat and general vigor diminishing towards morning. During the prevalence of cholera, both here and on the continent, it was often remarked that a large proportion of the attacks occurred early in the morning, in persons who had gone to bed apparently well." (Coombe.)

There can too be no doubt that digestion is much quicker with some persons than with others. And this is a matter which does not permit of more being said. It is a matter of experience with each individual.

In old days after a hard fight on a long march, or a day's hunting, came a good feast. But in our degenerate days we cannot do this. And with persons of weak digestion, it is a good rule to eat less, and to be particular about the food being easily digestible when tired. Perhaps to eat less, and take a little extra wine, is the best plan to adopt.

Meals must vary according to the individual. The Anglo-Saxon prefers a substantial breakfast. The Frenchman and the American like a cup of coffee and a roll on getting up, and then take a substantial meal about eleven a.m. They take a second good meal in the evening. It is well to have some food early, especially for those who study, or work for hours in the early morning. The first rum and milk first thing in the morning, recommended by the natives, often is of the greatest utility, and enables them to continue their

good breakfast: which otherwise they could not do—in consequence of the long fast. Times for eating must depend on the individual requirements. Families usually get on well together, because of family likeness to each other. When a stranger comes among them then somebody is apt to suffer.

The difference in individuals as to their food requirements has been noted by James Henry Bennett, who says, "In some persons the digestive process being very rapid, frequent meals are required even when abundant. For want of food at comparatively short intervals, for instance, four or five hours, sensations of sinking and faintness come on, and if the desire for aliment is not gratified, the appetite disappears, and general prostration with headache and even nausea ensues. In others, on the contrary, the digestive process appears to be very slow. When hunger has been appeased, the desire for food is long in again making itself felt. Such persons are generally satisfied with two, or at the most three meals a day. They are often not large eaters, do not require so much animal food, and can bear much better than the former, accidental or forced abstinence." Elsewhere in discussing the same subject, he writes: "The food requirements of adult human beings vary according to climate and temperature, and according to the activity of their organs: they also vary according to individual peculiarities. Some persons, even in health, digest rapidly and imperfectly the food which they consume. With them a considerable amount of it passes away undigested through the intestinal canal, and escapes with the feces. With persons so constituted, hunger soon returns, owing to the demands of the economy having been imperfectly supplied, and a fresh supply of food is required: it is as if half the coal placed on a fire had fallen through the grate. With others, on the contrary, the process of digestion is slow, and the elaboration of the nutritive materials contained in the food is complete. Such persons both require less food, and that food less frequently, inasmuch as they extract more nourishment from what they take." And such individual peculiarity is by no means confined to human beings. All who have had much experience of horses, know that some can do a great deal of work on moderate supplies of food, while others "eat their heads off in no time," in stable language, and yet are not capable of much work. And this seems to hold good of breeds of horses. Once I had two out of the same mare, which were just like their mother, in the small quantity of food they required, to do their work and keep in good condition.

It is also certain, that the temperament of the body influences the taste for and the amount of food taken. On this subject, Dr. Combe has made the following imperfect remarks "which, however, are not of quotation. "Where the constitution is of a mixed or compound of two or more temperaments, a diet composed of animal and vegetable aliment in nearly equal proportions is, in all circumstances the best. But where any one temperament predominates, the diet ought to be modified accordingly. When, for example, the sanguine constitution prevails, and is characterized by a florid complexion, great activity, strong action of the heart and blood-vessels, and a consequent liability to diseases of excitement and inflammation, the food ought to be habitually of a kind unexcited rather to soothe than to stimulate. Red meats, spices, wines, and fermented liquors ought to be used sparingly, and the principal support to be derived from refreshing soups, fish, mucilaginous vegetables, acidulous fruits, and diluting drinks. In the case of lymphatic persons, on the other hand, where the circulation is weak and slow, and all the functions are feeble, the system is benefited by the stimulus of a larger proportion of animal food, especially red meat and game; while vegetables, soups, and diluents of all kinds prove relaxing and hurtful. Aromatics and spices, however, are useful, as is also wine in moderation, and conformed with moderate exercise. If again, the individual presents a highly nervous temperament, characterized by delicacy, unsteady sensibility to impressions, and great excitability, without proportionate strength, such as is often seen in females and in men of genius, care ought to be taken not to make use of a heating or stimulating regimen. White meat, such as fish or fowl, are more suitable than the exercise of a heavy use. Spices are also hurtful, but mucilaginous meats, cooling vegetables and ripe fruits, are generally unobnoxious, and requiring that moderation in quantity is attended to, and that the course of life is in other respects regular and moderate. Where the phlegmatic or bilious temperament predominates, no particular diet, if the open air is enjoyed, more at times than the other, is necessary, unless food, than in any other constitution." When, however, we read, however, I can not get to agree with the author in his opinion, on the supposition, that he was proper, that the diet should be of a farinaceous food; a matter that will be determined. Otherwise what Dr. Combe has said is not a recommendation.

The habits and food of our colts.

culated, and form a matter of some importance, which those who meditate foreign travel too often neglect.*

Dr. Bennett writes, "Just as the stature, features, and bodily proportions assume a characteristic peculiarity in each individual during the period of growth, so the organic system also receives its stamp, the result of early habits which usually last throughout life. The habits of organic life, if we may so term them, are particularly marked in the digestive system. A man who has been brought up to the adult age on a very animalized dietary, in after life requires such a dietary more than one who has lived principally on vegetable food. Again, the one who has been accustomed until middle age to eat often, may not with comfort and advantage be able to supply the wants of his system by two meals only; and yet this system of feeding may perfectly agree with his neighbor early inured to it. Thus it is that the German, the Frenchman, and the Englishman become habituated to the food and food hours which obtain in their respective countries, and that their digestive powers often give way when they change their residence, and endeavor to conform to the novel habits of their new place of abode.

"These facts were well illustrated by what occurred during the Franco-German War (1869-70). Many thousand French prisoners were put on the usual dietary of the German soldiers—sausage meat, coarse bread, and beer. The common soldiers, peasants, and artisans, accustomed to a coarse and spare dietary at home, did very well, thrived; but it was not so with their officers. The latter became dyspeptic and ill, and in many cases their health was broken down and ruined. I have attended at Mentone several in whom the seeds of fatal diseases, principally phthisis, were then sown."

To a slighter degree the same thing was seen with the Anglo-Americans who were studying in Vienna the winter the writer was there. Himself, a hardy North-countryman, got on well enough; but nearly every other one complained of the food being indifferent, and were more or less out of sorts from it, if not actually ill.

On the other side may be quoted a case which Coombe took from Heidler. "A young woman of healthy constitution, brought up in

* When some of the inhabitants of St. Kilda's, a wild rock in the Atlantic far away to the west of the Hebrides, emigrated to Australia, it was found "they could only be kept in health by an occasional meal of the rank eggs and dried puffins they carried with them." While Finn recruits in Russia miss the birch-bark in their brown bread on leaving home,

all the simplicity of country habits, passed at once on to a less active mode of life, and to a much more elegant diet. In a short time she began to complain of irritability, loss of appetite, spasmodic sensations, and habitual constipation. Hysteria was soon added to other symptoms: her hope of becoming a mother being always deceived. An additional glass of wine, beer, and other tonics were ordered. The evil increased. The patient became melancholic, and believed that she was always swallowing pins. In the course of the year she became so emaciated and yellow that her mother, who had not seen her for eleven months, could scarcely recognize her. After an eighteen months' course of purgatives, and two courses of Marienbad water, she entirely recovered."

The same thing is commonly seen in country girls coming into towns to sedentary occupations; and young fellows abandoning an active life to wait behind a counter—especially in rooms indifferently ventilated. "The operation of the same principle is equally conspicuous in girls sent from the country to the work-rooms of fashionable milliners and dress-makers in the larger towns." With little exercise they require less food; but if the appetite keeps up, headache, sickness, biliousness, indigestion and constipation are apt to follow. Strong and healthy fellows who come up from the country to sedentary trades and occupations also require and get some unpleasant experience before they can adapt themselves to their new form of life with its novel needs and requirements. The same is found in the case of the traveller, who from a life of privation and hard bodily labor comes to luxury and dines out, while he settles down to a sedentary life to put his book through the press. He soon finds that his new life makes new demands upon him, with which he must comply, or pay the penalty. Light food of a kind likely to keep the bowels open (see a later chapter for this), is the aliment which will keep him in health.

Then comes the relation of food to climate. In tropical countries the carbo-hydrates form the staple of food. In temperate climates a mixed dietary has been adopted as the result of experience; while in Arctic regions the concentrated fuel-food has been found the only food that will sustain life amidst the intense cold. The stomach of an Esquimaux could not hold enough of carbo-hydrates to preserve him against starvation in the low sub-zero temperature. Liebig pointed out that the carbo-hydrates of the tropics contain only some 42 per cent. of carbon. The same is

of the Esquimaux, blubber and oil, contains from 60 to 80 per cent. of carbon; consequently we can readily understand the preference manifested for fat in Arctic regions—for heat-generating material indeed. There is no taste for sweets in these Northern climes, and Sir Anthony Carlisle found that even babies there had no taste for sugar, spluttering it out of their mouths, "while the little urchins grinned with ecstasy at the sight of a bit of whale's blubber." All Arctic voyagers have found the necessity for following the Esquimaux in the direction of eating fat in liberal quantities.

The importance of this matter of climate and food requirements is illustrated by the need for varied stores when bodies of men are sent out on expeditions. According to the temperature so must their food supplies be furnished—as to quality and quantity.

And closely linked with this matter comes that of the relation of food to work. It has been pointed out before that both with men and horses more nitrogenised food is required when active exertion and hard toil have to be endured. We see the farmer raises the allowance of oats to his horses at ploughing time. The Arab gives his horse barley and camel's milk, and if great effort be required, even camel's flesh. The horses used in Sahara to course ostriches get camel's milk and dried beans. (Payy.) This is a very highly nitrogenised food material, but the excess of nitrogen enables the horse to make rapid movements. The muscle itself does not wear out in exertion, as was once supposed. Traube held that the muscles burn up hydro-carbons, as fats or carbo-hydrates. Possibly the store of glycogen is the fuel of the muscle when at work. "The muscles appear to stand in the position of instruments for effecting the conversion of the chemical energy evolved by the oxidation of combustible matter into working power. While admitting that hydro-carbons are the fuel which produce heat and force, the necessity for albuminoid materials for hard exertion has also to be recognized: as albuminous food produces firm muscles, so exercise makes them hard. To sum up, science intimates that a liberal supply of nitrogenised matter is necessary to produce and maintain muscles in a good condition for work, and the result of experience is to confirm it." (Payy.)

This brings us face to face with exertion in cold climates. The Esquimaux consume much lean meat with their fat without injury. We can understand oxidation being very active in Arctic climes; and as has been said before, so long as the liver can convert

the waste or surplus albuminoids into the soluble products of excretion as well.

If the muscle itself does not wear out, what is the reason why nitrogenised food is essential for severe exertion? The answer must be found in the stimulant effects of urea, creatine, and kreatinine, the stimulant factors of beef-teen. These are powerful liberators, and so enable the system to get at its stored energy more rapidly and readily. And a certain proportion of them goes over to active effort. Navvies, lightermen, and others who work very hard have found that a considerable quantity of meat in their diet is essential to enable them to undergo their toil. Dr. James H. Bennett, in discussing this subject, says:—"I was told by an eminent railway contractor, that when some of his navvies were first taken to work on French railroads it was found that they could do in the same time double the amount of work got through by the French workmen, who lived in a much more sparing manner, and principally on bread and vegetables. The French sub-contractors were obliged, in order even partly to efface this difference wounding to their national pride, very considerably to increase the dietary of their workmen, and more especially to nitrogenise or animalise it. By adopting this course, the French workmen very soon increased their muscular powers. The materials for greater muscular development being given, and the muscles at the same time freely exercised, the organic nutritive changes became more rapid, the muscles increased in compactness and volume, and the work power increased in proportion." This reminds one of Pavy's story of a farmer who, when asked "how it was that he paid his laborers so well?" replied "that he could not afford to pay them less, for he found that less wages produced less work." The old Northern farmers always kept a liberal table, saying—"Work goes in at t'mouth."

Sir Lyon Playfair some years ago took up the subject "On the Food of Man in relation to his useful Work." He found that the "Subsistence diet" for persons who were not doing any work was about as follows:—

Nitrogenous matter,	2.76
Fat,	0.84
Carbo-hydrates,	11.60
Dynamic value in foot tons,		2.454

Diet of adults in full health with moderate exertion, as the soldiers of the great European armies:

Nitrogenous matter,	4.215	oz.
Fat,	1.397	"
Carbo-hydrates,	18.690	"
Mineral matters,	0.714	"
Dynamic value, 4,021 foot tons.			

Diet of active laborers, including that of soldiers in war time:

Nitrogenous matter,	5.41	oz.
Fat,	2.41	"
Carbo-hydrates,	17.29	"
Dynamic value, 4,458 foot tons.			

In this last it will be seen that the force-liberator—the nitrogen—is considerably increased: and that the concentrated heat, or force food (fat) is also notably increased, with a fall in less concentrated hydro-carbons.

Dr. Parkes found the food of the English soldier on home service to be:

Nitrogenous matter,	3.86	oz.
Fat,	1.30	"
Carbo-hydrates,	17.34	"
Mineral matters,	0.808	"

which is very close to Sir Lyon Playfair's estimate.

From these we can form useful broad rules as to the dietary of convicts (not doing hard labor) and persons in workhouses. While comparatively short rations will do when no labor is exacted: a more liberal dietary becomes necessary when work has to be done: and when very severe exertion is expected the dietary must be still more liberal both as to force-liberators and force-producers. The capacity of the English soldier in war has always been looked upon as closely linked with his commissariat. Prowess and food go together. Of course the question of food can never be separated from that of clothing in cold climates,—the more that is required for the heat the less is left for force-production. The native of Arctic regions is clad in furs in order to reduce his heat loss, and so lessen the demand upon his fuel-stores. Even then he needs liberal quantities of fat, (the concentrated fuel-food). About twenty pounds of meat and blubber per diem is the ordinary allowance. And Sir John Ross found that a large consumption of "Greenland food" was the road to success in frozen countries. "The quantity of food should be

increased, be that as inconvenient as it may;" and of persons who economise food by suitable clothing as far as possible.

On the other hand the Englishman who will persist in the roast beef and especially "the roast beef of old England" in tropical climates sooner or later falls ill with some hepatic trouble; because his liver cannot deal effectually or successfully with the albuminous food which he consumes far in excess of his body-wants. In Greenland he would hunger to death—he entirely burnt up "stock, look, and barrel"—in two or three days on a dietary amply sufficient for the plains of Bengal, or the Guinea coast. And *vice versa*, a Greenland dietary would pall on the appetite in tropical climates.

"There can be no doubt that a large proportion of the diseases of the digestive apparatus which are so fatal among European residents in India and other tropical climates result from the habitual ingestion of a much larger quantity of food, and this especially of a rich and stimulating character, than the system requires. The loss of appetite consequent upon the diminution of the demand for combustible material, is set down to the deleterious effect of the climate; and an attempt is made to neutralise this by artificial provocatives." (Carpenter.)

Taking his statement "that about two pounds of bread and three-quarters of a pound of meat are amply sufficient to compensate the daily losses of the system in a healthy man" (in a temperate area), we can see that considerably less than this is enough for the body wants in a tropical climate; and that much more will be required to maintain the body heat in the Arctic regions, especially if considerable exertion is also entailed.

Women as a rule eat very much less than men. They lead an indoor life, usually in warm apartments; they are smaller in frame, and lighter in weight, and consequently require less food. When women are out much in the air, either as ladies hunting or field hands, their needs are increased, and substantial meals are indicated, and usually are taken—by the ladies at least. But amongst uncivilized tribes, women often perform much of the hardest work, and probably take food accordingly.

The clerk, cooped up for many hours daily in a warm office (often up to 70° Fahr.), requires comparatively small quantities of food; the same may be said of the milch hand, and the cartman, or handicraftsman working indoors. But for the field labourer, the soldier, and the lighterman, and the soldier on active service, or a man on the scale must be allowed, if he has to keep his work up, a large amount of

persons performing little physical toil the food should consist largely of carbo-hydrates in the form of farinaceous food, with some nitrogen such as a fish diet could supply, and a little animal food. Of course, where there is any tendency to tissue degeneration (phthisis or scrofula), a certain quantity of fat is desirable. But with all this calculated, we must still reckon with the disturbing factors, the "palate" and the "appetite," which will rule the roost with many persons.

In a vegetarian dietary, the legumes (rich in nitrogen) take the place of meat, while oil supplies fat; and mineral salts are furnished by vegetables, as cabbages. No doubt a great deal of excellent work has been done by vegetarians; and a great many people would do wisely by becoming semi-vegetarians, at least, especially in warm weather—the writer's own personal practice. But then there are to be heard so many other voices, including those of the housewife and the cook than that of Science on this topic!

Then comes the matter of the opportunity of acquiring food. This is a matter which can scarcely be discussed at length here. It is a well-known fact that stockmen and miners have very often to live on undesirable food, as "damper." Flour cakes are indigestible, and pancakes have been the ruin of many a stalwart fellow, who ate them because he could not get anything else.

The subject of the food requirements of adults may fitly terminate with some considerations of the tendency of the age. We have not got the internal arrangements of our ancestors, and cannot deal with the quantities of food they ate apparently without ill effect. And this is specially true of meat and alcohol. There is a widespread movement towards "Vegetarianism" and "Teetotalism" among the Anglo-Saxon race at least; and this must have some acting agent. In a little work published some time ago, ("The Food we Eat") the writer speculated as to the dietary of A.D. 2000. What is put there in a light and jocular manner is however what he seriously believes will be the case, and what we are coming to. But of course the spirit of the axiom, "the future must take care of itself" prevents anyone from taking any steps for their potential descendants. For the bulk of persons a lighter dietary consisting largely of cooked farina (otherwise predigested starch) and milk will probably be found the best. But for those engaged in literary work some doubt may be expressed. "The grim old rufflers of the tongue and pen" were eaters of meat and drinkers of wine; and Carpenter states;—

"And while on the other hand it may be freely open to the advocates of 'Vegetarianism' that a well-selected vegetable diet is capable of producing (in the greater number of individuals) the highest physical development of which they are capable, it may be on the other hand affirmed with equal certainty, that the introduction of a certain proportion of animal flesh is in no way injurious, while so far as our evidence at present extends, this seems rather to favor the highest mental development."

So much then for the food wants of grown man under the various circumstances of occupation and environment.

CHAPTER XIV.

FOOD IN ADVANCED LIFE.

"A MAN loves the meat of his youth that he cannot endure in his age."

So said that most remarkable observer, Shakespeare. How he ever found time and opportunity to observe all he did, is a marvel to ordinary mortals. This observation too is perfectly accurate.

There are a good many reasons for the inability in age to cope with the dietary of youth. We saw in the last chapter the relations of the food to the individual, his work and his environment; and can apply its lessons here. As age advances the energies wane. The capacity to take exercise or undergo toil is enfeebled, and the inclination to do so also lessened. A man pleads he "is not so young as he once was." His limbs are stiffer and his wind is not so good. Some certainly retain their vigor till a very old age, and with it their digestion and their gastronomic ardor. In my days of early practice in the North of England, there was an old gentleman of the old school, and the habits and practices of the Regency. From early morn till dewy eve he sturdily walked throughout the day in order that he might enjoy his late dinner. His bell was his god; and he was an earnest faithful worshipper! It was not a lofty creed—but it could inspire conduct; and that is the highest terrestrial effect of a creed. He walked, he ate—and what is more, ate with impunity; and died a nonagenarian. After all it was a piteous sight to see the old patrician tramping along in all weathers with his dinner before him—an object, however, which could lure him to undergo all the inconvenience of exposure to bad weather in a tempestuous locality. His ambition, such as it was, was gratified.

At first "the turn of the hill" is scarcely noticed, and perhaps the only difference is that the individual does not come twice to the joint. Probably, too, the once cherished draught of ale is no longer so keenly relished. A slighter excess of alcohol is felt next morning than of yore. Perhaps gout has induced him to leave off port wine. But as years creep on the alteration becomes more marked; the ap-

petite generally is impaired; the palate loses its acuteness, and the skill of the cook is no longer so successful; the sense of taste and the appetite; but as the latter falls, so the former becomes more particular about his dishes.

The teeth, too, fall; and the strong meat "grinding machine" is no longer so easily masticated. Consequently, soups, broths, place of the joints, the steaks and chops, once so eagerly consumed. Spoon-meat is found more acceptable in every way. More condiments are required, and highly seasoned dishes find favour for more reasons than one. Flatulence is the bane of advancing years; and for this the cook can do much. There is a tendency to constipation; and that too is a matter for which much can be done in the kitchen.

Before proceeding further it may be well to quote from "The Diseases of Advanced Life" by Dr. Melachian, on the anatomical changes which go on in the viscera in age:—"Participating in the general wasting of the organs and tissues, the stomach and intestines lose bulk and become thinner in old age. Their glandular apparatus is also atrophied. Many glands seem to have entirely disappeared. The wasting of the tunics composing the stomach and intestines is more obvious in the duodenum, jejunum, and cecum. In some cases it is carried to such a degree as to admit of the contents of the intestines being distinctly seen through the attenuated structures. In striking contrast, the larger intestines occasionally preserve their natural thickness, chiefly through a compensating hypertrophy of the muscular coat. The mucous membrane is usually poorer than in the adult, and generally acquires an ash-gray coloration. In the stomach it is often traversed by enlarged vessels, assuming a varicose character, which become more numerous in the lower portions of the intestinal canal, and are particularly conspicuous towards the termination of the colon and rectum."

Such anatomical details tell us that the functional activity of the digestive organs must be much impaired in age. The gastric secretion must be crippled by the stagnation of the blood vessels. The glands of the small intestine are atrophied. Consequently, nutrition must be less perfect, and the food must be of a more readily assimilable character. No wonder, then, that the old man

"The fair remembrance yet of his youth is left."

shrinks, and that impaired nutrition leads to emaciation.

Then again, a return to nursery food is food in the form of stewed fruit. The toothless jaws cannot deal with the firm apple as can the sharp unblunted teeth of youth, and consequently it must be baked or stewed; and very good such fruit is for elderly persons.

There is a medicinal matter connected with stewed fruit. When cane sugar is added by the cook the resultant product is apt to turn acid in the stomach with many persons of middle age and advanced life. In such cases it is well not to mask the acidity by adding commercial sugar, but to neutralise it by an alkali. This leaves the natural sweetness, due to the levulose sugar of the fruit, which is quite pronounced enough for elderly palates. The amount of bicarbonate of soda, to be added when the fruit is put into the oven, is as much as will lie upon an English shilling, or a German mark, or an American quarter dollar bit, to the pound of fruit. This is quite enough even for unripe gooseberries. Such stewed fruit is excellent, and will often agree, where stewed fruit as ordinarily prepared (*i.e.*, by the addition of cane or beet sugar) turns acid.

This matter will be referred to again when the dietary of the gout is considered, only there it is well to substitute the bicarbonate of potash for the soda.

Such stewed fruit with cream forms, or ought to form a regular staple in the dietary of old age.

By keeping the principles which should guide us in the selection of food, well in view, much may be done to preserve the aged organism in health. These matters of mineral salts, fruit salts, and vegetable or meat salts, are of much importance; and more attention may be paid to them with advantage.

Milk puddings of all kinds and made with flour, maize flour, broken biscuit, rice, sago, tapioca, and semolina should form part of the dinner, at least, with all old persons. Porridge, or some similar dish of oatmeal, hominy, or any prepared cereals should always be a part of breakfast. The rule that all farinaceous matters should be exposed to a high temperature for some time before being used for the cook's purposes should no more be forgotten with culinary preparations for old folks, than it should be omitted in preparing nursery food.

The digestive ferments (like the rest of the body) are not so active in age, and so all food should be carefully prepared; and pre-digested starch is easily obtainable. As there is a difficulty often in

maintaining the body temperature with old persons the quantity of soluble carbo-hydrates must never be forgotten.

There is a prospect of milk sugar soon being in the market, and this will supply a need to delicate people which is very desirable. It can be added to tea, or to mineral water, or be prepared as a lemonade, or even be added to meat tea, not being too sweet to offend the palate when taken in such company.

When elderly persons dine out they often find the condiments of their hosts deficient in condiments for them, and consequently they are troubled with flatulence. They miss their wonted condiments. "A good old-fashioned plan of adding condiments to food without offending the palate was to make a peck or two of bread-crumbs, and cayenne pepper at table; and, so guarded, the circulation worked well. Some gourmets used to carry a small bottle of Negro pepper with them perpetually, so as to be prepared for any emergency in the way of lack of flavouring when dining where much seasoning is eschewed." (The Diseases of Sedentary and Advanced Life.)

All soups and all fish (except perhaps salmon) are suitable food for the old, so are oysters and clams; but lobsters and jellied shell fish are apt to disagree,—probably because imperfectly masticated. Heavy solid meats should be eschewed to a great extent. Light food as chicken, and rabbit, sweetbread, giblets and small birds and game are good. Sheep's head well cooked is admirable. Tripe too is excellent. Meat prepared by mincing or hashing (fresh meat being preferable to cold meat) is also good. A nice dish is prepared by stewing some hashed meat and adding a little flaxy potatoe with some pepper and salt: which gives vegetable matter as well as animal food in a readily digested form.

As the taste for sweets does not return with age, in the case of many old persons something of the kind of a relish is obtained. Ported meats of many kinds, beef, game, venison, are well liked. *Pate de foie gras* is excellent, and might be more used with advantage, to say nothing of the gratification of the palate, than it is. Livers indeed, fried and roasted, may be eaten hot or cold. The *Leberwurst* of Germany is a relish to many a table and furnishes a variety of food, too. Carriage has its claims to notice. The new and old fried on toast may be tried.

Oyster soup, a red mullet, a sweetmeat, a quince, a peach,

millet, followed by a digestive biscuit and butter with a glass or two of generous wine as Bucellas or Madeira, would form a "company" dinner for old folks,—to which the writer would have no objection to be invited.

Or a basin of whole beef tea with broken biscuit, followed by a milk pudding, forms a luncheon not to be despised.

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

1	2	3
4	5	6

...

and so there is not the acute craving for fluids. The food may consist of (whole) beef-tea, with a little broken biscuit in it, or milk, or milk and seltzer water; or milk-puddings—of malt extract, of stewed fruit and cream; and later on, any kind of soup. Then a little fish, a sweet-bread, or chicken, and so on gradually until the ordinary diet is ultimately regained.

But with fever present, as in the exanthemata, typhoid fever, or acute inflammation of the lungs, something more of the nature of a drink is desirable. The temperature is high and the fauces are dry, and the thirst is urgent. Consequently what is wanted is *Potus Imperialis*, Apple water, or Tamarind water, or Apple Toast water, or Apple Rice water, or the juice of fresh fruit as the red currant, the black currant, or the mulberry, diluted with water; all pleasant refreshing drinks; the nutritive value of all being very small. Rice water and barley water give something more nutritive than plain water and may be flavoured with any of the above. Home-made lemonade with a thin piece or two of the rind to give flavouring is excellent. Where cane sugar disagrees it may be made with malt extract, or Mellin's Food, or if the malt flavour is disliked, with milk sugar or levulose. Malt extracts with effervescing water are good. They supply maltose (grape sugar), soluble dextrine, some soluble albuminoids and some salts. Consequently they form an admirable food in pyrexial states. Then come the meat-infusions, as beef tea (as ordinarily made), mutton broth, veal tea, and chicken broth and eel broth, all pleasant stimulants, but lacking in food-value; and, unless some baked flour be added, not to be counted as food, but as pleasant beverages.

Milk may be given as whey, or diluted with Seltzer or other water. A great matter to be borne in mind with milk, especially in the case of typhoid or enteric fever, where the bowel is weakened by ulceration, is that it curdles as the first step in its digestion. When it curdles too firmly the curd does not fall to pieces, but travels along the bowel, a hard irritant mass, apt to lacerate the weakened bowel. This is no imaginary danger born of apprehension but a grim fact. At the London Fever Hospital a glance is always cast at the stools in enteric fever, and if hard white pieces of milk-curd are seen in the motion, measures are taken at once to remedy the hard curdling. A quantity of effervescing water is added, sometimes with some baked flour: so that too firm curdling shall not occur. Lime water is also useful. Sometimes a more pronounced alkali is indicated, as said before when milk was under

consideration. It is not enough to give a patient food in quantities; judgment is also required to select the food that will do no harm.

How much actual food is being gathered up by the body can quickly be gathered from a glance over the face of the patient, and this list should always be kept in mind.

There are other fluid foods which can be given when there is gruel, or oatmeal and water in view, none of which is so far from being used and being beneficial with care. A strained, served, subacid fruit as black or red currant juice, or orange juice with water, and on cooling furnishes a reasonable amount. A little honey or milk sugar may be added to give some more food value to the drink.

By ringing the changes on the same vegetable, fruit, or other easily masticated food, when an additional amount of food required is furnished to the patient. The grapes, for instance, are a good food of the body, which must be strong enough to meet the demands of the high temperature.

By similar alimentation the wasting process of the body does not become so reduced, and consequently a true anorexia does not develop. Where food-tarces only have entered the mind of the patient in apprehension that it is a sustaining food, the patient must look on its own resources, and too often the patient falls victim to the disease not dug for him by disease, but by the ignorance of those who substitute the nurses and attendants. But this matter will be taken up more fully further on.

At this point something may be said as to the amount of food in acute pyrexial conditions. We have good reason to believe that the digestive ferments are enhanced and present in a large quantity. Consequently we give such things as milk, cream, and soups, matters which have been predigested into a soluble form. This relieves us of any anxiety about the secretions of the stomach. We can furnish to the body its own food, neither of them presenting any difficulty; what is perfectly soluble presents no difficulty, and the food the body must have.

Fat is never relished by the patient, and is therefore never given so that it is carefully skimmed off the cream of milk. The fat supplied in the milk which is the best source of fat, is in a high state of solubility. However, we do not know. Nothing is yet known of the fat in the blood just.

Then comes the subject of the amount of food.

buminoids for histogenesis. This is a difficult matter and must be judiciously considered. Hippocrates insisted upon nutrition in fevers, and used barley gruel. Then came other views until the French medical men went to the other extreme and actually starved their patients to keep down the fever. Anything more abominable can scarcely be conceived. Under-feeding patients in febrile states however continued till Dr. Graves of Dublin set his face against the plan, and insisted upon more alimentation being given. The "feeding of fevers" was what he taught. This plan now universally obtains. "The great art of duly nourishing fever patients consists in giving a frequent, almost continuous supply of liquid nutriment, containing very soluble aliments in a dilute form" so says King Chambers. And now a very unpleasant task must be performed, and that is a searching criticism of what the worthy Doctor says next. "The physician sees that a large supply of nitrogenous material must be wanting. The nitrogenous tissues are devitalised and drained away in a disproportionate excretion of urea and other organic compounds, and nothing is taking their places. Shall he act antagonistically and try to stop the passage of urea by the kidney? I do not know exactly how he would set about it; but I do know that if he succeeded he would do positive harm: for the worst cases of fever are those in which metamorphosis is active (as shown by the heat), while the excretion of urea is arrested as shown by the lightness of the urine), they resemble cases of uramic poisoning from diseased kidneys. The other principle of treatment which I noticed in my introductory lecture would not perhaps be so directly injurious, but common sense would still allot the palm to restoration here. Let it be your chief aim to supply that which is passing away—nitrogenous tissue." Now the first sentence and the last sentence of this quotation point unmistakably to this, viz.: supplying to the body matter for histogenesis, for the repair of the wasting tissues. On referring to the lecture from which he quotes it seems he advocates "two or three ounces of milk every hour." He is not as lucid on this point as he usually is, but we must take the statement as it stands, viz.: that the "chief aim is to supply that which you clearly see is passing away—nitrogenous tissue." In the absence of context this means—If the sentence means anything at all—that we ought to supply nitrogenised matter to take the place of that which is passing away.

Now to this I demur. The tissues are melting down under the high temperature, and the danger to life is that the blood is becom-

ing surcharged with nitrogenised waste, and the system of excretion ("the typhoid condition") is set a-going. The system of excretion is the waste in the blood which is the danger-point of the disease. If we try to lower the temperature, to peptonise the food, to stimulate and keep the kidneys in action to get rid of the waste, and to get rid of the toxic matters. Now to give ample supplies of tissue-food, to give ample supplies to meet the waste going on, is a distinctly dangerous thing. It is, it appears to me, as tending to increase the amount of the waste, the matter of excrementitious character in the system. Products of retrograde metamorphosis are not solely furnished from the waste; they come also from the *tissue-examples* of our high-grade food, and to give ample quantities of animal foods is to give plenty of evil. Of course it is possible to contend that these products are never peptonised, and that they merely drift along the alimentary canal till they reach the point of excretion. But then we need not be certain that such is the case. Certainly if peptones reach the excretion, there is no such escape. If peptones reach the excretion, as is mentioned, and pertinently questioned, if the liver, which is the excretory centre, can carry on metabolism and convert some of the waste-matter conveyed to it into serum-albumen for its own needs. A downward career into kreatin, kreatinin, urea, etc., types, and the other products of retrograde metamorphosis (some of which are toxic) seems infinitely more probable. To give a large amount of tissue-food in ample quantities for histogenesis during pyrexia is, I opine, a mistake. We cannot be certain about it, but to repeat it is all Lombard Street to a Chinaman getting lost. Instead of saving histogenesis we only increase the risk of increasing the typhoid condition.

It is a matter which cannot be demonstrated; it is a matter of opinion and conjecture, at least, at the present state of our knowledge.

(From some observations made by Dr. Ord and Dr. Barlow at the opening meeting of the Medical Society of London, Oct. 18, 1885, it would appear that there is a certain relation between histogenesis and pyrexia. Where histogenesis is going on, the lowered temperature. This absorption of heat is due to the process, and the heat lost in histogenesis is added to the heat lost in the process; and so swells the pyrexia. The conclusion is that if there is no histogenesis going on in pyrexia, the heat lost in the process is on foot. The subject is one to which some further investigation.)

Probably Dr. Chambers has failed to express himself clearly; for all the men with whom I have interchanged ideas on the subject see it in the light in which it is put here. It is now widely recognised that the estimation of the bulk of urea passed in pyrexial states—once regarded as so important—is fallacious; because there exists no means of telling which of the urea is tissue-waste, and what proportion of it is derived directly from the nitrogenised elements of the food which have never passed through the stage of tissue.

Of the importance of “feeding fevers” no one now entertains any doubt; but of the desirability of larger quantities of albuminoid matter in pyrexia the gravest doubts may honestly be held and expressed. We know that the muscular structure wastes and breaks down into *débris* in pyrexia; we know that subsequent repair takes place by the development of young fusiform fibres within the sarcolemma; but this last is a matter of convalescence when the appetite is keen, especially for animal food. There is the pulling down of an existing building, and the erection of a new one on the same ground; but the pulling down precedes the building up—the two cannot go on together side by side at the same time. Histolysis and histogenesis go together in health where balance rules. When the normal processes are perturbed, upset by pyrexia, histolysis is augmented, to be followed in turn by rapid histogenesis. But I maintain, a rapid histolysis is not accompanied but followed by a swift histogenesis, or tissue-repair. It is the difference of a pair of horses side by side and tandem.

It is well for the reader to determine in his own mind what are the facts—or perhaps rather the probabilities of the facts—otherwise in well-meant endeavours to do the fever-stricken patient good we may be but reducing the chance of survival; and but adding fuel to flame, by still further lading the blood with nitrogenised waste.

Dr. James H. Bennett has something to say about “Defective Nutrition from the Existence of Acute Disease,” to this effect:—“In acute disease the functions of nutrition are partially suspended, the digestive powers are weakened, or even for a time arrested; hence the loss of appetite, or even the absolute loathing of food which ensues. The digestive organs being unable to transform and elaborate food, an all-wise Providence preserves them from the labour they are unequal to perform, by destroying the desire for nutriment, and rapid emaciation follows, the more rapid the more complete the inability to digest food. The process of organic destruc-

tion or disintegration inseparable from the use of it. If itself a condition of life goes on, accelerated perishing goes on; but the material to repair the loss not being affected by the dissipation it wastes in volume. Moreover, as the materials of combustion for the generation of heat are not coming from the digestive organs, the economy is obliged to re-absorb the fat of its own fat, —deposited, no doubt, in its cellular or adipose kind of reserve tissue. Thus it is that after an illness of one or three weeks the emaciation is sometimes extreme, more so when from the nature of the illness it has been impossible for the patient to digest any kind of food.

"In continued fevers, in which the inability to take and digest food is all but complete, and which may last for weeks, it is often found absolutely necessary to give wine, or other alcoholic beverages, in order to afford to the organisation the elements for heat-generation, or organic combustion, and to retard the process of destructive metamorphosis of tissue. In the absence of such aid, the extremities and then the body gradually become colder and livid, ensues, as much from cold as starvation. In this case the lamp of life simply goes out for want of fuel; like a coal or wood fire that has exhausted all the materials of which it is composed, and which expires for want of their renewal."

But all along it is "fuel-food" not "tissue-food" for which Dr. Bennett contends so eloquently, and what is about to be said is in contention on the same lines.

Imperfect supplies of "fuel-food" to the wasting organism have the effect of permitting great emaciation, which more liberal supplies would prevent; and on the present point of treating acute pyrexial maladies we certainly do not prevent wasting, and its consequential weakness, as we might. Beef tea (however good) contains practically no fuel-food, while milk contains but a comparatively small proportion of milk-sugar. Consequently our patients are apt to die, as they did under the French physicians, who deprived them of all food and called the starvation period *faux jeûne*; or openly starved them to death; while we allow them to eat, and give them insufficient food. The difference of course is the same as between the Frenchmen. We stand by the strong argument between the falling strength, milk-graveyard of the resistance of the powers, and do nothing to prevent the inevitable exhaustion. We see the wasting going on, and do nothing to stop it; we do "fuel" or nothing. We leave the patient to starve.

and unaided. We certainly give some alcohol—a readily oxidisable fuel-food—but we do not act efficiently. If the subject of exhaustion were only more fully and completely considered, we would know how to help and succour the sinking organism. At the recent annual meeting of the British Medical Association at Cardiff (1885) I read a paper entitled “When a Patient dies of Exhaustion what does he die of?” He dies because his body-store of fuel is exhausted; because we do not supply to him the grape-sugar which is the normal body-fuel, and his reserve stores of fat cannot be got at sufficiently rapidly. We starve our patients into exhaustion, and bemoan the fact; and then leave them to their fate, without doing anything to save them from it.

It is a grave and serious charge to bring against ourselves, and the matter, as the President of the Medicine Section (S. Wilks) observed, the matter is one of the very greatest importance. Do we, or do we not allow our patients to die from insufficient feeding? Not one person present at the reading of the paper disputed my proposition. It is true then that we do not feed our sinking patients as we should. We ought to supply to them grape-sugar in free quantities in order to conserve and economise the body stores. The beef-tea ought to contain well-baked flour (much of it soluble dextrine), or even milk-sugar or maltose, which is not too sweet for most palates. We should give lemonade made with milk-sugar, or levulose. Acidulated drinks with grape-sugar, or its equivalent ought to be given. Water is wanted to allay thirst, and drinks should be so made as to have a distinct “food-value.” If the patient cannot take such drinks and will only drink water, then of course we are not to blame. But when the patient does not get the chance of such nutritive drinks then that patient is a victim to ignorance—a sacrifice simply. It is not a matter for demonstration, it is one which reason can determine; and if after this outspoken protest any reader of this work does not feed his patient properly by the light of science he is a murderer! There are sins of omission as well as sins of commission, and to allow a poor patient to die unfed, is like allowing a man falling overboard to perish by not throwing him a life-buoy, or a rope when those are ready to hand. What should we say in this last case which would not apply to the first? Of course so long as we did not know what it was the sufferer needed there was no blood-guiltiness. So long as the different forms of grape-sugar were not to hand we were not blameworthy. But now that we know what is required, and, further, have the materials to

hand there is no palliation of the effect on food and drink. And the outcomes of systematic lectures on dietetics on the part of the proper and efficient feeding of patients in acute diseases, especially in pyrexial maladies: so that they shall not become exhausted, but exhausted further than we can possibly help. "Keep the hands but wisdom fingers" and now that we realise the importance of supplying grape-sugar to the economy in its state of transference as an efficient aid to the sick man battling with disease, and not going to exhaustion if we do not interfere.

What was written by King Chambers and quoted at the commencement of this chapter should burn itself into every reader's memory, viz.: that many sick persons pass into a critical stage from which many never rally—because they are insufficiently fed.

What was the denunciation on those who gave no food to the hungry, no drink to the thirsty? "Depart from me, ye cursers, into everlasting fire." And every man who has allowed a patient to die of exhaustion without having made a stout fight for him, deserves like Cain to carry the brand of murder on him for the rest of his days.

Even when the boathill for food is pronounced, it still remains possible to administer enemata of grape-sugar with meat broths containing salts and the stimulant extractives of meat, or when the typhoid condition is pronounced such feeding might still be continued. The patient must be supported in the hour of trial, and if this be properly done we may fairly expect a good percentage of cases to rally which now go on from bad to worse till death by exhaustion closes the scene.

I now add the Dietsaries in use at the London Fever Hospital, from which it is clear that the patients there are not starved but have ample allowances of food provided for them.

The whole subject of the feeding of persons acutely ill and in critical conditions is one on which some instruction is desirable. Milk does not agree with all. Beef-tea is a stimulant and not a food. What the patient needs is soluble carbo-hydrates, which may be given in various forms, which are not particularly expensive. What the Doctor wants is information on the subject of the matter. And if the Royal College of Physicians of London could to do this its history put its hand to a useful work and give the world some evidence of its utility, some excuse for its existence, instead of lumbering on and merely biding its time for a more useful and good thing. I fear it is vain to expect such activity from the Bo-

it seems the fittest body to undertake such a work. If it would only have one question on Dietetics in its examination for the superior qualifications of its Membership it would be taking a step in the right direction.*

* From a number of experiments performed of recent years it seems that the digestive fluids and their several ferments are gravely impaired in acute disease. This fact adds strength to the views advocated in this chapter.

CHAPTER XVI.

FOOD IN CONVALESCENCE.

"In the management of convalescence the patient should never be permitted to sit out of bed till the strength be considerably advanced. It is better that restriction should be imposed a little too long than that any risk of relapse should be run.

"The next point is of still greater consequence—the proper regulation of the diet. In our experience, by far the greater number of cases of relapse take place from indiscretion in diet. It should also be strongly impressed on the convalescent that it is as necessary to guard against the quantity as the quality of food, particularly when there has been gastric irritation in the progress of the fever. The stomach may be able to digest and assimilate a limited proportion of food, but the indulgence of an extra ounce or two may induce oppression and a renewal of the fever. This organ in convalescence partakes of the external muscular debility, and the convalescent may as well expect to be able to carry a heavy load on his shoulders, as to digest an undue quantity of food, even of a suitable kind." (Tweedie, Article "Fever" in the *Cyclopedia of Practical Medicine*.)

When the worst is over, and the critical period passed, the prostration is great. Care in the administration of food and the prohibition of stimulants is still required. But shortly the sores disappear from the teeth and lips, the brown fur clears off the surface of the tongue, and with these comes some return of the appetite, and a relish for food. The weakened condition of the organism, both in its whole and in each component part, must be borne in mind; and in all illness and debility the stomach is involved, and is functionally impaired; in other words, all sick and weak persons are dyspeptics, and must be fed accordingly. Small quantities of milk and seltzer water, of whole beef tea, of meat infusions with soluble carbo-hydrates are required at frequent intervals.

And there is one point to be attended to in the management of convalescence, and that is, not to rouse the patient too early, in order to be fed. The heart structure is extensively diseased, and

acute and still more by sustained pyrexia, and a great many of the muscular bundles are reduced to a mass of *debris* in the histolytic process which has gone on, so that the heart-wall is weak, and to raise the patient suddenly is to throw the weight of the blood-column in the head and neck abruptly upon the heart. Under these circumstances the left ventricle is unable to bear the weight so thrown upon it, and comes to a standstill in diastole—flaccid and distended with blood. Even after the patient has recovered sufficiently to walk about the room this danger of heart failure is not fully past; and after the high pyrexia of relapsing, or famine fever, it is not unusual for the convalescent to drop suddenly to rise no more—the weakened heart having stopped abruptly.

The appetite for food steadily increases, and must be met. Small quantities of milk puddings of very digestible character may be given; such as those made with broken biscuit, or with shredded maize, or cooked hominy, or indeed any farinaceous matter which has been already exposed to a high temperature. Milk or cream with cold coffee forms an admirable nutritive beverage. There is often a considerable amount of thirst lingering, and this calls for beverages, which can easily have a food value given them either by some form of maltose or grape sugar, or by adding some baked cereal. Gruel made with baked flour, or baked oatmeal is good. The lactated foods can be used, and if need be, may be flavoured by red wine of any kind, or by some other flavouring agent. Lemon juice suggests itself as a pleasant acid assuaging thirst effectually. Sago and rice milk may be added, and oatmeal and hominy porridge make a change. Blanemanges suggest themselves, but should not be made of raw or uncooked starch. The salivary glands still feel the storm which has swept over the organism, and have not regained their capacity to digest starch. To forget the fact is to run the risk of filling the enfeebled stomach with undigested starch—just about the most effectual means of upsetting it that could be devised. Dr. Tweedie warns us against errors in alimentation in convalescence, and this is one of the most common. A little later on, an overload or surfeit is apt to occur from the appetite being too keen, and the quantity taken at once too much for the stomach. A greedy feeder is most liable to this last risk of relapse; but all convalescents may experience a “back-cast” from a quantity of undigested starch finding its way into the stomach.

Then the patient advances to some white fish boiled, to a sweet-bread, to some chicken. It is not necessary to tempt the appetite

by condiments, the palate declares for simply dressed food; the plainer the food the better. The palate requires no artificial stimulation; the desire for food is keen, and eager enough to be gratified without any whetting. It is not like the invalid's case, in which the appetite is small and fastidious. Young convalescents, upon leaving disease, can generally eat anything offered to them; and consequently those in attendance upon them should exercise judgment in the selection of the viands supplied to them. They eat and go to sleep,—to waken up with a craving appetite.

There has been a severe demand upon the body stores of food which have to be built up again. There has been extensive destruction of the muscular tissues by a rapid lysis, the tissues melting down rapidly under a high temperature. All this has to be met and repaired; and the appetite is keen, and the digestion active and rapid. Nevertheless it is well to commit no error; as all persons familiar with disease and convalescence know that a relapse is followed by a slower rate of repair. Once a "back cast," and the process of repair and the progress of the convalescent are a ke slowed. A maxim with Sir Frederick Roberts in his Afghan experience was, "When you have once got an Oriental on the run, keep him on the run." If his flight was not kept up he might rest somewhere where it might not be so easy to dislodge him; consequently, it was best to keep him going. So it is with convalescence; it is best unbroken and uninterrupted. Once a check, and the onward progress has again to be initiated. And all initial movements start even in a Great Northern Express. Every stop involves a fresh start. Carefully then, and with caution liberal supplies of food may be given to the convalescent. But it must be borne in mind there are convalescents and convalescents. There is the robust hearty youth who has had a smart attack of enteric fever; as soon as his appetite returns he can eat anything, a sheep in its wool and all. But there is too, a delicate girl recovering slowly from an attack of scarlatina, with kidney complications. She is a very different convalescent: there is the injury done to the kidneys to be kept in mind, and never forgotten. The kidneys are a great factor in the excretion of nitrogenised waste. While they are still feeling the effects of the disease which has passed over them, they are called upon to great demands upon them; and this function, delicate and delicate, is overlooked or even underestimated. Such caution in the giving of food alone are safe here. The hungry youth who eats like a pig, and the sheep is skinned would be torture, and a relapse would be the

meals, sparingly composed of animal food, which would be proper in the other case, —while aliment suitable enough for him would be fatal to the girl in all probability. “What is one man’s food is another man’s poison.”

The importance of bearing in mind the function of the kidneys in the dietetics of convalescence is illustrated by the following case. In my early days of village practice in Westmoreland, a case of nephritis came under my care,—a typical case of acute Bright’s disease, not post-scarlatinal. By suitable measures convalescence was secured. I was but a young practitioner, and though the case was doing well somebody thought it might be doing better. So an older man was called in, who thought I might allow some more food and administer some steel. Accordingly, this was done, and for a day or two all went well. Then came a change sudden and rapid; in spite of all that could be done uræmia set in, and the girl perished. That case has been one corner-stone in the building of my medical experience. Regret was unavailing; and I went on a sadder but a wiser man.

In like manner after enteric fever it is well to remember that there has been ulceration of the bowels, often extensive, sometimes deep, and that hard pieces of food—as unripe apple, or insufficiently cooked potato—might readily cause a perforation of the weakened bowel. My repertoire contains no warning case telling of forgetfulness here, which could be placed at the reader’s service. A medical experience involves a certain element of patients sent to their doom by some sin of omission or of commission, and mine is no exception.

But my experience can tell of a very instructive case ament convalescence; and the recital may help some reader in a difficulty. In my early days an old peasant woman was seized with complete suppression of urine. She was a hale old dame, and after three days or so of active treatment, the urine reappeared. Of course a careful examination of it was made, and some tube-casts were found. Taking what Sir Thomas Watson said about tube-casts as my guide, I gave it as my opinion the old crone would not long survive the attack. This coming to her ears made her very wroth, and she at once transferred her professional confidence to an irregular practitioner in the neighbourhood, —who was always in demand whenever the regular practitioner’s opinion was unfavourable, or his advice distasteful. His opinion was that she only wanted something to eat and drink, and she would do well. This advice was promptly followed, and carried out with the most satisfactory results to the

patient. She glowered at me with evil eyes, and refused to reënter her cottage, and held the poorest man of her professional capacity. And well she might, for she had passed and gone since then, and she still refused to acknowledge to the erroneousness of my opinion of her case.

A very common cause of interruption to the progress of convalescence is an excess of food. There is a disposition to rise of temperature, sometimes vomiting, commencing with the result of clearing the system. In strong organisms these intercurrent derangements produce little or no effect; but in delicate individuals they are fraught with a certain risk to the patient, and often have a very malignant effect upon the onward progress of the case. In those who have been very gravely ill, and in those who have any constitutional deficiency or special gastric idiosyncrasy, the very greatest care must be taken. Such a case came under my notice lately. A delicate girl had a slight pneumonia with a good deal of gastric irritability. By very careful dieting she went on very satisfactorily and seemed likely to weather the storm; but just as she seemed "out of the wood" an injudicious meal inaugurated a rapid downward progress.

Light food, as chicken, rabbit, game, and small birds must prepare the way for chops, steaks, and a cut from the joint, together with suitable vegetables. The dietaries of the London Fever Hospital may be adopted until the whilom sick person is out of the doctor's hands.

The amount of food a strong and previously hale young man will eat in a day, after an acute attack of fever is something surprising. We hear of eight or ten meals a day, and pretty good meals too, being taken under such circumstances. In fact, the convalescent wishes to be back to his place in the world and to his work, and he expedites his progress thereto to the best of his ability. But there is an old adage which says, "The more haste the less speed," and sometimes this is the case.

It is not always desirable that the plentiful dieting of the valescent be too rich in albuminoid material. No doubt it has been tissue-waste, often extensive, and nature has been called into painful play. But this end is not always served by over-indulgence in rich food. Some time ago a young gentleman, who had suffered severely from a typhoid fever, had a violent attack of rheumatism. His heart was considerably weakened by the fever, and the rheumatism being on in the pectoral and humeral joints, he was unable to

could to a French cook, whom he found in Bombay. Here he went in for liberal meals supplemented by generous Burgundy. But to his great disappointment he did not improve as he had anticipated. On arrival at Aldershott, he found a friend deep in the perusal of my work on "Indigestion and Biliousness," who drew his attention to some remarks therein, as to the difference betwixt merely swallowing food and its digestion, and the importance of light food where the liver cannot deal with a large quantity of proteid matter. His own plan having failed egregiously, he determined to accept my suggestions. He put himself upon farinaceous food and fish—matters his liver could deal with—forthwith, and the results were so satisfactory, that he made my acquaintance in order to report to me the (to him) wondrous results.

Where the liver is weak it is not well to overload it by an excess of albuminoids, for the burden only deepens its inability to carry on the metabolism of proteids.

CHAPTER XVII.

FOOD IN GASTRIC AFFECTIONS.

IN this chapter it seems desirable to commence with the lighter forms of trouble, and to proceed, later on, with the consideration of the graver forms. Indigestion will be dealt with in a later chapter at full length; the present consideration being devoted to positive disease of the stomach.

The main maladies of the stomach are catarrh, atrophy of the glands, gastric ulcer, and finally cancer. In the first there is a quantity ofropy mucus given off, and as the food in the stomach is rolled over and over it becomes gradually covered with a wrappings of this mucus, upon which the gastric juice is impotent to exert any digestive effect. The food with some gastric juice is taken rolling in a bag, and feels like a dead weight in the stomach. It is so obvious that in such a case all solid food must be rejected, and such food only be taken as can not be so encased by mucus. Fluid food alone must be taken, that is obvious.

Atrophy of the glands of the stomach involves functional incapacity, and only food of the most digestible nature should be taken. Food which is not acted on by the stomach so much as to be absorbed is here indicated, and that means that milk is not taken in large portion of the dietary. Farinaceous matters, such as bread, but not so predigested, or soluble carbohydrates.

In gastric ulcer there is little or no distress when the organ is at rest, but as soon as food enters a pain is set on foot. This is partly to the acid gastric juice irritating the raw surface of the ulcer, partly to the dragging upon the base of the ulcer caused by the peristaltic movements of the stomach, and to the pressure of the food. Vomiting at once gives relief from both causes, and the food which is collected, and which contains the irritating acid, is rejected, only food that can be taken without suffering.

In cancer of the stomach pain is at once a very distressing feature, but there is also pain when the stomach is empty, and the food is entirely sent at the pyloric sphincter, and is rejected, and the passage of food out of the stomach. However, it is not the pain which

foods can be taken without suffering, and towards the end all food aggravates the suffering.

To show what may be done by a suitable dietary, the following case is instructive. An old lady at Brixton suffered much from pain and sickness caused by cancer of the stomach. A lump as big as a doubled fist could be distinctly felt. Yet by putting that old lady on a strict dietary of fluid food and predigested carbo-hydrates the stomach quieted down, there was no sickness and no pain, and at the end of a week the friends asked the regular medical attendant if there was not some mistake in the diagnosis. So great was the relief afforded that the patient lived on until cancer broke out elsewhere; indeed it was some months before she succumbed, when she had only seemed about a week from the inevitable end when the change of food was begun.

The great matter about any disease in the stomach, any gastric irritability which may crop up as an intercurrent trouble in the course of other maladies—as phthisis for instance—is to introduce but a small bulk of food at once into the stomach. That is the fundamental matter to be kept ever in the centre of vision, and not only that but well focussed. It is too, the central affair round which all the rest turns. Metaphors and similes might be multiplied indefinitely, but no amount would be too great for the subject matter. “Bulk” is the first thing to attend to.

Small quantities of meat broth thickened with baked flour but so thickened that the eye can detect no mass; milk with Vichy or Vars, or other like waters, these are the staples of the dietary. To these may be added malt-extracts (marvellous food in bad gastric cases) where the patient does not dislike them. Small quantities of the syrup of stewed fruit are often grateful to the patient. But in the milder cases, some boiled white fish may be borne and forms a change. Sometimes a very strict diet, with the patient in bed to economise the powers and reduce the body-expenditure to a minimum may be enjoined for a definite time, as a curative course in ulcer or catarrh. Such a severe regimen, eked out by nutritive enemata, will do much towards a cure, even in very bad cases, and achieve one in slighter cases.

When there is a great excess of acidity, vomiting gets rid of much of the offending acid. Milk is returned in firm curd; and must be given with some fixed alkali, as prepared chalk, or light magnesians, as much as will lie on a six-pence, or sometimes even a shilling, to the half pint—as lime-water or aerated alkaline waters

are not strong enough to neutralise the action of HCl , &c., are not enough to secure the desired effect, are essentia. *Alia*. Many of the prepared carbo-hydrates, may be exhibited. Care is to be taken in these cases. When slight perturbation of the stomach is all that is to be feared, cane sugar must be abandoned, and other saccharine matters being substituted for it. Indeed, broadly speaking, all the dissoluble sugars, which readily undergo actions forming acids, must be given up, and glucoses, or anhydrous sugar, taken in their stead.

In those cases where thirst prompts the sufferer to give up a good draught of fluid, which the stomach promptly rejects as objects, it is well to give the patient small pieces of ice to suck. The melting ice cools the dry fauces; and, introduced in this gentle manner, the stomach tolerates the presence of fluid. It is possible, by the use of ices containing fruit syrup, or cream, to introduce a small quantity of nourishment into an irritable stomach. When the stomach is in active revolt and objects to all contents, this is a matter of life or death, very often, how far food can be introduced into it. Hunger and thirst both are experienced; where, in many cases the patient manifests an intense dislike to food, &c., or not. Where the latter is well borne, the difficulty is small, but where too objection to enemata is insuperable, the greatest difficulty may be encountered.

When the quantity or bulk of food which is tolerated at once is small; the meals must be the more frequent. If they are sufficient to be called meals, they should be of the full number, being, fast, eleven o'clock, luncheon, tea, dinner, and supper. Half-meals by no means too many. When only two or three ounces of the food can be taken at once, then the feeding times should number a dozen in the twenty-four hours. Where only a very small bulk is tolerated, every hour is not too often. The smaller the bulk, the shorter the interval, else the patient will starve.

Sometimes the patient loathes the only food which is suitable. He hates beef tea, cannot take milk, eggs disagree, and butter, &c., of an egg is good, where little can be borne at a time. What is to be done? Try a teaspoonful of malt-extract every hour. If this new food, not handicapped by memories of association, will pass a fair trial, and often succeeds, especially with food of the agreeable, the size of a Barcelona nut, of *Cake of Holland*, &c., &c., tried hourly.

The resources of the physician are often exhausted.

troublesome stomach cases, but success often is the reward of extensive acquaintance with food of all kinds and qualities. But then, this acquaintance is only possible by long, watchful attention to the subject.

The principles of feeding in gastric cases are the same, whatever the form of disease, though in minor points some variation may be suggested by the individual exigencies of each case. The digestive act remains the same, whatever the precise form of disease. Small quantities of readily digestible food at once, is the secret of success. Enemata of soluble carbo-hydrates and peptonised meat come in very conveniently—there is no palate to be offended in feeding *per rectum*.

Of course, there are always the individual likes and dislikes to be encountered; as also the fact of certain foods disagreeing with the patient: two matters which often tie one's hands very awkwardly in dealing with cases. And it is in women—in whom, as compared to men, assimilation is imperfect—that these disturbing factors are encountered most frequently. When the assimilative power is very small, the patient must be put to bed and the room kept warm, to reduce the body-expenditure to a minimum. The amount of food which will "sustain" life under these circumstances, is surprisingly small. And after a period of comparative rest, the appetite and power of digestion return. Such a case was put on record by Professor Seegen, where for thirteen days, but thirty-five grams (515 grains) of milk were the sole food. After that, the appetite returned, and considerable quantities of milk, an egg and some arrow-root, were taken daily. It is possible, too, to give some food to such patients by having them rubbed with oil. Absorption by the skin, even if small, is of priceless value in an emergency of this nature, when, indeed, every little helps.

Sometimes violent and irresistible cravings for food of the most apparently unsuitable character are experienced. It becomes, at times, a matter of no small difficulty to decide upon whether these cravings shall be gratified or not. Experience has shown that frequently the food so craved for has agreed, despite all *a priori* objections; and consequently furnishes an argument for yielding to the patient's desires. Granting these exceptions to the rule, the broad rule remains of small quantities of liquid food at frequent intervals in all cases of disease of the stomach. Of course, this does not thrust out of sight the fact that the stomach requires its own intervals of physiological rest; but the smaller the meal and the more digestible its nature, the less this interval is.

CHAPTER XVIII.

FOOD IN STRUMA.

THIS is a very important matter at the present time. Struma is the word now in use in lieu of scrofula. Scrofula was a name for that suppurating of the cervical glands which followed from enlargement of these glands; and got its term from Scroph, a scow, from the porcine appearance of the neck. This it was not, however, till the Prince Regent adopted those voluminous neckties which were worn in the early part of the present century. It was vulgarly known as "the King's Evil" because it was believed it was cured by the royal touch. Probably there was something in this, but it was not, not directly. The child which succeeded in getting a kiss from the royal touch would probably after that get the special treatment and regimen which is so useful in strumous, or scrofulous children.

Struma is closely linked with, if not really a dependent condition, with tissue-degradation. There is a certain enlargement of the epiphyses, often noticeable in the knotted fingers; among the bones, the bones, as a rule, being small and light. The muscles are small and apt to be flabby. The lips are thick and seem to curl about the nose. The eyebrows are often well-marked, and often straight or highly arched. The little fairy child, usually young, with brown hair, well vaulted forehead, cleanly defined eyebrows, pointed nose, and archness of expression, with long slender fingers, is a strumous child; and will usually not survive childhood. It is in contrast to the heavy plain child of coarse features, thick lips, or short or stunted eyelashes, enlarged cervical glands, and a strumous nose, not uncommonly seen; both are alike, if not equally strumous. The great lexicographer, Sam Johnson, had a strumous countenance, with the knobby head of confirmed struma. He was the last who was touched for the "King's Evil." He did not seem to have benefited much therefrom, was a devoted and promising true-blue Tory all his days. Professor Lushington, a person of the strumous clutches, thus describes the "vital force manifested by defective nutrition, and by defective and deficient function." And adds, "the child of struma is a

it is "of a retrogressive type towards the infantile or a lower ethnic form." Of the nervous system it is "irritable, feeble in tone, mind precocious, but mental powers imperfect, or if good, soon exhausted," a "lower ethnic form" of individual marked by precocity and early exhaustion. In its slighter forms struma gives us our type of beauty; but it is an unhealthy type. The precocious fairy child will never make a strong healthy woman, fit to be the mother of children.

Of what may be called the cause of the strumous taint, the deterioration of the physique, the following illustration is fraught with instruction. In my early days of practice in Westmoreland a family tainted with struma^s were among my patients. The father was a man of fair physique of the nervous diathesis; while his wife belonged to a moderately strumous family. In their early years of married life they were prosperous as working folk, and their first children manifested little of the strumous taint,—had indeed just enough of it to give piquancy to the expression. Then the father took rheumatic fever which injured the valves of the heart, and crippled him so that he became very poor. The children born during this time had more marked struma; their noses were stubby and their eyelashes broken and defective. Then the father had some money left him, and their circumstances became easier and their regimen better. The two children born after this improvement in their circumstances resembled the first-born children in the slightness of their strumous taint. The family group was a most instructive one.

One curious matter there is about strumous children, and that is the amount of uric acid in their urine. They also are liable to acid, or sour perspirations.

Struma is a departure from the normal in a downward direction, both as to development and nutrition; and, consequently, has to be met by good food, fresh air, and indeed, everything which will tend to raise the *physique*.

In the first place, it is very undesirable that such a child be suckled by the mother; especially if the struma come from her side. When a wet nurse is sought she must first be a perfectly healthy woman, and then Dr. Combe ("Physiological and Moral Management of Children") thinks "that the nurse should resemble the mother in all the characters in which the latter can be considered healthy; she should be nearly of the same height and bulk, for it has been noticed that the child of a tall, thin woman

rarely thrives upon the milk of a woman who is a "thickset." It is also of great importance that the milk be delivered as nearly as possible at the same time as the new-born child is nursed upon milk a few months after it becomes serofulous."

Now that the wet nurse is going out of fashion, it is less necessary than it was. But the milk should be of excellent quality, and if possible got from cattle in town dairies. "It is a curious fact that the milk of a tuberculous cow—an animal which is very subject to these diseases, when confined in stables, as the case frequently occurs—has been found to contain seven times as much protein as the milk of a healthy animal." (Baudeloque quoted by Ingou, "On Serofulism.") The milk should be from a country-fed cow, brought in what may be termed a normal manner, and feeding a ration of food. Not only should such children be carefully fed, but they should be brought to the country under circumstances the most favourable for encouraging development. And without this Sir J. Clark thought that our efforts to improve their health will fail. "The good cannot be backed up by every other advantage." "The child, however, should not merely live in the country but should be as near as possible in the fresh air. I have frequently had occasion to note the extraordinarily good effects, which an out-of-door life produces upon serofulous disease in the case of children who during the season of gleaming are in the fields from sunrise to sunset, far removed from serofulous glands, as well as other forms of strumous disease. They frequently undergo a notable amelioration during the summer." (Ingou.) The advantage of sending such children to the coast, where the whole day is spent in the open air when the weather permits, is obvious. "The seaside and coast-weather," is the best recommendation for such children when affected with some form of strumous in their joints.

There is in strumous children much to be learned from the notes. "The ribs have thickened and project at the ends, the extremities 'the rachitic garland' is formed, the epiphyses of the enlarged costal epiphyses."

Melancholus is also common, the child is morose, and does not reach height, being an extreme example of the hypochondriac. Deformities of the spine are frequent. The spine is angulated, the ribs are

single or double curvature, or even caries. The music-stool giving no support to the back has been held largely to blame for this. In the old starvation days for school-girls, the imperfect osseous development of struma was very frequent. In a large school visited by Sir John Forbes, not one of the forty girls was free from some crooking of the spinal column. And of these school-girls Sir J. Clark said—"While the natural form of the body is destroyed, the general health suffers In short, all the requisites for the production of struma may be found in a large proportion of boarding-schools when the system described was pursued." But any such expression of opinion about boarding-schools at the present day would be incorrect and unjust. Still the matter of feeding delicate children, and especially girls, is a very important one, in the matter of girls more than boys, because the advent of puberty and the inrush of new ideas consequent thereupon seems to affect more the palate in them than boys. Slate pencil and chalk seem the articles most craved after by the depraved appetite of school-girls; and neither have a food-value, while the chalk locks up the bowels. Consequently, great care and supervision is requisite to prevent girls eating what they should not have; as well as to see that they eat what they ought to eat.

Strumous children, from an early age exhibit a strong tendency to form an excess of uric acid.* It is seen in crystals in the chamber utensils; and gives rise to sour acid perspirations. (It is not asserted that uric acid is found in the perspirations, only that they are sour.) Probably this has something to do with the good effects of potash upon strumous children, to which Lugol bore his testimony. He observed that the administration of potash had a good effect upon the maladies when present, "without in any way lessening the tendency to a fresh tubercular deposit."

Where it seems desirable to administer potash (which is not very palatable), it may be given in the following manner, without offence to the palate. To each pound of fruit when put into the oven to stew, add as much potash-bicarbonate as will lie on a dime or a quarter bit (in English, a six-pence or a shilling.) This will neutralize the excessive acidity without the taste of potash being present. Then some sugar may be added. Such stewed fruit with cream, would form a desirable factor in each day's food.

* Professor Laycock pointed out how struma is a return to a "lower ethnic form;" in this tendency to form uric acid, we see a return to the primitive form of urinary products—the urates of reptiles and birds.

Then comes the question of fat. Fat is necessary for the formation of healthy tissue; and when there is a rapid rate of tissue-degradation fat becomes even more necessary. But then the strumous child turns away from the butter, and revolts from a piece of sweet animal fat; it turns away from bread and butter when a stout slice of beef is offered, and from a layer of butter. How then is the need to be satisfied? By means of cod liver oil, given in some palatable form? Certainly; but is cod liver oil the best form of food in the form of fat? That may be questioned. Prof. G. A. Hays Bennett introduced the use of cod liver oil, but Dr. J. C. Wood, in his experience of it, he says, "Its use soon became general in all scrofulous and tuberculous affections; and what is more, it has stood the grim test of time, and holds its ground as firm as ever."

The fishy oil is relished by many children, who, on other grounds it might be thought the child would dislike. On the contrary, it is taken by many delicate children, and even by the case of white barbarians in Eskimo lands. The fat seems to be in a distinctive way. Dr. King Chambers tells the case of a plump cod-Skye toddler who whined for his cod liver oil for months. Other children take greedily the oil from canned sardines, a form of oil very good for them.

Then there are other ways of introducing fat without offending the eye or the palate. Of these, milk stands first and foremost; either plain, or as milk puddings, or as an adjunct to porridge. Then cream may be eaten with stewed fruit. Cream too may be served up at table as cream moulds variously flavoured; containing cream with some farinaceous matter; or as meringue. As whey and cream it is a pleasant accompaniment to many cold puddings. Then there is butter, which can be added to bread, and some potato, or in thin sandwiches, where something tasty is polished; or it can be added freely to mashed potatoes, or eaten with milk puddings, hot or cold. It is to be found in omelette. When of some importance of moment then stout puddings suggest themselves, thick cracked and stewed with cream and sweetened with sugar, or, when some, the marrow fat being never detected, as in the case of the baked loaves of New England farmers, in the neighbourhood of a peasant farm, not likely to offend the palate. This is an economical food.

Then there remains one more food, now seldom used, but of some point of view, not to be forgotten. This is the fish-bone

and sugar is acceptable to the palate of most youngsters. The old original toffee, made when butter was cheap and sugar dear, is the form to be adopted when it is desirable to afford fat to the system. The increasing price of butter has reduced toffee (in most instances) to the level of a mere carbo-hydrate; well enough as a fuel food, but inoperative for tissue construction. In my experience many a little strumous child whose palate revolted at cod liver oil has been saved from death by liberal supplies of toffee;—though its mention usually provokes a smile of derision and a sceptical remark from the mother. She soon however sees for herself the beneficial effect it produces. Incredulity gives way to gratitude: just as of old those who came to scoff remained to pray. There is other conversion than religious conversion!

In struma advantage can be taken of the ready absorption of fat by the skin. Children readily admit of cutaneous injections being applied to them. In a warm room the child's trunk and limbs can easily be sponged over with warm water and soap so as to cleanse the surface, after which oil, which has previously been warmed, can be applied. It should be rubbed over the skin freely. This can be done one or twice daily. Cod liver oil probably is most readily absorbed, but its smell is objectionable. Then comes neat's foot oil, also an animal oil; olive oil is less offensive than either. Such injection in cold weather forms a good protection against cold.

After a suitable regimen under favourable circumstances, as an outdoor life at the seaside, the system will often greatly outgrow its strumous tendencies, and a fairly healthy organism be attained; where if no steps are taken the system simply gravitates deeper and deeper into confirmed struma; and sinks ultimately under some tuberculous or scrofulous disease. The preventive treatment of strumous affections is a great matter. When some joint has been excised or mischief shows itself in the mesenteric glands, or the lungs, the above scheme and regimen have to be adopted as a curative measure.

The great food for the strumous is "Fat."

CHAPTER XIX.

FOOD IN ANEMIA.

There is a tendency in youth and especially in girls to be anemic, or bloodless. Why this should be so is very clearly shown. It seems probable that the demands, of growth, of intellect, of work, school or other, made upon the organism, lower the capacity for sanguification. No doubt in some cases inherited soil is a factor. Then exposure to malarial influences, a perfect arrest of formation of blood-corpuscles, as well as, probably, causing the rapid disintegration, or wearing out of the red blood cells. A blood-poison carries with it a tendency to anemia, and often to the "pallid" form. It is the prayer of a nerve for health, "blood, blood!" Rheumatism and neuralgia, especially neuralgic pain, is never the result of a pure, or deficient in quantity. Consequently, we find anemia under the most opposite circumstances, viz., blood-poison and on the one hand, and poverty of blood on the other. Anemia, therefore, may be based upon defective blood-formation or corpuscles, when there is a blood-poison operating injuriously on blood-formation, poisoning even also destroying the red blood corpuscles.

It may be said for the instruction of non-medical persons that the blood-corpuscles do not feed the tissue, they are to be fed by the serum, or *liquor sanguinis* of the blood. Still, when the quantity of blood is good the general nutrition is not impaired.

It has been pointed out in a previous chapter that the general nutrition of the body is but mental son of nutrition of the blood, the food, or out in two matters, 1. the quantity of the blood, and 2. the quality of the blood corpuscles. Hence, the general nutrition of the body, which must be maintained, is not more, the administration of energy to the blood, but the essential part of the treatment must be to keep up the quantity present in the blood, as supplied by the formation of new corpuscles, and to cause the blood to give the nutriment to the tissues in the proper order as nature to which, however, the physician is not to be true. The special poison must be met by the removal of it from the organs involved by sanguification, and the removal of it from

readily seen by the inutility of steel and good food in a case of anæmia in which syphilis is a factor, until the proper treatment of the taint is superadded; and then the difference is noticeable at once. The treatment of anæmia often fails from ignorance or forgetfulness of this fact. It is not uncommon in cases of gout and biliary disturbance to find that weak anæmic persons have been taking steel and quinine most carefully for some considerable time without any good result; and yet, as soon as the gout poison is met by potash, or the liver put right by hepatic stimulants and cholagogues, improvement at once sets in. When a certain point has been reached, then the iron comes in most usefully.*

These matters, lying somewhat outside pure dietetics, must be pointed out in order to prevent disappointment. Valuable as a knowledge of dietetics undoubtedly is, it must be remembered that feeding cannot do without medicinal aid in many cases. A man who has lost both legs would do little with one crutch.

Purity of blood and iron to enable hæmaglobin to be made are essential to the treatment of anæmia: but proper food also is requisite. The food should consist to a considerable extent of animal food (except in gout and cholemia) at least when the case has made some progress. At first in anæmic cases in girls it is well to send the patient to bed. The assimilative powers are feeble and cannot digest enough to feed the system, and meet demands on the body as well at the same time. To send the patient to bed to lessen the body-expenditure is the first step. By so doing even a very feeble digestion can supply the wants of the organism, with a little to spare for improvement. "You cannot eat your cake and have your cake" says the old adage. No more can a starving organism use up each day's income and expect to store and garner strength. A few days in bed is the proper start in marked anæmia, especially with girls. This is a lesson experience has taught me with some humbling of self-pride. Pain is of various kinds; and wounding of the self-pride may make as enduring an impression as the infliction of physical pain—as when they flogged certain healthy boys in order to imprint the locality on their memories in the old rude method of "riding the boundaries." Such a case occurred to me when attached to the West London Hospital. A pallid nurse-girl was an out-patient

* Why iron is less useful in anæmia linked with the formation of blood poisons within the body than in other forms, it is impossible to say, but the fact remains.

be of a character to keep the bowels open. Oatmeal porridge, dishes prepared from maize, vegetables, fruit, fresh and stewed, and meat, and meat broths, or soups. These should all be given according to appetite and palate and body-wants.

There is a difference betwixt anaemia and convalescence from acute disease. The latter is a case of quickly down quickly up again very often. In anaemia there is more chronicity, a slower failure of the powers and a slower recovery. A person regaining flesh and blood, and with them, strength after anaemia, has not the recuperative powers of a previously healthy and robust adult laid prostrate by the stroke of acute disease. The appetite is not so keen, the digestive powers are not so vigorous. The regimen appropriate for the one will not fit the other. Circumstances alter cases. Certainly! And light food of readily digestible character should in each case precede the solid meals which are appropriate later on, while it is almost needless to say all body-outgoings should be reduced to a minimum.

were readily taken, and that the animals so fed, did better than those who got no shavings. A certain amount of woody fibre in their food seemed essential to their health. ("Fruits and Farinacea.")

Conflicting views are held about the dietary of constipation. Some writers advise avoiding food which will load the bowels as vegetable parenchyma, while others again discountenance such food. That the bran of wheat is good there is no doubt. But if the hypothesis put forward to account for this, *i.e.*, that the mechanical irritation of the bran-scales excites the vermicular action of the bowels, is true and well founded, then it would seem the bran, or much of it is not digested. Certain it is that the coarser the flour and the more bran it contains the better it is, as has been the experience of sundry mariners. And the custom in Westphalia of eating bread made from unbolted flour has been introduced into England (Liebig) with advantage to the bread-eater. The addition of treacle (which is largely levulose, or uncrystallisable sugar) to brown bread is credited with increasing its aperient effect; and "brown bread and treacle" is in vogue amongst the students of the impecunious class whose sedentary habits clog the acting of their bowels.

It may be said as a broad rule, that flesh meat, and meat infusions have a distinct tendency to keep the bowels open; while milk and farinaceous matters tend to constipation. Cheese has a reputation of its own in this respect; while eggs, especially if hard boiled, also bind the bowels.

As to the peculiarities of individuals they cannot be discussed here. One person is potently affected by a small quantity of food which in the bulk of human beings has a purely negative action. For such persons abstinence from the articles which disagree with them is all that can be advised.

Thus, some persons are readily acted upon by fruit, especially fresh fruit, when ever they indulge in it; while nearly all are affected by rhubarb when first eaten in spring.

Under certain circumstances foods which ordinarily tend to lock up the bowels, set up active purgation. Thus milk, ordinarily binding, when the stomach is acid so as to set up excessively firm curdling, starts a sharp diarrhoea; a natural eliminant process to get rid of the peccant matter. In the same way, other like matters act. Or so, at the Leeds Dispensary, a woman brought a child ill with diarrhoea. On being asked what she had given, she replied "Hard boiled egg," and seemed to feel herself decidedly ill-treated by

without calling in the aid of strictly medicinal agents. But frequently these last are indispensable.

In more chronic conditions of looseness of the bowels, milk and farinaceous foods are still to be made the staple of the dietary. Stiff puddings, as suet dumpling and plum puddings, are held to be constipating, and the same is said of pastry; but if such is really the case, certainly neither are suitable for conditions of intestinal irritability, whether catarrhal or otherwise. In chronic conditions of diarrhoeal character, some objection lies against animal broths and meat infusions; and it is well to give them, or soups with sago, biscuit-powder or vermicelli.

Creeled rice is a very good food in such conditions; and the juice of cranberries is useful, as there is some astringent (tannic) in cranberries. The juice of the pomegranate is also pleasant and good in any looseness of the bowels.

Then come the astringent wines, rich in tannin—as Claret, Carlowitz and Catawba. These may be drunk undiluted; or may be made into a nutrient food by adding them to solutions of grape or cane sugar, or even to lactated foods. In many cases the small amount of alcohol is desirable.

It may be laid down as a rule, that farinaceous matter is useful in diarrhoea: the soft starch being not only non-irritant, but actually soothing to the morbid mucous membrane of the bowel.

No doubt in many cases of diarrhoea, acute or chronic—just as in constipation—the resort to medicinal agents is often necessary. But, granting this and admitting the numerous remedies in our possession for the relief of both conditions, still, their action can be potently aided or thwarted by a suitable dietetic regimen. Indeed, in the milder cases, regulation of the dietary is sufficient to keep the bowels in a satisfactory condition.

In the diarrhoea which belongs to travelling, it is well to take a full dose of rhubarb—which first opens the bowels, and then locks them up as a secondary action—the day before travelling; and after this to take milk and farinaceous matters only till the journey is over.

(In consequence of this secondary binding action of rhubarb, an artificial constipation is far from uncommon. By substituting some other aperient for the rhubarb for a time, the bowels commonly come to move naturally and of themselves. Rhubarb is indeed of all aperients that least fitted for habitual use.)

CHAPTER XXXI

FOOD IN PHTHUSIS

resistive power to the tubercle bacilli inspired by all;" or to state it thus—"Certain families manifest a tendency to lung-changes in which the tubercle bacilli find a favourable nidus;" we must confront the fact that there is a something in some families which renders its members liable to become phthisical, which is not present in other families. Taking a broad view of phthisis, it can be honestly affirmed that the families which exhibit this tendency to tuberculosis are those in whom the strumous diathesis is existent. It is the "lowly ethnic form," which carries with it the tendency to tissue-degradation, which gives us our phthisical families.

Laennec laid it down that tubercle was a neoplasm—a new growth of tissue—unknown in the healthy organism; and when it was found the case was hopeless. To such dismal view has succeeded another and truer one, viz. that there is nothing in tubercle—no histological element—not found under other circumstances. Tubercle is a rapid proliferation of connective-tissue corpuscles—the lowly basement-membrane, or packing material of the body—the more rapid and excessive the production, the more huddled together the cells, the more liable is the mass to die. The sparser the production and the nearer the new growth to normal tissue the better its chance of survival.

This pathological disquisition is not the pursuit of a side-issue of the question, but an essential matter which the reader must firmly grasp if he, or she, is to have a correct and vigorous hold of the dietetic management of phthisis. When the cell growth is rapid and extensive, the crowding of the individual cells upon each other means death to them. It is like crowding a host of individuals upon a limited area so closely that it is impossible to feed them; those in the centre die first from starvation. Rokitsanski speaks of this "central softening" and of "the ocular proof that the softening commences at the point most remote from surrounding textures" in tubercle. Such softening or caseation of the mass is accompanied by hectic fever which may, or may not exhaust the powers; and wear the patient out in a shorter, or longer time according to circumstances. If the softened mass can be expectorated a cavity remains, which may include the whole morbid area, leaving the individual short of so much lung, but otherwise little the worse. At other times the expectoration of the softened tubercle is accompanied, or followed by a new development of tubercular growth doomed to soften and die; and the long continuance of the hectic fever exhausts the powers.

But the course run is not always so good. "The growth may not be so free and the cells may not be so another so fiercely, and so some nutrition of the cells may be possible; in which case death does not occur." A lung, in truth, belongs as little to the nature of the tubercle as itself" (Rokitski). But if enough of nutrition can be secured, the mass of lowly texture it may make an ideal material for some cases only a mass of lime salts taken a piece of meat, or even to tell of the by-past tubercle; which may stay in a lung expectorated long after all evidences of lung mischief have disappeared. Such is the history of tubercle. But what does the clinical pathologist tell us? "The cure of tubercle may take place in various ways. Each of the metamorphoses of tubercle may become associated with the character of a healing process. Still neither the absorption of tubercle, nor its ejection through the medium of ulceration, nor local healing processes, are fraught with a value to the system, unless accompanied by the extinction of the tuberculous process, or producing crisis." This is the great matter to attend to, says Niemeyer. Opinions differ as to when the term "tubercular" can be applied to the mischief in the lung. Some hold it may be used when consolidation of the lung is on foot and in certain cases this is so, when lung consolidation is found in a person of tubercular taint, who has been wasting for some time and had night sweats. Others hold that a mere consolidation of the lungs is not enough, and that it is only when there is evidence of softening or about there that the term "tuberculous" can be truly applied. In the language of Niemeyer "we have no hesitation in stating that the greatest danger for the majority of consumptives is *that the cure ought to be made tuberculous*."

He says too-- "Prophylaxis against consumption can be made in the first place, that when an individual shows signs of defective nutrition and a delicate constitution, especially if he have given evidence of unusual delicacy with a tendency to diarrhoea, or to morbid excretions, *he should be put on a course of food which is calculated to invigorate the constitution, and to counteract the morbid tendency*."

The prophylactic treatment of tubercular affections, therefore, is to prevent, or remove that disintegration of the body and its tissues which gives the tubercle a field for degraded connective tissue composed of cells that are not organised to live. There is a general tendency to the formation of

the inherited constitution is a great matter. Evil surroundings and the mode of life tell on the parent-organisms, and what is acquired by the parent becomes inherited by the child. Lagol said the population of Paris of his day was eaten up by scrofulosis, yet their ancestors three or four generations back were healthy people from the country. Dr. Walshe, and still more recently James Cantlie have pointed out the degeneracy of the populations of large towns. Vogel says "Children bring no existing tubercles into the world with them," though "there is no disease so positively inheritable as tuberculosis." Granting this he goes on to say—"The following are generally considered the external causations of tuberculosis: bad air, confinement in close, imperfectly ventilated, dusty rooms, damp houses and bad food—by which living exclusively on rye bread and potatoes, and the deprivation of animal food, is understood."

With the other matters than bad food we cannot deal here: but with it we can deal to some purpose. If the tendency be a matter of inheritance let the physical links betwixt parent and offspring be severed as soon as may be. The mother should give place to a healthy wet nurse (chosen on lines and principles laid down before). In households where this cannot be accomplished artificial feeding must be wholly or partly carried out. In the matter of struma insistence was laid upon the necessity for fat for the formation of sound healthy tissue. Phthisis is an outcome of the strumous habit; and to improve the strumous habit is to reduce the risk of phthisis.

Fat is requisite for the building of healthy tissue-elements. Milk gives that form of fat which is most desirable, viz.: the emulsion. Fat in milk exists in a very fine emulsion, the fat-globule being very minute. Milk is most acceptable to all young creatures, and is usually taken by them freely. The dietary should be so arranged as to include a quantity of fat-bearing milk. To this may be added some predigested starch if needful. But as soon as the infant can chew, and has some organs of mastication it becomes a question how far it is not good policy to give it some food on which it can exercise its jaws. A biscuit, or cracker, or piece of bread munched would tend to thirst calling for a draught of milk. Such starchy matter well chewed would develop the jaws and their accessories. Too much soft food, whether of insoluble, or of predigested starch, leaves out of sight the natural means of developing the salivary glands. The gnawing which a child delights in is started by the irritation set up by the teeth coming through the gum. During the whole time of the growth of the primary or milk teeth this irritation

wonders for that little starveling. It is not uncommon in the North of England to see a child fed on pap for its meals with a little bag to suck or chew betwixt meals, containing butter and sugar, called a "sugar-pap." This is a capital plan when milk cannot be obtained, and supplies an excellent form of fat. The utility of ordinary toffee, as an admixture of carbo-hydrates with hydro-carbons possessing a high nutritive value, has been pointed out before.

Such then are the various modes of introducing fat into the youthful organism. There remains the injunction of fat to fall back upon.

Then most authorities advocate a supply of good vegetables for children—not in lieu of, but,—in addition to other food. These supply phosphate of soda and other useful salts. The same end might be met by animal food, if it were not for the fact that in children of strumous or tubercular constitution, they are given to the formation of uric acid in larger quantities. Potash, a uric acid solvent, is advocated by all authorities on tuberculosis or serofulosis.* Phosphates must be supplied freely to children of this diathesis, as they readily escape by the urine; and, of course, if voided in the urine, are not available for the tissues where they are so needed. Well-cooked vegetables should be allowed—and form part of the dietary. For the supply of other salts, as the citrate-malate, or tartrate of potash, for instance, fruit in various forms should not be forbidden; indeed, the eating of it should be distinctly encouraged.

A healthy child can extract what its system needs from very unpromising materials; but not the child which comes into the world with an impaired physique. It may not bring with it into the world ready-formed tubercles; but it brings with it the tendency to their formation. And with a thorough knowledge of this fact, we must do our best to combat the inborn tendency.

The milk of the Alderney cow is best. Then that of the black-poll'd Angus breed; but the milk of any healthy, properly fed cow

* There is a curious and interesting point about the formation of uric acid by children of strumous or tubercular family history. Professor Laycock pointed out that the skeleton of the strumous is of lowly ethnic form. I have elsewhere pointed out how the formation of uric acid by a mammalian liver is a case of reversion, or turning back of the liver to the form of nitrogenised waste found in the birds and reptiles who have a solid urine. The mammalia have a fluid urine, largely composed of the soluble urea. We see then, in the strumous, osseous degradation as well as hepatic degeneration; both of which must be borne in mind in their dietetic regimen.

will do. Mixed with the various farinaceous articles, a milk pudding, milk should be a staple of the dietary. Butter should be freely supplied. Bread and butter pudding is good. Butter should be added to milk puddings at the table with no objection. Fat should be encouraged; and suet puddings are good. The salted cod and pork of New England form a capital food, rich in fat. Puddings should be well buttered; and for a change, some milk warmed with a knob of butter melted in it, could be served at table with the boiled potatoes. Pastry made of French puff paste may be a luxury (when it does not cause indigestion), in reasonable quantities. Where the appetite is capricious, even chocolate creams have a value. Sweets may be essential; but so far as possible the eating of sweets betwixt meals should be avoided. It may be necessary to give out small meals by sweets; but a watchful and even suspicious eye should be kept on the practice in order to avoid the error of stimulating the appetite for meals by sweets. Indeed it is only in rare cases that the plan of "any food, rather than none," is necessarily adopted; it only should be adopted when no better alternative can be devised. Of course toffee (containing butter) is the best form of sweet; and I know at least one comely damsel who owes her life to toffee in the early years of puberty.

In some of the Northern parts of the U.S.A. a cure for phthisis in vogue is to go lumbering where plenty of exercise in the open air, the sharp chill air of winter, gives an appetite for fat pork. Such a plan is only suited for phthisical patients with large and energetic frames. The principle is sound, and the practice may be modified in a less severe and gentler form, by many young persons threatened by phthisis. Plenty of exercise in the open air, with bonny and pork for victuals, would save many a life; especially amongst that wretched town-population doomed to live in small, crowded apartments, where the air is foul and laden with miasma and germs; who form a large proportion of the phthisical patients who crowd the waiting rooms of all hospitals for diseases of the chest.

And now a few words about the use of cod liver oil. As pointed out before, fish oil is not the most desirable, or the best tolerated, but it has the advantage of being the most digestible. It should be taken after food; and in order not to needlessly overburden the system, the interval should be about an hour, i.e. as long as possible, after the food passing out of the stomach, and the process of the digestion of fat begins. As pointed out before, the digestion of fat does not commence until the food has been in the stomach

quently if given before, the oil only swims about the stomach, offending it, and causing spasmodic contractions, or excitations. Whenever the stomach resents its presence, it is well to administer the oil at a distinct interval after meals. In other cases it is convenient to take it immediately after food (as in the case of working-people going to their work), and if it does not disagree it may be taken then.

After ten years' experience at the City of London Hospital for Diseases of the Chest, and my other experience at the West London Hospital, and at the Leeds Public Dispensary, to say nothing of over twenty years of private practice, in the thirty years which have elapsed since I was bound apprentice (as an apothecary) to my father, who was in general practice in Westmoreland, I may say that in encountering phthisis my greatest dread is a distaste for fat. Where the nutrition is good, the prognosis is, as a rule, favourable; even if a large area of lung is implicated. While on the other hand, in those cases where there is slowly progressing emaciation with a distaste for fat, the prognosis is as dark as Erebus, even before any lung complications set in; which commonly enough is not till an advanced stage is reached. Whatever may have been the teaching of the experience of others, that is the lesson my experience has taught me. When a patient is steadily wasting with a tainted family history, and a personal history of a loathing of fat intensified of late, then the forecast of the future of that patient presents no difficulty; it is, indeed, only too easy to tell what it will be. The practical outcome of this is the insistence of fat—in some form—being taken without waiting for lung-mischief to show itself. That lock which opportunity carries in front of it (being bald behind) must be energetically grasped; and to be forewarned is to be forearmed. If the patient can be induced to take fat for the tissues, all may yet be well. The absence of fat (somehow or other) leads to a dyscrasis which favours the production of tubercular growths. Even when tubercle has developed and softened, cod-liver oil has snatched many a patient from the grave—yawning before him, or her. If fat can be taken in sufficient quantities, healthy tissue can be built up around the diseased area, and the ravages of the invader arrested. Where the absence of fat leads to a depraved tissue-growth and repair for a considerable time, then a condition of much danger exists. Prof. Laycock called this—"the pretubercular stage of phthisis." Any tubercular growth then formed will certainly start on a downward progress in the least possible time. The impaired tissues

In many cases it might well and fittingly be preceded by some milk with a little sugar and a small quantity of rum, or brandy (and a suspicion of nutmeg) in it, about 7 A.M. Once this "rum and milk" was a far-famed and sovereign remedy for consumption. And there is no doubt that with many persons such "break-fast" is good, as breaking the long fast of the night, after which the ordinary breakfast can be enjoyed; where without such preparation the stomach would feel itself unequal to a hearty meal. Put in scientific phraseology this furnishes to the stomach the matters requisite to the formation of the gastric juice—known as "Peptogens."

At 11 A.M. might come one of the little interludes just spoken of.

For lunch a cupful of white soup, or vermicelli soup, with a digestive biscuit and a small sweetbread, or a small fish as a sardine, or anchovy on toast. A glass of good red wine would not be out of place. Then a rest on the couch would not come amiss. At 4 P.M., a cupful of chicken broth with plain biscuit broken into it, or some milk with Mellin's food might be taken. Dinner at 6 P.M. may consist of a whiting or some flat fish, followed by a chicken or some game, a little bread and butter pudding and, if fancied, a little cream cheese or a nip of other cheese. A glass of good port, Bueellas, or Madeira, or Marsala (according to the purse) may be taken. At bedtime some treacle posset (milk made hot with treacle and some ginger) may be taken on getting into bed; and the rest be put in a jug with a lid (under a big tea-cosy) for the small hours of the morning. Such would form a liberal dietary during the hectic of softening tubercle; and as soon as the fever abates, some cod-liver oil may be added.

The food when the patient is confined to bed should never be allowed to remain in the room one moment more than can be avoided. Perhaps grapes are the only exception to this. Being covered with a stout impervious skin, the sweet flesh is kept from taint. Only just as much food should be taken into the room at once as the patient is likely to consume, and no more! The contrast to this is a weak, wasted patient with a wan, worn face, and a hand almost transparent, languidly looking at a table on which stands a cup half-full of milk; a saucer with some canned apricots in it; half an apple with a cut surface going brown; a glass with some half melted ice floating in water with a metal spoon in it helping to melt the ice; half a glass of port, which has been there an hour or more;—and trying to fix the resolution as to which of the group he will try. No wonder the appetite is languid, and that choice is diffi-

cult, and that commonly enough the attempt is
frigid.

The physician may be doing all that he
can, may arrest the night sweats, control the fever,
regularize the bowels, and the stomach in good condition, but he must have
a nurse naturally gifted and carefully trained, and he must be
feeling as well as nursing; the results are unsatisfactory. The
physician must exercise a general supervision, and
praise for the good nurse, and a word of admonition to the
capable or careless nurse; and for this he too must be carefully
trained.

When those times of gastric irritability come, as often they do,
and do it, most cases of chronic phthisis suffer, the feeding becomes
a cardinal matter. Small quantities of good food at frequent
intervals are essential. After some studies of these matters, and
of gastric irritability, the conclusion arrived at is, first, the more
extreme and rigid the dietary the more quickly the stomach
gets over it. On a very strict dietary this may be done in four or five
to ten to seventy hours; whereas on a moderate one it may
only it may be a week or ten days before the gastric disturbance
is overcome.

A small nutritive enemata of treacle and warm water (one
half) or a syrup of milk-sugar or grape-sugar may be given
daily, to help to maintain the general nutrition. Or the enemata
be rubbed with oil, as a makeshift.

With modern knowledge of digestion and body-chemistry,
with a more perfect grip of the essential wants of the patient, and
the importance of arresting all outgoings, as well as that of the
production of food at once digestible and nutritive, the
of the phthisical are much brighter than they were at the
of the present century. A suspicion that there is an
twain merely getting food down into the stomach, and that
is absurd; and that a tablespoonful of milk, or a
is digested, is really better for the patient than a
"milk-purposes through the alimentary canal." The
much-trusted organ, in that which it really does
efficient help to it. For per se it can do nothing, and
there is gastric irritability on foot, it is a good idea to
food." Bullock's food and raw milk, and
offend to the consumptive, and when the
to swallow, have no advantage over the
much has been said by this means.

All along the line, whether during the periods of comparative health, or times of acute peril, with patients with a phthisical tendency, the great cardinal matter is the nutrition of the organism. To build up healthy tissue is our aim; and for that, fat is essential. And usually, fat is loathed and avoided. Imperfect nutrition is what we have to contend with, and the side from which danger is ever threatening, more or less seriously. Cod-liver oil, oil emulsions, preparations like *Cremor Hordeatus*, and *Oleobyne*, all strengthen our hands, as regards the permanent treatment of the phthisical. In those periods of gastric disturbance to which such patients are liable, prepared foods are invaluable. "The subjects of pulmonary phthisis resemble in one respect, hand-fed infants. Like them, they are liable to repeated attacks of gastro-intestinal catarrh, which gives rise to indigestion and flatulence. These attacks, by the influence they exercise upon general nutrition, may produce very serious consequences." So says Dr. Eustace Smith, in the article on Chronic Pulmonary Phthisis, in his classical *Treatise on the Diseases of Children*; and his experience at the Chest Hospital, Victoria Park, agrees with my own, that intercurrent attacks of disturbance of the alimentary canal, and with them, failure of the nutrition, are among the most serious complications we have to encounter in the treatment of Consumption. And this is even more true of children and very young adults than of older phthisical patients; but even with senile phthisis, the difficulty of maintaining the nutrition is the great stumbling-block in their treatment.

CHAPTER XXII.

FOOD IN CHRONIC HEART AND LUNG DISEASE.

SOME critical reader may be surprised, or be inclined to scoff at the distinction made herein. Is not Pith's a lung disease? he may ask. Certainly pulmonary complications are very common with pith'sis, and frequently form the main trouble and the cause of death; but all the same, but admitting this, pulmonary pith'sis has no right to be classed as lung-disease. In the proper sense of the *cast*. That would be misleading. Pulmonary Pith'sis is not a primary lung disease, as chronic Bronchitis and Emphysema certainly are. Pleuritic disease may certainly exist in a person with pith'sis or tendencies, and then a condition resembling *pith'sis* *bronchialis*, with the same indications for treatment, may be set up. But what is meant by the above heading are cardiac affections and related state of the lung, which interfere with the flow of blood through the pulmonary circulation, and the obstruction, working backwards, produces congestion of the portal circulation, and the viscera connected therewith.

Acute pleurisy or pneumonia requires the dietetic management laid down in the Chapter on "Food in Acute Disease;" while condition of pleuritic effusion, or empyema, would rather be better treated by the lines laid down in the present chapter.

What is dealt with here is a condition which may more properly be said to be the opposite of (or antagonistic to) pith'sis. Both cardiac valvular disease, and chronic bronchitis and emphysema are connected with tubercle. Tubercular disease of the lung may, and does, co-exist with valvular disease of the heart, or with emphysema, or with all three; so very many that the very fact that a patient has tubercle of the lung. But Rohdendorf, and the late Dr. Penock, both have shown that the disease of the lung, which relies upon in some sort of tubercle, is not really connected with tubercle. While the disease of the heart is certainly, if you can, a case of emphysema or chronic bronchitis, or a combination of pulmonary pith'sis, or a variety of conditions, it is not tubercle.

The point to be well borne in mind is, that the disease of the heart and disease of the lung, are not the same thing.

offered to the flow of blood through the pulmonic circulation, and the effect of this upon the blood-current and blood-vessels behind the obstruction. A certain damming of the current is the result, and with that, congestion of the liver (impeding and impairing its functional activity), and behind that again congestion of the portal vein and its venules. The effect of this venous fulness is felt in the stomach and intestines; and feeble digestion and sluggish bowels are the consequence. The condition of the mucous membrane is favourable to the production of gas, giving flatulence. Flatulence is one of the greatest troubles of persons with a crippled heart or chronic lung-disease. The pressure of the elastic gas prevents the descent of the diaphragm and so further embarrasses the breathing. So pronounced may the difficulty of breathing, so set up, become, that the term "asthma" is applied to it by the sufferer. The disengagement of gas in valvular disease of the heart is sometimes so great that it pours up the œsophagus producing loud noises, and continuing for some time. The term "heart-wind" is applied to this explosion of noisy eructations.

Another consequence of this venous fulness is that the patient feels "full" even when the stomach does not contain food. It is necessary to go over this pathological ground as a preliminary to the discussion of the food appropriate thereto for the instruction of the non-medical reader, in order that he may the better grasp what is here said.

From the venous congestion of the whole abdominal viscera impairment of function follows. The stomach is enfeebled, and primary digestion is lowered, while the working of the liver is embarrassed, and much of the proteid matter which reaches it is never carried forward as the albumen of the *liquor sanguinis* for the nutrition of the tissues, but proceeds on a downward career toward uric acid and urea. The condition of the urine, more or less laden with lithates, tells of the faulty assimilation. It is impossible to feed these persons up by giving them large quantities of animal food; this simply adds to the liver embarrassment. The amiable intention merely links a condition of lithiasis to the pre-existing trouble. It is well to bear this in mind in feeding persons with chronic disease of the lungs and heart. The more severely ill the less the assimilative capacity, and the soluble carbo-hydrates, all but independent of the digestive act, form almost the only food that can be absorbed.

When fairly well persons, with old-standing valvular disease of

the heart, or with chronic bronchitis and emphysema, can share the family meal, provided that certain foods disagree with them, and that, for example, cats, some even pigeons, being sensible and rational, are not beyond this it is not necessary for them to go. There are many seasoned dishes, as the carnivorous people are good for the patience. Mustard too they patronise, and they are good for of ginger with their noses. Chutney they find suits them, and they always welcome mulligatawny soup, and still with an air of satisfaction when a curry appears on the scene. Fish *en papillote* suits them, especially when well seasoned. Cauliflower and vegetable marrow *en papillote* can be treated in the same manner and be on intimate terms with the pepper custard. All soups and made dishes to which seasoning can be added are in vogue with them, and all kinds and varieties can be made for them. Potatoes can be served up with a light boiled egg, and peppered to taste. Fresh meat, mutton, and seasoned with a wash of mashed potatoes, would form a fitting dish. Mixed collops would not come amiss, with a Scotchman would doubtless advocate a haggis, or a sheep's head, or

which can hardly be over-peppered. The toes of a swine could be prepared to meet their requirements. An Irish stew, or pepper-pot, or hotch-potch, would not be rejected by such persons, if well-seasoned. Probably they will prefer pepper and butter with a green apple rather than sugar. Pig's feet or tripe could be prepared to their mind. A fat little pig's cheek cut in pieces and served up hot with a sauce pagament would form an excellent dish, even if it were there is any gouty complaint on a very common occasion. They would probably be more likely to pay their attention to ginger-sticks, confection cookies, than to cranberry tarts, or huckleberry pie, or jumbles. Muffins and "mother's crullers" they would be likely to pass unnoticed, but their eyes might move to a ginger-bread cake. Chocolate cake and marie cake would not go to their minds at all. With milk puddings they would probably prefer rice pudding, or ginger or nutmeg, and eat it over a bowl of tea, or a bowl of eggnog, or a bowl of punch. Chocolate is good for them, but they will not eat it, or for that drink it, or swallow it. Sugar is not good for them, and pepper and nutmeg are good for them, but that is not persons. Must, therefore, be prepared to feed them with nearly anything, and to give them

tional pepper. Potted fish should be specially prepared, for added pepper cannot always take the place of seasoning put in at the proper time, as in meat puddings and pies for instance. Their whitebait must be devilled, and their prawns be served as a curry. Not that their stomachs are to be made a perpetual spice-mart, but their wants as well as their palates must be consulted.

The three things to be aimed at are, readiness of digestion, a proper proportion of carminatives, and a suitable bulk,—the latter being a matter entirely under the control of the individual. Small meals at no long interval should be the practice; and a light supper is essential to a good night. A substantial supper is very liable to be followed by an attack of difficulty of breathing in the night, and is a matter of certainty with many persons suffering from disease of the heart.

As to the amount of alcohol such persons should take, the answer is a difficult one. King Chambers says of alcohol in disease of the heart—"Let the alcoholic drink be limited to that quantity which increases the appetite. In a great many instances this quantity may be very shortly written down—0." No doubt this is quite true of some persons, especially those persons to whom alcohol is a poison, no matter what is the matter with them, whether it is their heart or their liver. Some generous wine will very commonly be found to agree well with persons with old-standing heart or lung mischief, and will help them at meals. In attacks of dyspnoea some alcohol as a stimulant is of incalculable value. As cordials alcoholic compounds are excellent, *Alcohol de Menthe* has done me excellent service.

When the condition is marked and the patient gravely ill, then a dietary of liquid food is necessitated and meat infusions, predigested starch, and milk, become imperatively called for. King Chambers also points out, that with some persons who have dilated hearts a "sloppy diet" does not always agree. Certainly there are persons who can not get on with liquid food, but they are the exception and not the rule. The observation is the outcome of the writer's extensive knowledge on the subject of dietetics, which makes him more observant of the exception than the rule.

When a very advanced condition is reached then the admixture of alcohol with the food becomes imperative. Milk with some brandy; a little beef-tea, or "invalid's soup," also with some brandy, not forgetting to add some soluble carbo-hydrates. One caution again must be observed, and that is not to push the alcohol when

symptoms of acute gastric disturbance appear. Sometimes it is a very difficult choice to make in deciding what is best for the patient, whether to risk the withdrawal of alcohol, and when to remember the irritable condition of the stomach. The introduction of soluble carbo-hydrates into common use is helping to settle the question. While it was a case of choice betwixt food that required digestion, and the alcohol which passed readily into the blood and was burnt in the organism, then the alcohol had the preference. But now, when so many preparations of soluble carbo-hydrates are at hand, it becomes possible to limit the amount of alcohol to the requirements of the system as to stimulant, other readily combustible hydro-carbonaceous matter being substituted for it as a food.

And this is a matter of cardinal importance when conditions threatening exhaustion are present. As for instance, when some fresh acute bronchitis is added to old-standing bronchitis and emphysema, or is super-imposed upon a condition of advanced mitral disease with permanent congestion of the pulmonary circulation. It has seemed to me that when the capillaries of the lung (which are normally of comparatively large calibre) have for long been distended by a block at the mitral ostium, they readily give way, or yield when some new force is in action. Hepatisation of the lungs is very apt to spread and extend where there is old-standing mitral disease, and then a condition of acute danger is developed, requiring the most watchful care, and calling not only for appropriate medical measures but for skilful measures of feeding. The system and with the general condition the digestive organs are in no condition to meet any demand upon them for assimilative purposes. Small quantities of readily assimilable food must be supplied to support the sorely taxed organism. As to the assimilation of fat, it is very doubtful how far it is possible at all in these conditions. A granule at the urine, high-coloured and laden with lithates, tells that the liver is in no condition to deal with large quantities of a fat material, even if the stomach could digest them. Then the patient has to be considered: fat is loathed, and predigested albuminoids are not enjoyed. How then, and upon what is the patient to feed? The needful grape-sugar must be supplied to it, else the organism will burn out like a fire imperfectly supplied with fuel. There are no doubt medical men to be found who regard with indifference the matter of food and its digestibility, and who are content with, but very little discrimination as to what these small quantities of these are, as a rule, men who have not deeply considered the

ject of digestion and its processes. The advanced physiological knowledge of the present day is demonstrating what it is the body requires when in straits, and how to supply its wants in a form calling for the minimum of the digestive act. They remind me of those persons who could not see the advantages of George Stephenson's invention of the locomotive; and only held a railroad to be inimical to the coaching interest and the breeding of horses. These persons will inevitably become rarer and rarer, as man is mortal—any "rarity is the precursor of extinction" as we know—to make way for younger men more intimate with the facts of physiology.

CHAPTER XXIII.

FOOD IN BRIGHT'S DISEASE.

ONCE more a certain amount of pathological consideration must precede the subject proper, this being especially necessary for the medical readers.

"Bright's disease" is a disease of the kidneys; it is commonly taught; and its evidence is to be found in an albuminous urine. This is about as accurate as to say that the Iberian peninsula is Europe, and the Ebro the chief river of Europe. "Bright's disease" of the tubular nephritis, or large white kidney type, and anæmic or lardaceous kidney variety, may fairly be spoken of as "kidney" diseases. But of the granular, cirrhotic, contracted, non-granular kidney—by far the most common form of Bright's disease, it is impossible to speak as a mere localised "kidney" disease. It is a matter of widespread change, in which the circulatory system has a marked share, and in which the kidneys are also involved. The condition is one of waste-laden blood; where the blood is surcharged with nitrogenised waste. There is imperfect action of the liver, which fails to convert the albuminous matters into serum albumen (for the nutrition of the tissues) and urea (the soluble form of nitrogenised waste). Instead of the latter there is a distinct tendency to form uric acid, the primitive form of nitrogenised waste. Mammalian livers normally form soluble urea; but when the liver develops a tendency, or rather and more correctly, reverts to the back to uric acid formation, then the kidneys suffer severely. Professor Geo. Johnson, F.R.S., put it happily in his well-known work on Diseases of the Kidneys: "*impaired digestion of the food, and consequent accumulation of the products of food in the blood, through the kidneys.*" As albuminoid matters as proteids enter the blood, this faulty digestion does not occur, and the proteids enter. Soluble peptones alone enter the portal vein, and are "faultily digested," we now know, as with the liver. The point of Bright's disease, then, may be said to be this:—How far naturally inefficient kidneys, or kidneys rendered inefficient as scarlet fever, by their inadequacy to extract the nitrogenised

crates, help to the condition of the blood, and to accumulation therein of excrementitious matter, is a subject on which opinions may differ. How far the first departure from health is mental worry or overstrain—which notoriously affects the liver and the assimilative organs injuriously—we do not yet know. Clifford Allbutt regards Bright's disease as having frequently such a mental origin.

When these mal-products are formed, and exist in excess in the blood, two things may happen:—

1. The kidneys may become injured by the output of these lithates; or

2. The excrementitious matters are retained in the blood, giving rise to lithiasis,—or gout in its widest sense.

Frequently and not uncommonly more or less of both is found. Certainly, gouty persons have gouty kidneys. But a well-marked condition of Bright's disease may exist which all physicians would class as Bright's disease, and not as gout. In such cases there is generally a certain pallor, (such is the Bright's disease seen so much in Germany) while a gouty aspect is more common in England.

In this widespread change the urine may or may not be albuminous: or only albuminous at times: or it may not be albuminous (Mahomed). Many hold that in this chronic granular condition of the kidney albuminuria is not essential to the diagnosis, and that, when certain other changes are notable, the absence of albumen from the urine carries with it no comfort. By the overwhelming importance attached to one symptom and the neglect of the other semela, a great deal of error, confusion, and misapprehension,—aye, and it is to be feared much unnecessary mental discomfort,—have been occasioned. Surely there is enough unhappiness in the world without adding to its mass unnecessarily. Albuminuria, too, has other associations than Bright's disease.

Such then is Bright's disease as ordinarily encountered.

I am old enough to remember when the teaching was—meet the drain of albumen by liberal quantities of albuminoid food. This, it is needless to say, was an outcome of imperfect acquaintance with the nature of the malady, and a superficial conclusion. It stood on the same chronological platform as the advocacy of the flesh of carnivorous animals for diabetes. It was at that time thought that the great matter was to keep the albumen from passing out of the body; and, when this could be successfully done, to introduce plenteous supplies of it.

Now a great change has come over our views. The teaching of

Good, Johnson has had a profound influence upon the theory, and the most recent works on the subject are based upon the view of nitrogenised waste by the blood in the case of Bright's disease. Prout did good work in this connection, and the younger men of this generation are all familiar with the view of chronic Bright's disease to the albuminoid element in the blood. Probably, too, they are all, more or less, of the opinion that in many cases a mental factor plays a prominent part in the origin of the morbid condition. In other cases the starting point is excessive indulgence in animal food, *deceit* beyond the necessities of life. Where there exists evidence of renal inadequacy, the diet becomes necessary to limit the work thrown upon the kidneys. What amount of evidence would be accepted as conclusive, and what as merely suspicious, but not proven, is a matter upon which opinions necessarily most probably would differ; and I am by no means disposed to step out of my way here to attempt to settle the point. If any case it exists, it will be well to err on the side of caution.

In the first place, then, it is well to restrict the amount of animal meat consumed. The flesh of the larger animals, beef, mutton, pork, no matter whether as adults or juveniles, must be taken in strict moderation; and with due regard to the needs of the body and its tissues, and not to the cravings of the palate, nor of the stomach, which leans to meat. The joint is best avoided, on account of the fat of it. Made dishes are often permissible, and especially those made which gratify the palate without taxing the liver. Baked and broiled, for instance, is a suitable dish, though some would take exception to the albuminoid element in the liver. Stews, with plenty of vegetable matter and fat, are preferable to joints. It is so, in fact, with legumes, which are rich in nitrogen, should not be a large constituent of the stews. Hashes with plenty of vegetable matter, as turnips, carrots, celery, onions, leeks, and with a limited amount of lean meat, are good and palatable. A vegetable soup is also a good one-course dinner. Provided the digestion is good, and is not contradicted. Cakes, crackers, biscuits, sweet corn, and the like are admissible. Savors should form a distinct element in the diet. Fruit, raw or cooked, may be taken by those with weak digestion.

Certainly there are some cases of Bright's disease, especially of the pronounced anuria, to which the above diet is not applicable, and where a more strict animal flesh diet is necessary, and even the bulk of persons with Bright's disease, where the dietetic principles of these remarks apply.

Fish in all forms, all kinds and however cooked, are admissible. But here again we must remember that fish is albuminous, and that unlimited fish even may be harmful. Shell fish and molluses form an agreeable change.

Soups of all kinds may be taken, a preference being given to those where the amount of "stock" is small. Those prepared with cream, and those thickened with farinaceous matters, are to be preferred. Soups can be made thick with vegetable matters, and such soups are good.

Persons with Bright's disease will do well to learn to eat fat. They should put cream (as understood in England) in their tea and coffee, eat plenty of butter on their bread, should be given to biscuits and butter at lunch and dinner; should have real melted butter to their fish; and should not be niggardly with the oil in their salads. A basin of porridge, either of oatmeal or hominy, followed by a rasher of fat bacon, or fish with melted butter, would form a suitable breakfast. Soup, as described above; some well-mashed potato or fish, followed by a vegetable entrée, with biscuits and butter, would form a desirable lunch. Then dinner might consist of soup, fish, an entrée, pudding of any kind not too rich in eggs, milk-puddings, blancmange, pastry, or creams are permissible when they can be procured; again, biscuits or crackers and butter, with a nip of a savoury cheese to finish with. Dessert might be liberal—though I am afraid some of the older practitioners will not approve of this indulgence. Then if some food were required later on, a plain biscuit and a glass of milk, or of wine, would be sufficient.

Of course much would depend upon the features of each case. As has been pointed out before in these pages—and may possibly even be referred to again—it is not what is consumed, but what is digested, that is the measure of nutrition. It may be found that even where some meat would seem indicated by the anemia it does not attain the desired end. I have at the present time a patient at the Victoria Park Hospital, who has other evidence of Bright's disease than mere albuminuria, in whom this is well seen. She is a well-built, strong woman, but pale. So sure as she eats any animal food, so certain is she to put in an appearance, complaining of not being well, looking ill, and with a quantity of albumen in her water; while she restricts herself to farinaceous food, vegetables, fish and milk, she is well. Again and again she is tempted by some foolish, well-meaning person to take some meat, and, being a good-natured obliging woman, again and again has suffered for her compliance.

Indeed I have found with a certain class of persons afflicted with this disease that a milk and farinaceous dietary (crisp bread, &c. bathed) has not only cured the urine of a chronic case, but that the sick persons feel better.

Still, there may occur cases where from special considerations a dietary may be removed with advantage, but then the alimentary system should be strictly watched. And if there be continuous morbid action it should be put in abeyance. Then a modified dietary may be of considerable and indeed of advantage to the individual. There is to the use of alcohol in Bright's disease, opinions differ. Some two or three years ago the matter of the part played by alcohol in the production of Bright's disease was hotly discussed, some very eminent authorities taking sides. Of course it may be laid down as a general proposition that alcohol has a bad effect upon the liver, and one will deny that, and that therefore when the working of the liver is impaired alcohol is contra-indicated. But then sometimes a little alcohol has a beneficial effect upon the system generally, and it may be questioned how far it can be wisely withheld, especially if good results ever result from its use. Any accumulation of facts in regard to it, if he cares to make the experiment, that a certain amount of alcohol will send up the proportion of albumen in the urine, though probably few will be tempted to test the accuracy of this statement in their own persons. Each person who is the subject of Bright's disease must study the matter for himself or herself. I am not of opinion that it is well to make a hard and fast rule to the effect that under no circumstances should alcohol be taken by a person with chronic renal mischief. This would be unwise. But as a broad rule, it may be said that when genuine Bright's disease is on foot it is well to keep a strict watch over the amount of alcohol taken, just as it is kept over the amount of animal food consumed. The majority is found among water-drinkers, and therefore has not a purely alcoholic origin. All the same, alcoholic indulgence will aggravate the state of matters when it exists, and therefore persons who take much alcohol who do not, on the whole, suffer lessened by their kidneys.

Without negating other causal relations, it may be said that the excessive and drinking habits of the majority of our population is an important part in the production of chronic Bright's.

With some persons a milk dietary is effective in removing the symptoms, and the removal of animal food is of great value. Milk and oatmeal gruel are not so good as milk and oatmeal porridge.

To remove the albumen is not, however, to restore the kidneys to a normal condition. Nevertheless, it is quite possible to stay the progress of the kidney mischief, and to arrest its march by wise and judicious measures, of which a careful, regulated dietary is a factor of cardinal importance. And in some cases it is possible to make the advance so slow as not to be measurable.

CHAPTER XXIV.

FOOD IN ALBUMINURIA.

IN putting albuminuria into a chapter by itself my design is to emphasize the fact that it and Bright's disease are not synonymous or convertible terms.

Albuminuria is commonly enough really the escape of peptones by the kidneys. Instead of being converted, or turned back into proteids in the portal vein, a quantity of peptones find their way into the general circulation and slip out of it by the kidneys, just as readily and as easily as they found their way into it by the portal venules. In such case the system is ill fed, and the individual is pallid; and a condition closely simulating true Bright's disease is set up. Such, indeed, usually are the cases of Bright's disease which are "cured;" for the real disease (as spoken of in the last chapter) does not admit of cure. Conditions of acute "Bright's disease" are, however, compatible with recovery; leaving the kidneys either free from permanent injury, or in a more or less disabled condition. Thus the consequences of scarlatinal nephritis are most varied. Even more permanent states are compatible with considerable length of days. But with conditions where the presence of albumen in the kidneys is not accompanied by co-existing evidences of perturbed health, the prospect is far less grave.

The following is an illustrated case in point: A patrician youth, who like many others of his class indulged a good appetite, had albumen persistently present in his urine. A hasty medical opinion to the effect that he had "Bright's disease" was accepted by his family; and in so far he was an object of interest to a large circle of friends. One day, when riding in a London park, he met with an accident, and, breaking a limb, was taken to a neighbouring hospital. There he was put upon the ordinary dietary, with the result that he lost his albuminuria. The case was greatly discussed in polite circles as a "wonderful cure of Bright's disease." This is an instance of the loose, inaccurate gossip that goes on about diseases all over the world.

When a diet is too rich in albuminoid matter the urine is apt to

be albuminous. When such albuminuria exists, it is but reasonable to suppose that some modification of the dietary is desirable. First, however, ascertain what the precise facts are, and if the dietary is really too rich in albuminoids. If this be so, it is well to reduce the amount, and put the patient upon farinaceous matters, fish, fruit and fat. The whole matter was too thoroughly gone over in the last chapter to need repetition here.

If, on the other hand, there is evidence of mal-nutrition present, it may be well to combine a liberal dietary with a course of tonic and haematics.

Mere albuminuria, as differentiated from Bright's disease, is not a bodily trouble which calls for special dietetic treatment, beyond what has just been said above. If too much labour in the digestion of albuminoids has been thrown upon the digestive organs, then we may find the nitrogenised matters slipping away in some form of albumen, just as they may be cast out as some imperfectly oxidised form of urinary product.

Indeed the whole treatment of a case of albuminuria, alike medicinal and dietetic, will turn upon the diagnosis; and for accuracy greater pains are desirable than are usually bestowed upon it, too commonly the worst being rashly assumed.

CHAPTER XXV.

THE FOOD IN DIABETES.

If the dietetic indications in mere albuminuria are comparatively unimportant, the case is widely different in diabetes mellitus, where they are of the gravest importance.

In diabetes the patient wastes, because the grape-sugar, the food of the body, is slipping away by the kidneys and so is unavailable for the body-needs. It is not the presence of the sugar in the urine which is of importance, though it would seem like it from the attitude of a considerable section of the medical profession. The patient is not ill because sugar is present in the urine, but because his body is starving, and he is tortured by the symptoms set up by the presence of an excess of grape-sugar in his blood. An urgent thirst impels him to drink copiously to wash this excess out of his system. There is some prospect of success in dealing with a salt-thirst by drinking; but with a sugar-thirst the case is widely different, as anyone who has had experience of it well knows. The distention of the vascular system adds to the blood-pressure in the hepatic arteries, and so increases the output of sugar from that viscus. The suffering patient drinks and drinks, and grows worse and worse too frequently; and unless some impression can be made upon the functional disturbance in the liver, the case goes on from bad to worse until the final end. In other cases the progress is less disastrous, and by proper dietetic arrangements the patient keeps the malady at bay and lives on— a harassed and miserable creature. In other cases a better career is found, the “*diabetic storm*” passes over, leaving a comparative calm behind; when there is more or less sugar in the urine continuously, but no very distressing symptoms of diabetes. In such cases any disturbance, as a stroke, or a needless worry, may bring on an acute diabetic attack. Such cases then the varieties of what may fairly be termed “*Diabetes*,” compared to mere “*Glycosuria*.”

It is well to bear the distinction in mind, when we consider an individual case where sugar is present in the urine.

In the management of diabetes proper treatment is of the greatest

cardinal importance. By indulgence in any food which will form grape-sugar, or rather from which grape-sugar can be formed, a distinct aggravation of the symptoms is at once set up. All carbo-hydrates must be carefully avoided. The consequence is that a rigid dietary must be laid down by the medical man, and observed by the patient. Indeed the latter is only too glad to secure relief from the torturing thirst endured. The dietary preserves the organism from perishing of exhaustion until the disturbance which sets up the diabetes is abated, if abatement be feasible. By supplying food material which does not form grape sugar, two distinct ends are served.

1. The organism is fed; also

2. The suffering caused by the presence of sugar in excess in the blood is relieved.

These are great matters, truly. But the question may be raised:—Does a strict dietary exercise any curative effect over the malady? Does it merely keep the patient alive till a cure is wrought? Or does it tend directly to that end?

If we believe in “physiological rest” to any organ impaired or injured, or functionally incompetent—and such a broad rule is recognised in medicine—then the non-carbo-hydrate dietary has a distinct curative tendency. Physiological rest as regards its glyco-genic function must favour the regaining of its lost or weakened power by the liver. It is at least a reasonable assumption, even if it cannot be demonstrated as a proved fact. Such assumption falls in with one’s experience on other matters medical.

Where carbo-hydrates are contra-indicated, then we find the dietary restricted to hydro-carbons and albuminoid with the alcohols, as ordinary alcohol or spirits of wine and the alcohol glycerine.

The chemist has striven to give directions to the baker in this matter, and by depriving wheaten flour of its starch, a gluten bread is made which can be eaten with butter. Gluten bread is not particularly acceptable to the palate, still with salt butter it is not very objectionable. Some gluten biscuits are the most unpalatable food I have ever tasted in a wide range of experience. Nothing but a monstrous fear of death, or some tremendously strong motive to live, could enable any one to get them down.

Then the oil of the almond is utilised as formed in almond flour, and still more in almond biscuits. These are palatable enough, and form pleasant eating.

Amongst of almond biscuits, a certain little matter may be related

which throws a strong light on the genesis of diabetes. Taking one day with Mr. Van Abbott, who is so well known in connection with almond food for diabetes, I asked casually, "What are your chief customers, Mr. Van Abbott?" His face assumed an earnest look as he replied, gravely, "Grey, worn, business men, looking older than their years."* The almond biscuit is rich in oil and therefore forms an admirable "fuel-food."

Then, of course bulk is grateful to the stomach and must be sought in the parenchyma of vegetables.

Before proceeding to the practical discussion of the food of the diabetic, it may be well to give Dr. Pavy's table of the various foods:

DIETARY FOR THE DIABETIC.

MAY EAT

Butchers' meat of all kinds, except liver, ham, bacon, or other smoked, salted, or cured meats. Poultry, game, shell-fish and fish of all kinds, fresh, salted or cured. Animal soups not thickened, beef tea and broths. The almond, bran or gluten substitute for ordinary bread. Eggs dressed in any way, cheese, cream cheese, butter, cream. Greens, spinach, turnip-tops, French beans,* Brussels sprouts,* turnips,* cauliflower,* broccoli,* cabbage,* asparagus,* sea-kale,* vegetable-marrow,* mushrooms.

(Those marked with an asterisk may only be eaten in moderate quantity, and should be boiled in a large quantity of water).

Water-cress, mustard and cress, cucumber, lettuce, endive, radishes, celery, vinegar, oil, pickles. Jelly flavoured, but not sweetened, savoury jelly, blanquette made with cream and not milk, custards made without sugar. Nuts of any description, except chest-nuts. Olives.

MUST AVOID EATING

Sugar in any form. Wheaten bread and ordinary desserts of all kinds. Rice, arrowroot, sago, tapioca, macaroni, vermicelli, potatoes, carrots, parsnips, beet-root, peas, Spanish onions.

Pastry and puddings of all kinds. Fruits of all kinds, fresh and preserved.

* Of the association between persistent night-sweats and increasing diabetes we are beginning to have some knowledge. It is already a fact, between some sugar-smokers and the appearance of the nocturnal sweating symptoms.

MAY DRINK

Tea, coffee, cocoa from nibs. Dry sherry, claret, dry Sauterne, Burgundy, Chablis, hock, brandy and spirits that have not been sweetened, soda water. Burton bitter ale in moderate quantity.

MUST AVOID DRINKING

Milk, except sparingly. Sweet ales—mild and old—porter and stout, cider, all sweet wines, sparkling wines, Port wine, unless sparingly. Liqueurs.

Such is a fairly complete list of foods forbidden or permissible.

Of course all the predigested carbo-hydrates as prepared foods are contra-indicated. However valuable elsewhere, they must be banned from the diabetic's dietary. On the other hand, all oil emulsions can be cordially accepted.

All fishes preserved in oil form admirable food.

With Dr. Pavy's material to hand, it is possible to construct the different meals of the day, it not being necessary to have all the articles mentioned upon the table at once. A variety, of course, must be provided.

BREAKFAST.

Tea, coffee or cocoa with cream, gluten bread and butter, almond biscuits, fish, with oil or melted butter, caviare, mutton-chop, rasher of bacon, cold meat, ham, game, pressed beef, tongue, water-cress, salad, pickles.

LUNCH.

Soup, lobster salad, potted trout, celery, raw or cooked, radishes, scrambled eggs, cream cheese. Claret, hock or sherry.

DINNER.

Salmon, with lobster sauce, brill, anchovy sauce, cod, shrimp sauce, herrings, sauce tartar, plain white fish and melted butter, trout with parsley butter (not flour paste), stewed mushrooms, pigs' feet. A joint of any kind with spinach, cabbage, cauliflower, Brussels sprouts, celery, celery, endive or cucumber.

Entrées. Sweet veals, pigs' or calves' cheek boiled, then cut in pieces and served up with a sauce piquant. Mayonnaise. Game of all kinds.

Salad and cheese, cream cheese, pickles. Creams (unsweetened).

Gluten bread, butter and cheese, almond biscuits, claret, hock,

Chablis, Capri, Carlowitz, Catauba, red Australian wines, — W. S. S. or brandy, with any aerated water.

By such means, variety—so essential to the palate—can be secured, and an ample sufficiency for the nutrition of the body, too, furnished.

And now to some criticisms of the rigid dietary:

In the first place, Dr. Pavy bans liver; yet he allows shell-fish, though the oyster and the mussel consist largely of liver. This is obviously only an oversight.

Then do not turnips contain some sugar? He states they do. That is another oversight.

The more serious criticisms are these:—Is it always desirable to subordinate the dietary to the diabetes? Are not sometimes slight divergences from the rule permissible? Though milk does contain sugar, is it always advantageous to bar it? Of course when the patient is gravely ill, any addition to the sugar in the blood is offensive and objectionable; but when the condition is less pronounced sometimes milk may be allowed freely, and the same may be said about ordinary bread. In one case lately, where the patient chafed at the rigid dietary, a little bread came in advantageously. The relief it gave to his stomach more than counterbalanced the extra sugar in the system.

But of course such deviations from the rule should be narrowly watched, and with a full consciousness on the patient's part that the change is an experiment which may, or may not, turn out successfully. What the writer desires to put clearly, if he can achieve this, is "That while it is necessary to adopt a rigid dietary in grave cases of diabetes, and when acute symptoms are present, is it necessary or desirable to pursue such dietary when the symptoms are quiescent? Is it, or is it not, the fact that many persons, fairly to be regarded as diabetics, suffer much more from their dietary than their urine, and that many break bounds without apparently suffering in consequence thereof? The histories related by various writers would favour this view in numerous instances. But, all the same, great caution should be exercised, and the effects of each individual article be narrowly watched.

The patient must be dieted in order to be kept as long as possible. Liver can regain its lost power as regards its glycogenic function, if possible. But how is this to be attained? It is to be gained by the avoidance of sugar and what will make sugar. The case of a patient, years ago, known as "the skimmilk treatment of diabetes," is a case

vigorously by Dr. Scott Donkin, had a large measure of success; and yet the milk sugar was not removed in the cream. The actual amount of sugar had something to do with the result, I feel certain. A glycosuric patient of mine recently sent me a sample of her urine for examination which was absolutely free from sugar. Yet at that time she was living upon vermicelli pudding and arrowroot—two farinaceous matters most likely to have produced an excess of sugar. There seems some ground for holding the view that the liver can deal normally with a certain amount of sugar, and a certain amount only; and any excess beyond this remains in the blood, giving rise to diabetic symptoms. But the subject is one on which light is coming in slowly, and time must elapse before we can see it in the full day-light of complete knowledge. Some years ago a retired army medical officer of eminence consulted me for severe attacks of angina pectoris, and on examination he was found to have a gouty heart with disease at the aortic valves. On taking down his history it turned out that some years previously he was (in India) testing the urine of a patient for sugar. He was in perfect health and strength, and wishing to be sure of his tests he determined to examine the suspected urine side by side with his own. To his astonishment he found his own urine heavily laden with sugar. So alarmed was he that he made all haste home to put himself under an eminent authority on the subject of diabetes. On this his health soon began to fail in reality, and after some months he grew so weak and ill that he begged for some raspberry jam, as he could not be worse than he was. His request was complied with, and he felt the better of the carbo-hydrate; and the result was he gave up his dietary, ate and drank what he pleased, and got well; ultimately losing his glycosuria entirely. There was no grounds for questioning the truth of his narrative: he was a man of known probity, belonging to a distinguished family. It is a curious fact that several brothers of this family have at times passed sugar. They are all gouty.

In what has just been written it is not intended to cast doubts upon the value of a non-saccharine dietary in diabetes—that would be absurd; but to point out that we have yet much to learn about the conditions under which sugar appears in the urine. Of the value of a suitable dietary, not only as a means of furnishing relief, but as a stepping-stone towards cure, in many cases, there can exist no doubt. Indeed in diabetes now, thus we see one of the most marked instances of the value of dietetics, and of the food treatment of disease.

CHAPTER XXVI.

FOOD IN GLYCOSURIA.

THE main reason for the existence of this chapter—as a separate chapter rather than as an appendix to the last chapter—is to keep glycosuria well apart from true diabetes. There exists such an unfortunate entanglement of the two, not only out of the medical profession but even within its ranks, that a complete divorce of the two is most desirable.

Glycosuria is quite common with corpulent persons, especially where some gout is present. In such persons, indeed, it seems a species of waste-pipe running off the surplus carbo-hydrates, and without which the body would become embarrassed with an accumulation of fat. Indeed the laying on of fat from the surplus of carbo-hydrates, while a condition of glycosuria is co-existing, is not unknown.

A marked instance of this came under my notice about a year ago, when a series of experiments with prepared foods (consisting mostly of predigested starch), instead of increasing the out-put of sugar, resulted in a marked increase in weight.

What the relations of gout and glycosuria are we do not precisely know. The one is a disturbance of the glycogenic function of the liver, the other of the second function of that important viscus the metabolism of albuminoids. There is some linked association evidently, though what its nature is still shrouded from us.

On the other hand, one of the great dangers connected with the abundance of animal food allowed in cases of diabetes and glycosuria is the production of interstitial nephritis, or Bright's disease. There is no proof that sugar in passing out of the kidneys does any violence to those organs, any more than it does to the portal vein by which it enters the circulation at every meal; but the large amount of albuminoids not only allowed but positively forced upon the glycosuric patient is certainly destructive to the kidneys. The other aspect of food, viz.: seeing that we do not do harm by it. It is not well to engraft a state of permanent injury to the kidneys upon a condition of glycosuria, which, as yet at least, is not nec-

victed of evil, even if it have been under some suspicion. That is the substitution of a morbid certainty for what is so far unproven.

There are not only the benefits to be derived from a correct dietary in certain maladies; there is the harm which may accrue from dietetic error. The kidneys are far from being beyond suspicion in gouty persons of middle age, and if a liberal meat dietary be adopted there is a distinct and measurable danger of aggravating the renal trouble—a most undesirable matter. The patient must choose (under medical direction) what he will do, whether he is more afraid of the gout or the glycosuria; and if he is a prudent, thoughtful man he will choose the glycosuria—as the less evil of the two.

If the teaching of this chapter be mainly negative, it is not, therefore without its value.

CHAPTER XXVII.

THE FOOD IN GOUT.

BEFORE proceeding to the subject-matter proper of this chapter, it may be well to clear the ground somewhat as to what is meant by the term, "gout." It once had a definite meaning, and was applied only to the deformities of the small joints of the extremities caused by the deposits of urate of soda in the articular cartilages of these joints. They were produced *goutte per goutte*, that is "drop by drop." So long as our ideas of gout were restricted to these obvious morbid changes all knew what was meant by the term.

But when the chemist discovered that the said deposits, or "chalk stones," consisted of urate of soda, the word began to have an extended meaning. Gout was seen to be a blood-disease, *viz.* that uric acid (as urates) circulated in the blood, and wherever the blood circulated there might gout follow. For instance, it was found that gouty deposits were laid down in the valves of the heart just as much as in the joints of the fingers and toes. Urate of soda is commonly formed in nodules on the pinna of the ear (Otolites.) Bronchitis has been found often to have a basis of gout. Certain skin diseases were recognised as having gouty associations. Then various disturbances of the action of the heart have been discovered as being outcomes of gout. Thus the range of the term was found to be constantly extending. From uric, otherwise lithic acid, being the *materies morbi* of gout, the terms "lithiasis" and "lithæmia" have been used to designate the gouty condition. They are more accurate terms, but it will take some time for the public to become familiar with them; whereas the vagaries and internal complications of gout are fairly generally known.

The various foods of the body, whether fuel-food or tissue-food, undergo within it an oxidising process which produces heat and force. The fuel-foods burn easily; the tissue-matters oxidise with difficulty. Worn-out tissue and surplus tissue-matters undergo retrograde changes which fit them for excretion by the kidneys, and to a less extent by the skin. These changes take place to a great

extent in the liver—the furnace of the body—the resultant products being cast out by the kidneys.

When kidneys first make their appearance in the animal world we find the form of nitrogenised waste to be uric acid. Uric acid, as urates, belong to animals with a three-chambered heart and a solid urine, *i.e.*, birds and reptiles. When the mammalia appear we find a four-chambered heart and a fluid urine: the form of nitrogenised waste being the soluble urea. There still remains, however, traces of the primitive urinary product, uric acid. So long as the liver can convert the waste and spare albuminoids into the soluble urea all is well. As the soluble urea the excrementitious matter is got rid of without difficulty, and albuminoids far in excess of the body-needs can be eaten with impunity. But when the liver fails in the formation of urea, and reverts to or falls back upon the primitive uric acid, a change comes over the scene. The comparatively insoluble uric acid lingers in the blood, finds a habitat in the articulations of the extremities, in the valves of the heart, on the pinnae of the ear, while it irritates the air passages and teases the skin. A certain proportion finds an exit by the kidneys, and worries these organs into interstitial nephritis, known as the cirrhotic or gouty kidney.

At other times the condition assumes a complexion which entitles it to be designated Chronic Bright's Disease.

When urates are formed by the liver one of two things happens: 1. the urates are cast out by the kidneys, which become injured by this out-put, and chronic kidney disease is set up; or 2. the uric acid remains within the system, giving gout (in some of its many forms); or, as very frequently happens, somewhat of both occurs, as has been said before.

From this we can understand what is meant by "the gouty habit" or "a gouty man" or "latent gout," and realise that the individual is the subject of hepatic reversion, with a tendency to the formation of uric acid. We can also comprehend what is meant by "gouty families," and why Cullen called gout "*Morbus Hereditarius*." When there has been great indulgence in animal food for generations the liver becomes more or less worn out by its toil, and the scion of such a family starts life with an insufficient liver. "The children of gouty parents ought more especially to follow the hygienic and dietetic laws laid down in these pages ("Nutrition in Health and Disease") if they wish to escape much suffering. As a rule they ought to be water-drinkers throughout life. They have to pay the penalty of their progenitors' excesses or dietetic errors,

If the gouty diathesis is strongly marked, they should "live a moderate and abstemious in their food habits, and lead as far as possible a muscular a life as possible." (James Henry Bennett). While, on the other hand, the humbly born alderman who started life with a liver in all its integrity, eats and drinks with impunity into old age before his Nemesis overtakes him; or, perhaps, himself escapes so free to leave a tendency to gout along with his wealth to his descendants.

By such consideration of gout (in the widest sense of the word) we can all the better understand the food-requirements of persons afflicted with this tendency to form uric acid. What the relations of uric acid are to "blue blood" it is impossible to say definitely, but both run in the veins of long-descended patricians. Gout has had a history of good eating at some time in the history of the individual or his race. The liver has become impaired by the labours it has had imposed upon it for generations. And this liver-failure brings with it in its wake all and sundry of those disagreeable consequences,—the various and numerous forms of gout. It also points out to us in an unmistakable manner the food to be adopted for the relief of this tendency to form an excess of uric acid.

1883.

LONDON FEVER HOSPITAL.

DIETARY TABLES.

ADULTS.	CHILDREN (<i>Under Ten Years of Age</i>).
1.	2.
Milk, 2½ Pints.	Milk, 2½ Pints.
Bread, Eight Ounces.	Bread, Eight Ounces.
3.	4.
Milk, Two Pints and Half.	Milk, Two Pints and Half.
Beef Tea, One Pint.	Beef Tea, One Pint.
Two Eggs.	Two Eggs.
Ice.	Ice.
5.	6.
Fish, Eight Ounces (Uncooked, Fresh Fish).	Fish, Five Ounces (Uncooked, Fresh Fish).
Bread, Twelve Ounces.	Bread, Twelve Ounces.
Potatoes, Six Ounces.	Potatoes, Four Ounces.
Milk, Two Pints.	Milk, Two Pints and Half.
Milk Pudding.	Milk Pudding.
Tea, Half Ounce, or Cocoa, ¼ Ounce.	Butter, One Ounce and Half, or Treacle.
Sugar, One Ounce.	One Egg.
Butter, One Ounce and Half.	
One Egg.	

ADULTS.

7.

Meat, Six Ounces (Cooked, and without Bone).
 Potatoes, Eight Ounces.
 Bread, Twelve Ounces.
 Milk, One Pint.
 Two Eggs.
 Pudding.
 Tea, Half Ounce, or Cocoa, $\frac{1}{2}$ Ounce.
 Sugar, One Ounce.
 Butter, One Ounce and Half, or Treacle.
 Cheese, One Ounce.

9.

Fish, Eight Ounces (Uncooked, Fresh Fish).
 Bread, Twelve Ounces.
 Potatoes, Six Ounces.
 Milk, Two Pints.
 Milk Pudding.
 Tea, Half Ounce, Coffee or Cocoa, $\frac{1}{2}$ Ounce.
 Sugar, One Ounce.
 Butter, One Ounce and Half.
 Two Eggs.

11.

Fish, Four Ounces (Uncooked, Fresh Fish).
 Meat, Eight Ounces (Cooked, and without Bone).
 Potatoes, Eight Ounces.
 Bread, Twelve Ounces.
 Milk, One Pint.
 Pudding.
 Tea, Half Ounce, Coffee or Cocoa, $\frac{1}{2}$ Ounce.
 Sugar, One Ounce.
 Butter, One Ounce and Half.
 Two Eggs.

13.

Meat, Six Ounces (Cooked, and without Bone).
 Potatoes, Eight Ounces.
 Bread, Twelve Ounces.
 Milk, One Pint.

CHILDREN (*Under Ten Years of Age*).

8.

Meat, Three Ounces (Cooked, and without Bone).
 Potatoes, Eight Ounces.
 Bread, Twelve Ounces.
 Milk, Two Pints and Half.
 One Egg.
 Pudding.
 Butter, One Ounce and Half, or Treacle.

10.

Fish, Five Ounces (Uncooked, Fresh Fish).
 Bread, Twelve Ounces.
 Potatoes, Four Ounces.
 Milk, Two Pints and Half.
 Butter, One Ounce.
 Two Eggs.

12.

Meat, Four Ounces (Cooked, and without Bone).
 Potatoes, Four Ounces.
 Bread, Twelve Ounces.
 Two Eggs.
 Milk, Two Pints and Half.
 Pudding.
 Butter, One Ounce and Half.

ADULTS.

Two Eggs.

Pudding.

Tea, Half Ounce, Coffee or Cocoa,

1 Ounce.

Sugar, One Ounce.

Butter, Two Ounces, or Treacle.

Cheese, One Ounce.

Beer, One Pint a day, or Beer

Money 1/- a week.

DIRECTIONS FOR STEWARD.

Beef Tea.—1 lb. of lean meat without bone to the Pint.*Strong Beef Tea.*—2 lbs. of lean meat without bone to the Pint.*Chicken Broth.*—One Chicken to each Pint.

One Fowl, 2 lbs. for Three Persons.

Mutton.—Leg and Shoulder clean and best end of neck, special order.*Beef.*—Salt Silverside (other joints, special order).*Rice Pudding.*—One Ounce Rice, Quarter Pint Milk, Half Ounce Sugar, One Egg for Three Puddings.*Sago.*—Quarter Pint Milk, 1 Ounce Sago, 1 Ounce Sugar, One Egg to Three Puddings.*Custard.*—Two Eggs, Half Pint Milk, Half Ounce Sugar.*Bread Pudding.*—1 oz. Sugar, Bread, One Egg for Four Puddings.

When the poison of gout was first discovered to be an acid, a curious instance of careless and slovenly reasoning was furnished to us. That sugar will ferment, and turn into an acid, is a notorious fact. The poison of gout is an acid. Therefore sugar must be eschewed. This illogical conclusion, however, derived a certain sort of *se-hantia*, or colouring of reality, from the fact that with some gouty persons sugar does turn acid in the stomach, and very quick a too. But because such is the case with certain dyspeptic gouty persons, it is not well to lay down a broad rule thereupon applying to all gouty persons. As a rule it may be affirmed that gouty persons are not partial to sweet food, and pass the sweets in dinner with very few exceptions. But all the same, rule and is not derived from the saccharine elements of our food. The old school who have sugar are giving way to younger men, who find the chemical-physiological changes in the nitrogenised elements of our food.

In the present consideration the gouty is not a child, and requires all the resolve appropriate attention when his digestion is disturbed in a proper manner. To him a child, or her, would run off the subject for more complicated than is desirable.

Then again some persons object to fat for gouty people. On what their objection is based is unknown to me. For my own part my experience, personal and other, runs on all fours with Ebstein's directions to eat fat. Once fat has passed the stomach, all trouble connected with it is apparently over. It is burned readily, nor can the objection lie in the tendency of fat to make fat in the body. For we have every reason to hold that fat does not make fat, and that the fat in the body is derived from the surplussage of carbohydrates.

The food element which furnishes uric acid is, then, the albuminoid, or proteid matters. The flesh of animals is the main source of gout-poison. It is not possible to dogmatise on this matter; but on comparing notes with a number of good observers of extensive experience, I find there is a considerable agreement as to the flesh of the larger animals being gouty. The white meats, as chicken, do not seem to possess the same strong tendency to pass into uric acid as is found with the brown meats. Caseine, whether in its animal or vegetable form, seems very free from this tendency to uric acid formation. Of course what has just been written must be taken for what it is worth—neither for more nor for less. An impression may be fairly correct in the main even if it cannot be represented in figures. A diet of lentils, if rigidly adhered to, is a very valuable matter in cases of confirmed gout. But this fact does not rest solely upon the comparative freedom of caseine from a tendency to uric acid formation; there is another important factor at work. Such food does not tempt the palate to excess; and thus the liver gets that physiological rest—as regards its function in the metabolism of albuminoids—which is so important in the recovery of impaired functional capacity. No doubt many gouty persons would get great relief from their tormentor if they could adhere to a dietary of lentils; but then human endurance falls under such a gastronomic trial. The late Earl of Derby, it is said, once had some sherry sent him, which, it was asserted, "would not make gout." He returned it to the wine merchant, saying, he "preferred the gout." So it is with a dietary of lentils. The Hebrew Patriarchs had flocks and herds, but they lived on milk and lentils and farinaceous matters. Indeed Isaac marked an important family ceremony by first partaking of savoury meat. The Zulus, who possess a most magnificent physique, live on milk and "mealies," or cooked maize.

When we come to the practical details of a gouty person's dietary,

we must keep the cardinal matter of the avoidance of *excess* well in view.

Soups need not be forbidden, as the amount of govtage is not very great, and possibly little of it passes out of the alimentary canal. White soups made with cream are good. Vegetable soups are excellent, though a caution may be required about those made from legumes, albeit the albumen is caseine. Fish soups are permissible. All soups may be thickened with broken biscuit, or contain quantities of vegetable matter, as *Julienne*, *hotch-potch*, *Spring soup*, and *green-pea soup*, *asparagus soup*, *corn soup*, *pea and tomato*, *turnip and potato soups*. Many persons prefer a quantity of onion, and others celery in their soup, both of which are unobjectionable from a food point of view.

As to sauces there is little to be said. Fish forms a large portion of the dietary of the gouty. Fish does not differ from the flesh of animals in chemical constituents so much as in its comparatively small food value. The amount of actual proteid matter in fish is calculated at about one fourth that of animal flesh. The consequence is, a comfortable, satisfying meal of fish does not overload the liver with albuminoid material, as would be the case with an equal bulk of the flesh of warm-blooded animals. The difference lies in the proportion of water, which is very large in the flesh of fish.

Many soups are made from fish. First comes the attractive *cray-fish soup*, after which follows at a short interval (gastronomical) *oyster-soup*. *Lobster-soup* is good. A white soup is prepared from any of the white fishes down to the *conger-eel*, and notably the *haddock*, which is admirably adapted to the body needs as well as the palate of the gouty.

Then there are cooked fish, the whiting, the haddock, the cod, the turbot, the brill, the skate, the halibut, the shad, the John dory, the gurnet, and lesser known fish. The coarser fish are the eel, the hake, the chub, the bream, and of course the pike. The trout, the salmon trout, the sewin of South Wales, the lake trout, the salmon, the grayling, and the white lake trout of the St. Lawrence are all excellent. The mackerel, the herring, the *recommode*, *cod* are too rich in oil for many stomachs, but otherwise are harmless. These fish may be boiled, or baked or fried, or made into *croquettes*, *fish-balls*, *fish-cakes*, etc., with potato or served upon *croutons*. The smaller flat fishes, the sole, the flounder and the *plaice* are good, but if fried should be prepared with care, and even then, when eaten are more offensive to a tetchy stomach than when boiled or baked.

with the fibres soaked in fat. "This is the source of many an attack of acute indigestion, or "even gout at the stomach."

The shell-fishes, as the oyster, the clam, the mussel and the scallop, are all excellent. So are the lobster, the cray-fish, the shrimp, the prawn and the crab. Many other shelled-fishes, molluscs and segmented fishes, are eaten in different parts of the globe; while the Chinese prize highly the sea-slug. The edible snail has its votaries, but this is not "a fish."

To a gouty man of perfect digestion a lobster-salad is a typical food. Its food-value in nutritive material is not high, while of its acceptability by the palate it is needless to speak. Fish in many forms lends itself easily to curries; and a well made fish curry with plenty of rice is an excellent dish for a gouty individual.

As to the flesh of animals, the great matter to remember is that this is a very concentrated food, rich in albuminoids and therefore to be eaten sparingly—as regards the lean at least. The question of digestibility is not being considered here, but the different forms in common use may be arranged pretty much as follows, mutton, beef, lamb, veal, pork; the last two being partaken of cautiously. Bacon and ham seem less objectionable. When these concentrated animal foods are eaten they should be in company with a fair store of fat, and a liberal supply of vegetables of all kinds. The rabbit furnishes a capital change, especially when cooked with a quantity of fat pork or bacon.

The same may be said of the flesh of fowls of all kinds, those with white flesh being to be preferred to those with brown meat on their bones. Thus the chicken is better than the duck or goose. Turkey seems to be midway.

Then, as to game, the same may be said of white and brown meat, whether in skin, fur or feathers. A little game to finish off a meal will scarcely be objected to by the most rigorous food-disciplinarian. But again it is a question of bulk, and little must be taken.

Then as to strong matters as sweetbreads and other glands; tripe, pig's foot, pig's cheek, souse, or as brawn, liver and bacon, they are all permitted in proportion to the bulk of fat, and the small quantity of lean in the dish. Ox-yeal head is not objectionable. Vegetables form a very considerable factor in the dietary of the gouty. Some objection might be taken to mushrooms and other fungi because of their meat-like composition. Vegetables containing a quantity of oxalic acid disagree with many gouty persons, who stand of course on old rhetoric, some, tomatoes, asparagus and even

broad beans. But for ordinary gouty persons potatoes, turnips, carrots, turnips, parsnips, beet, broccoli, Brussels sprouts, spinach, spinach, endive, lettuce, beans, peas, haricots, green corn, and yellow corn, kale, leeks, onions, salsify, sea-kale, samphire, asparagus, vine shoots, cucumber, gherkin, vegetable marrow, and other generally used vegetables, are excellent. Many admit of being deliciously prepared, so as to furnish that desirable matter, viz. vegetable Salads should be freely eaten.

Puddings are to be measured by the amount of eggs used in them. The egg is rich in albuminoids, and therefore objectionable to the gouty. All milk puddings as rice, tapioca, sago, semolina, &c., flour, malt-preparations, and other farinaceous matter should be prepared without eggs, or at most only the yolk should be used.

Suet dumplings are not to be forbidden, or bread and butter pudding, if the cook will make it without eggs. But custards, and omelettes, sweet or savoury, are to be eschewed. Pastry is not to be so strictly forbidden, nor yet cakes, if the amount of egg be moderate. All kinds of creams are permissible, but blanc-manges, sauces, jellies, lemon sponge, and articles of food containing much gelatin are to be avoided.

Fruits of all and every kind, raw or cooked, are suitable, whatever views may once have been held about sugar. A quantity of dessert fruit is desirable, except for those gouty persons with whom sugar disagrees and turns acid. Pale yellow lemon juice, and lemons, oranges, seedlings, and the like subacid fruits, are at present rather in vogue with persons with a gouty or rheumatic tendency.

Then, as to cheese and cheese preparations. Every preparation consisting of caselle, cheese in large quantities is not permissible, or is similar. Such dishes as toasted cheese, Welsh rarebit, cheese soufflé, or macaroni, consisting of cheese and eggs, are inadvisable. Macaroni, cheese and cheese stews may be eaten in moderation. For those who hanker after the flavour of cooked cheese, a small portion is permissible. Hominy, sago, or rice well boiled, and mixed with cheese freely grated, may be eaten to the satisfaction of the browser, and is a very good and digestible soup, or farinaceous. Not specially large quantities of uncooked cheese to be eaten.

Cheese omelette, which contains only a small amount of cheese. Biscuits, crackers or riced bread, with butter or oil, and cheese, should form part of the diet of the gouty. Nuts, but being eaten, would be too hearty of course. The following are good:—Ginger-bread, Nonchalant, Cannon-bread, or Brio.

The small green Swiss cheeses mixed with herbs and grated on bread and butter are unobjectionable. The poorer skim-milk cheeses are contra-indicated.

As to milk itself, it should be taken in moderation, unless it is taken as the sole food—a regimen which agrees with some individuals.

As to dessert, nuts and fruits, fresh or dried, are permissible for those with whom they agree.

And now as to beverages. Few gouty persons get on well with malt liquors. Ales and porters brewed after the English fashion are so much poison to gouty persons of thin flank. Bulky, gouty persons can only indulge their liking for such malt liquors by a compensating abstinence from the flesh of animals. No gouty man can drink malt and eat meat at the same time without paying the penalty for such indiscretion. Beer brewed after the German plan, as lager beer, does not upset the liver to the same extent as English ale; and many gouty and bilious persons can enjoy a draught of lager, whether Bavière, or Milwaukee, without suffering therefrom. In my opinion, indeed, the lager system of brewing will ultimately prevail,—at least among persons who have to be careful as to what they eat and drink.

Wines, too, differ in their qualities. All good generous wines must be given up. Old Port, Madeira, brown Sherry, Tokay, Constantia, Chateau Yquem, Bueellas, all must be abandoned. Port, old in wood, is less pernicious than old bottled port; but most gouty men eschew port wine sooner or later, no matter how fond of it. Champagne, Moselle, Hock, and Burgundy are scarcely less harmful, and I have seen gout simply blaze out in a few hours after Maecobrunner and Rüster.

A poor, thin wine, as the poorest claret, or corresponding Italian, Spanish, or Australian wine (I do not know the wines of the Western Hemisphere and therefore cannot speak of them), is all the vinous drinks a confirmed gouty man can take with impunity. He may still pride himself on his cedar—but it is for his guests, not himself!

A little spirit, brandy or whiskey, Scotch, Irish, Bourbon, Old Rye, or Monongahela, with water, is the best drink beyond doubt. Mineral and aerated waters may be taken freely, and help to wash the uric acid out of the tissues.

When a gouty man's palate fails him, it is rather a fortunate accident. So long as the pleasures of the table are regarded as among the good things of life, so long is the tendency to form uric acid fostered and favoured. An old clergyman was known to say "I've

been a dyspeptic for fifty years. "Thank God for it!" The cause of his pious gratitude for what most people regard as a curse, was the fact that his brothers had all perished from gout, while his dyspepsia had prevented him from overworking himself.

When the liver manifests a return to a reptilian condition, its owner must alter his dietetic and gastronomic performances, or suffer the consequences.

There is no nice way of having the gout; and, what is worse, there is no pleasant way of avoiding it.

CHAPTER XXVIII.

FOOD IN NEURAL AFFECTIONS.

IN dealing with affections of the nervous system there is one point which must never be forgotten, and that is the supreme importance of fat. For long the Germans have recognised this fact, especially in reference to the treatment of neuralgia by cod-liver oil.

This matter of the relations of fat to the nervous system receives an increment of interest from the physiological fact that the nervous system is rich in a certain phosphorised fat known as Lecithin with a formula $C_{41}H_{100}NPO_8 \times 3H_2O$. As pointed out before, in the chapter dealing with anæmia, there are only two bodies actually made or (as M. Foster puts it) "manufactured" within the animal organism, and which are not found in the vegetable world.

These are linked with the function of the blood-corpuscles and with the nervous system, two matters with which vegetable life has nothing to do. The one is Hamoglobin, which enters into the composition of the red blood-corpuscles; and the other is Lecithin, "a conspicuous component" of the brain and nerves. Just as under certain circumstances we find some persons becoming anæmic, and the red blood-corpuscles either are not made, or break down, or may be somewhat of both; so in others we find the nervous energy waning. How far this is due to exhaustion of the store of Lecithin is a matter on which as yet we can only speculate. But the hypothesis that such is the case stands in a curious and suggestive relationship to the clinical facts that fat and phosphorus are so serviceable in the treatment of atonic conditions of the cerebro-spinal system.

In the busy struggle for existence of the present day, when, as Sir J. Crichton Browne has pointed out, "the battle of life is no longer fought with thews and sinews but by the brain," the demands made upon the nervous system are excessive. "The spirit indeed is willing but the flesh is weak." Certainly the spirit is so very willing that the flesh must give way, even when not weak. It is all very well for Clifford Allbutt, F.R.S., to charge his neurotic patients with a lack of endurance. Whatever powers of endurance the body

may possess the restless tenant would exhaust them. The sword will wear out the scabbard! And especially is this true of those small or slightly built persons with well-varnished foreheads and small visages—the Arab type of beings of the neurosal diathesis. Their restless energy brings about, sooner or later, a break-down. The assimilative organs cannot supply the nutritive pabulum of the nervous system in sufficient quantity; the store becomes exhausted and then the brain-power gives way; the work that once could be accomplished with ease becomes a grievous task, and a little later on the capacity to discharge it is no longer present. There is a distinct break-down; and the poor patient, bereft of his whilom power, is sent away for rest—to do nothing but recuperate. Such a story is sadly frequent at the present time, and threatens to become even more common in the future. After a long rest and appropriate food the brain is once more equal to its labours.

In such a case the individual has distinctly drawn a bill upon himself—borrowed so much of his intellectual capital; and then comes when that bill must be met; after which follows a long slow process of paying back into the body-bank till the working capital is once more sufficient for competent action. There has been a body expenditure in excess of body income, and so the reserve body-capital has been heavily drawn upon, until indeed it is too small for successful enterprise. By reducing the outgoings to the minimum, and raising the incomings to the maximum, a new balance or capital is regained. It is just like a man who exceeds his income; he gets into debt, and then he has to live on very little until he has cleared off his debts;—and then he can go on again. When the pabulum of the brain is exhausted a long process of recuperation is necessitated.

Having secured for such a patient the requisites for rest and recuperation, the next question arises, viz., How is he to be fed? Experience has taught us the fact, before physiological chemistry told us the why, that fat and fish are the foods specially indicated. Fish is rich in phosphorus. A phosphorised fat has to be furnished to the nervous system. But to supply in liberal quantities of cod-liver-oil, tanning phosphorus and fat, is not necessarily to restore the depleted assimilative powers the capacity to build up and assimilate the new materials. It is one thing for the contractor to supply the materials for a house; there are the masons and bricklayers, the carpenters and painters, the slaters and gables, the joiners and the fitters of the required for its erection. If these are not given to the work

of construction will proceed but slowly, no matter how much material is to hand. So in the body. If the constructive powers are but feeble, then time is essential to the accumulation of a sufficient store of lecithin for working purposes.*

There are the capacities of the assimilative and constructive powers to be calculated in each case. When they are fairly vigorous comparatively rapid progress is possible; when they are gravely impaired, and still more when the individual has but small viscera, the recuperative process is and must be a slow one. The digestibility of the food must not, then, be lost sight of in the calculation. Small meals of suitable food at short intervals are the line of feeding to be taken. Then fish with butter should form a conspicuous factor in the dietary. Milk puddings are good. As a beverage cream with seltzer water, or other aerated water, is capital. Cold coffee with cream forms an excellent combination. Creams of all kinds are good. Cod-liver oil and oil-emulsions may best agree with others.

The yolk of an egg in a little sherry would make a little change. Salads with oil suggest themselves. Stewed fruit and cream ought to form a staple dish.

When we consider that the pabulum of the nervous system is a phosphorised fat, we can comprehend why the plan of treating cases of cerebral exhaustion by liberal quantities of lean meat has turned out a failure. Albuminoids do not supply the requisite material for the intended purpose: while in their metabolism they furnish matters which may be called hepatic mal-products, or "liver-stuffs," which possess very irritant or toxic properties as regards the brain. Consequently a highly nitrogenised dietary is not only without advantage, but actually possesses positive drawbacks. The brain is not fed thereby, but in its weakened condition is annoyed and vexed by these "liver-stuffs."

This matter of the toxic effect of liver-stuffs has another aspect which is well worth consideration. An American patient told me that, after the perusal of my work on "Indigestion and Biliousness," he found that he lost his bilious headaches by striking out of his dietary the flesh of animals except fish. He lives on a milk and farinaceous dietary, with butter, peas the fish. On this restricted regimen he has excellent health, feeling quite well. But any diver-

*What is here written must not be taken as accepted demonstrable fact, for it rests on a working hypothesis, and here even without its utility.

gence from this dietary brings with it its own pain, and Dr. A. Hulg, one of the casualty-physicians at St. Bartholomew's Hospital, has put on record a similar experience of his own, arrived at quite independently. Long ago, when attached to the West London Hospital, a patient came to me whose expression told of continual suffering. She was tormented by severe and almost ever-present bilious headache. She averred she could only take a little lean meat, dry bread and tea. She was prevailed upon to give up this regimen, not without considerable difficulty, and try farinaceous matter and fat. Much to her surprise she soon was considerably better. Ten years later she turned up accidentally, when it was found that the improvement had been maintained, and her face was free from the "pained" look it bore when first seen. The theory and the clinical facts hang together at any rate, if a more positive expression of opinion is not yet warrantable.

Some persons, and especially women, of highly developed nervous system and small digestive power, are liable to recurrent storms of neuralgia, specially facial neuralgia. Sometimes it is facial neuralgia (as ordinarily understood) at other times there is intense pain in the eyeball with hemimeralgia.

When the storm is blowing they can be got to take medicine, but food they cannot be induced to take. "Pain is the prayer of a nerve for healthy blood," said Romberg; and this exquisite sentence is well worth bearing in mind. The blood must be good in quality and sufficient in quantity, else the nerve gives out its inarticulate cry. One sufferer of this kind came under my notice lately. Milk was objected to and meat infusions alone were available. But such infusions, as pointed out elsewhere, have (practically) no food-value. Some "inert food" was available, and by the addition of it to the meat-infusions the strength was maintained. Not only did the system derive support from the food to bear the suffering, but the nerve, being better fed, sooner ceased its wall. Shortly the opportunity offered itself for a second try, to press her new convictions upon a friend suffering in a somewhat, and with the same satisfactory result. When the patient is a chronic sufferer can only object to every proposition, and be set to work at once. The patient, however, when so suffering, is not a competent judge of what is best to be done, and must be guided by the doctor. One patient suffered severely from a sudden cold, and catarrh of the nose. The cold air of the night caused them to wake, and the cold air of the

neuralgia. Some food in the small hours of the morning warded off these morning face-aches in a most satisfactory manner.

(The effect of cold air in the production of these face-aches seems as follows: Cold causes the cutaneous vessels to contract, and the cutaneous nerve-fibrils thus have their blood supply cut down, with the consequence of neuralgia in the starved nerve. Protection against cold is, of course, one preventive matter; while a sufficient supply of nutritive blood is another. Our ancestors in draughty houses had bed-curtains and wore night-caps. No doubt this saved many a face-ache. Their descendants, with curtainless beds, and ignorant of night-caps, suffer for their advance in taste. The facial neuralgia which has its origin in a cold drive is a similar case of nerve-starvation.)

This view of the relation of food to neuralgia, and especially superficial neuralgia, is well worth bearing in mind. Such neuralgia belongs to persons with small viscera, and who can only take small quantities of food at once. They cannot endure the long unbroken fast of night, and should have something last thing on getting into bed, and also something first thing in the morning before commencing to dress. In cold weather, beyond this, they must have some food in the small hours of the morning. This may consist of milk with some malted preparation; or of a meat infusion with some predigested food. This may be kept warm under a tea-coscy; or on the hob, if the fire be kept up, or a spirit lamp may be used.

And, by attention to the matter, many neuralgic sufferers will be able, if not to escape their plague altogether, still to avert much of their pain. At times alcohol in some form may be indicated as an addition to the food just mentioned. Sometimes indeed a little alcohol seems necessary to digestion in the small hours of the morning.

Dr. King Chambers describes a curious neurotic condition, linked with an erroneous dietary thus:—“There is a peculiar form of *hypercholesterolemia* which arises from eating too little vegetables and too much meat. It is distinguished by the high specific gravity of the urine (1.025 to 1.030) dependent on the presence of urea alone without sugar. There is, in the cases often a remarkable lassitude, accompanied by a paralysis of the limbs suddenly occurring after exertion. Sometimes there is emaciation. Both these symptoms usually go to the patient and his friends to attribute the morbid state to insufficient nutrition, and to increase much and more the proportion

of meat in spite of the aggravation of the ailment. "A rapid recovery attends the diminution of the meat meals to one daily, and the supplying their places with plenty of porridge and green vegetables."

There is another piece of evidence, that while the ferments produced by normal digestion are agreeable—hence the practice of charity-dinners—when there is an excess of albuminoids in the dietary, or the liver is unequal to its work, mal-products are formed which possess a toxic effect upon the nervous system.

CHAPTER XIX.

FOOD FOR CHRONIC INVALIDS.

As might, *à priori*, be expected, the dietary of a chronic invalid differs materially from that of persons in robust health. Yet this obvious matter does not always receive the attention it merits, and the invalid has often "to take what is going," and which is adapted to the body-needs of the rest of the family. Of course this does not occur in thoughtful families; but is apt to be the case where one member of a robust family returns to the family group in broken health. It is all very well for a family to enjoy unbroken health; but lack of familiarity with illness carries with it certain drawbacks; and of these, ignorance of what an invalid requires, is one. There is no intention to neglect the sick member, or disregard his or her requirements; it is simply a case of want of knowledge. Whereas, on the other hand, in families where illness is not unknown, an acquaintance with the wants of sick persons is found. This is, in the hour of need, the silver lining to the dark cloud of illness. Few physicians but have had the above facts forced upon their attention from time to time. Called to a sick person, in one household all are ready to lend their services, but the willing volunteers are undisciplined mobs, destitute of any knowledge of what has to be done, and how it has to be carried out; while in another family the physician finds the different members ready to fall each into his or her place like well-disciplined soldiers. How they have purchased their knowledge it may be well not to inquire; but it is there when it is wanted—and that is something.

With the chronic invalid the position is somewhat different. That one of the females of the household which has the kindest heart, the best head, and the most willing hands, takes upon herself the task of tending the sick member. But no amount of willingness can compensate for the lack of actual information.

No affection for instance can ward off the evil effects of bread crumbs in the bed, which soon become hard, gritty particles readily productive of bed sores. Ready and affectionate as she is, this devoted woman, has her lesson to learn, and probably learns it—as

much other knowledge is acquired through years of experience, long-taker; but all the same it has its disadvantages. The cook, or the nurse must pore over her cookery book, and select the best she finds there. If some of the details seem unnecessary, or superfluous, unnecessary, she will do well not to exercise her right of interpretation and make modifications of her own. It is dangerous to prophesy about viands. The proof of the pudding is in the eating. And it is not well to assume that some cooked product will be to one's taste. It may turn out far otherwise. Taste and try. Paul's advice to the Thessalonians to "Prove all things, hold fast that which is good," applies here very pertinently. She should make a private trial of each dish before serving it up to the invalid. She may be puzzled what dishes to select. It will be well for her to give her attention to light rather than solid food. The invalid is not equal to much exertion and therefore does not require the substantial repast which is desirable for the hale members of the family, and which they enjoy. The body-requirements are smaller. On the other hand lack of exercise leaves the appetite unimpaired, and the zest for food is not so keen. Something tempting is indicated: a small bird on toast, a dainty sandwich, a herring's roe, or a bit of cod's roe, a sweetbread or a little fish, a small quantity of a milk pudding, or a little stewed fruit and cream, or a custard in a glass, something small in amount and tempting in character.

Very often the cook dislikes the notion of these "frivolous temptments," in which case it is better to try to prepare them independently of her than to persist with her. No woman yet ever made a dish palatable which she prepared unwillingly. A cook must give her mind to her dish— if it has to be a success.

If the volunteer experiences a difficulty in her selection, she may find considerable help in a little book "Food for the Invalid, the Convalescent, the Dyspeptic, and the Gouty," published by Macmillan and Co., containing some three hundred out many pages, and selected from various cookery-books and compiled by the author.

She must learn, too, to prepare small quantities at a time. Few foods improve by keeping, and few bear warming up without losing much of their toothsome-ness. Sometimes the turn of mind of the invalid is such that a dish of which the invalid may fancy a little, or even a small portion, something special must be prepared. The invalid is not always eating out a feeble appetite, or even being dainty. He may, to all doubt, the invalid bears much quietude with the food, and nothing about it. Perhaps he is peevish. There is a great deal to be said

As the amount of food which can be taken at once is small, the invalid must not be restricted to the family meals, but have little gastronomic interludes. A cup of milk, a cup of beef-tea with some prepared food in it, a little custard, the yolk of an egg in sherry, or some chocolate, or possibly even some sweets. But here we have to confront a difficulty, viz.: How far do such minor meals destroy the appetite for the regular meals? This must form a matter for private judgment in each individual case.

Sometimes the invalid may like highly seasoned dishes, but as a rule plain fare with condiments in moderation is preferred by sick persons. As Sir Wm. Roberts has pointed out, the taste for savoury food is diminished in sickness, and simple food, unattractive to hale persons, is relished by those who are not well.

In consequence of lack of exercise there is always a tendency towards sluggish action of the bowels; and this can usually be met by a baked apple, a few stewed prunes, or figs occasionally. Especially is this necessary where milk forms a considerable portion of the dietary. Meat-infusions have a tendency the other way. According to the state of the bowels, milk or meat infusions must be resorted to.

Then another matter is the avoidance of fat by invalids. In all illness fat is objectionable to the palate. Whether this is an instinctive avoidance of an article of food which cannot be digested or not, may not be affirmed. But of the impairment of the pancreatic function in acute disturbances there is much proof. This, however, is certain, viz.: that the necessity for such concentrated fuel-food as fat is much diminished in the case of invalids. Confined indoors to a great extent, and incapable of much exertion, there is not the demand for hydro-carbons which is normal in health.

(Of course this statement does not apply to strumous and phthisical persons, who require a certain amount of fat to build up healthy tissue).

Salted and dried meats are rarely relished by invalids unless it be in the case of bacon and ham. Tinned or canned meats must be used at once, else they turn against them. All greasy meats are disliked and dry meat preferred: thus a chicken is in request, while a succulent duckling is rejected. Boiled or baked fish is preferred to fried fish, as a rule. An egg is a favourite food with invalids usually. Plain boiled, or boiled lightly, and turned out into a warm cup containing a little knob of butter melted with some pepper and salt, either are attractive. Or a light boiled egg, turned out upon

some nice mashed potato and mixed therewith, form a more acceptable to the palate of an invalid.

As to drinks and beverages it is impossible to say at present, so much depending upon the likings and dis-likings of different persons, and so much upon the actual requirements of each individual.

Great cleanliness and neatness in the serving of food is essential, otherwise the appetite flees at once. No attractiveness of viands can compensate for a dirty plate or a soiled cloth.

CHAPTER XXX.

FOOD IN OBESITY.

CONTRARY to what might be expected, fat eaten as food is not deposited in the body as fat. Part of it is utilised in tissue-formation, while the rest is burnt. The fat of the body is the surplusage of carbo-hydrates beyond body-needs. Of course it is obvious that if the dietary be rich in fat this will burn to maintain the body-temperature, and thus the demand upon the carbo-hydrates consumed is lessened, and the more remains over for adipose tissue.

Fat is deposited in the body in an irregular manner: sometimes evenly distributed, at other times certain localities are selected. The eye lies in its orbit with a packing of fat around it. The salmon has its fat in its tissues, the cod locates its fat in its liver. Some persons are fat all over like a fattened animal; while others carry their fat in a thick pad over the wall of the abdomen. The camel, the dromedary and the buffalo have a hump of fat on the back.

The King of Dahomey's wives are fattened on rice and butter, previous to the nuptial ceremony. A certain plumpness is admired in some animals; and a reserve store of fat is often very convenient, carrying with it a certain staying power. A sleek, well-fed horse is an instance in point.

On the other hand fat may be a burden, as many obese persons know full well. It certainly may be so excessive as to be unsightly; and fat persons are regarded as legitimate objects of ridicule by the vulgar. Consequently, many and divers plans have been laid down for the reduction of redundant corpulence.

Lewis Cornaro, who was a corpulent, gouty man, put himself upon a spare dietary, viz., twelve ounces of food, mainly vegetables, and fourteen ounces of light wine daily; and on this lived in good health for over half a century. Few persons, however, could undergo such a severe regimen. Another obese man fed largely upon apples, on which he reduced himself considerably. The great idea in the different plans of treatment is the avoidance of carbo-hydrates in the food so far as is possible.

There are two plans before the public at the present time: (1) Bantingism, and (2) Professor Elstein's plan.

It will be well to take the plan advocated by Mr. Banting first, because it must be criticised rather than recommended. He cuts out carbohydrates; no sugar, no starch. So far so good. But on the other hand he allowed unrestricted indulgence in albuminoids, as the flesh of animals for instance. This reduction of a human being to the status of a carnivorous animal has been attended with unfortunate results. All along in this treatise the danger of an excess of albuminoids in the food has been steadily insisted upon. The dangers involved therein have been pointed out, especially in the chapters on "Bright's Disease" and on "Gout." And this danger of kidney injury is no myth or imaginary risk. Not long ago a very illustrative case came under my notice. A man of sixty, whose kidneys were probably not above suspicion, was advised to reduce his corpulence, he then being in good health and feeling well. It was well to take off a couple of stones, his medical man said, and Mr. Banting's plan of reduction was recommended. The desired reduction was accomplished but at what cost? No suspicion of any impairment of the kidneys seems to have suggested itself to any one, to put them on their guard; and the untoward result was, a condition of lithiasis was established, and in a few months Bright's disease had claimed its victim. Whenever any reduction in a corpulent person over middle age is contemplated, the possible condition of the kidneys should never be forgotten. There is much reason to believe that many persons have fallen victims to their desire to reduce their corpulence, and the adoption of Bantingism for that end. It is, really, a very dangerous practice.

Professor Elstein's plan is free from the dangers which lurk around Mr. Banting's method. The Göttingen Professor, who has given much attention to, and has written a small book on, the subject, sums up as follows: "The permission to enjoy excellent sweetmeats, always of course in moderation, as for instance salmon, paté de foie gras, and such like delicacies, to our corpulent gourmet to his other sacrifices. These consist in the exclusion of the carbohydrates. Sugars, sweets of all kinds, and food in every form, I forbid unconditionally. The quantity of bread is limited at most to 5 to 6½ ounces a day, and of vegetables to 1½, asparagus, spinach, the various kinds of cabbage, the cucumbers, whose various conveyers of albumen, as Asparagum, are to be avoided. Of meat I extend none, and therefore, the Beefsteak is

to be avoided, but on the contrary, sought after. I permit bacon-fat, fat roast pork and mutton, kidney fat, and, when no other fat is at hand, I recommend marrow to be added to the soups. I allow the sauces as well as the vegetables to be made juicy, as did Hippocrates, only for his sesam oil I substitute butter." This is a summary of Ebstein's creed, which is a distinct improvement upon the regimen advocated by Mr. Banting. Some modifications, however, may be suggested, such as further development of fish in the dietary. Fish of all kinds, fresh, dried, and canned, should be largely taken. Then a caution about the amount of lean meat and legumes is desirable, as being albuminoids and needed in small quantity. Salads of various kinds are good. The lettuce, the endive and celery may be added with advantage to his limited list of vegetables. The onion, kale, sea-kale, turnip-tops, do not contain so much saccharine matter as to exclude them. As to the potato, of course an objection lies against it theoretically, but when the stomach has to be filled a small quantity of potato will help to do this in a satisfactory manner. The palate must be consulted, and so must the stomach; and a small quantity of forbidden matter may in many cases be desirable. Only of course more time for the desired reduction is then essential.

Bread, biscuits, crackers, corn cakes, milk puddings, other puddings, all must ruthlessly go if the reduction has to be successful.

Malt liquors of all kind must be foresworn, and a light wine or spirits and water alone be indulged in.

As much exercise as possible should also be taken, as exertion counts off so much food.

But do not let the new regimen to be too severe, or so rigid as to affect the general health. If ill effects are being experienced, some relaxation for a while may be advisable. Where obesity is the outcome of indulgence in rich food in unlimited quantities, combined with generous drinks and habitual laziness, the regimen requisite for its reduction will be beneficial generally. But where a person is merely corpulent and yet well with it, it is a matter for deliberate thought as to how far any sudden change of regimen is desirable. Some families are normally stout just as others are lean. And as it is impossible to fatten the latter, so it may not be wise to try to reduce the former.

The treatment of obesity is the avoidance of carbo-hydrates; while in emaciation carbo-hydrates, in their most digestible forms are what is required to give a certain roundness or embonpoint.

There is, however, a certain atrophy seen in females at or

about puberty, and then again at the menopause, or the change of life, which is linked with anæmia. Girls or women, therefore, whose pallid, the blood goes thin, and fat accumulates upon them, especially the subcutaneous fat. With girls a quantity of red meat is also essential for cure as are chalybeates. With women at the change of life, an excess of albuminoids may tell upon the kidneys. The fatness may depend and does depend evidently in some cases, upon the deficiency of red blood-corpuscles, leading to impaired oxidation, and from that again to the deposition (as fat) of the unconsumed carbon hydrates. This impoverishment of the blood may be due to the pressure of some poison in the blood breaking down the red blood-corpuscles, or hindering their formation, and in such cases the removal of the toxic element is the first step to be taken; after which, the cause being removed, all goes well. Such a case came under my notice some time ago. Cont here was the underlying causal factor; and the clearing away of the toxic agent by uric acid solvents was in one case followed, and quickly too, by the re-formation of red blood-corpuscles, and a reduction in the adipose tissue. But the reduction of the fatty tissue in ladies at the change of life—especially if some malnutrition of the muscular wall of the heart be also present—is not a matter to be lightly undertaken, and requires considerable judgment.

When a person is naturally stout and leads a rational life, taking exercise and eating in moderation, any great reduction in bulk is a questionable advantage. But where obesity is the outcome of gluttony and laziness, then a change all round is certainly desirable. But even then a sudden reduction in bulk is not without certain drawbacks, and entails an exercise of the judicial powers.

When a reduction in bulk has been achieved, consistency in the dietary is necessary in order to prevent increase in weight; and a regulated dietary is, in any case, a pleasant matter to say the least of it, and must become the rule of life.

CHAPTER XXXI.

THE FOOD IN INDIGESTION.

PROBABLY none of the topics in the connection of food with illness is more interesting than the one now to be considered. It will be necessary to go into some detail in the matter, as "indigestion" covers a large field.

The first matter in this consideration is the physiology of digestion, which has been spoken of before, but must be glanced at again. First so much farinaceous matter is rendered soluble by the action of the saliva. The saccharine elements of our food and the soluble outcomes of starch pass readily through the wall of the alimentary canal; and so the stomach is relieved of their presence. What starch remains untouched by the saliva passes through the stomach (unaffected by gastric action), and finds in the intestines the pancreatic solvent. In the stomach the albuminoids of our food alone are acted upon. That is the second section of the digestive act. The third and last is the pancreatic digestion, where starch, albuminoids and fat, all are dealt with.

Gastric indigestion then may be due to

1. Too much undigested starch in the stomach; or
2. Albuminoids too hard to be dissolved; or
3. Fat floating about in the stomach, offending it and interfering with its proper work.

In these cases the stomach may be equal to its duties if the dietary be adjusted properly.

Then there is indigestion based on the stomach having more food in it, *i.e.*, a larger bulk of contents than it can deal with. Here smaller quantities of food must be taken at once.

Finally the stomach itself may be at fault and be unequal to its duties. This will take two directions.

1. Deficiency of gastric juice;
2. Defective muscular movement.

The late Dr. Arthur Leared held that pain is the proof of the first; and a sense of distension (with flatulence) the evidence of the other. In many cases the two are combined.

These last causes of indigestion may be due to organic disease, but by no means necessarily so. Functional incapacity of the stomach may, and often does, exist without organic disease. It may or may not be followed by structural or gross alteration.

When the stomach is at fault, there is no need of any measure to be taken for, which cannot be considered here. Frequently, however, artificial pepsin in addition to the amount furnished by the stomach will enable the labouring vessels to get over its difficulties. Often the addition of carminatives and condiments to the food will relieve the difficulty. But granting all this, and not attempting to underestimate its importance, the main matter with primary or organic indigestion is the food taken in to the stomach.

In the first place it must be of such character that it can be properly assimilated by the teeth, and be so reduced by mastication as to be in a fit state for the stomach to act upon it. If such dental preparation has not gone on, then the food reaches the stomach in an unfit condition, and consequently that vessels is incapable of dealing with it. Defective teeth, the habit of forcing the food unchewed, and unsuitable character of the viands, are the main causes of this form of indigestion. Certain things, as pastry, cake, fat and flour are kneaded together into a mass not readily disintegrated; cheese, nuts, bits of hard meat or gristly hard pieces of potato, cabbage or other vegetable matter all vex the stomach.

In the case of Alexis St. Martin, Dr. Beaumont found that such articles offended the stomach, arresting the flow of the gastric juice, and causing the living membrane of the stomach to be dry and raw. From this we can understand the notorious old fact of the above-mentioned articles provoking indigestion in a delicate, sensitive, or feeble stomach. Persons who can eat most articles of food with comfort, and these indigestible articles do cause them suffering. Consequently they should be scrupulously avoided.

Where mastication is defective owing to the loss of teeth or the decayed condition of the old of the dentition, it be broken. When the teeth have become out right and are in working order the food can no longer be clogged by lumps of food upon which it cannot come to exert its action.

When the habit of eating quickly obtains, and the food is not chewed or masticated, and thus the stomach is offered an unfit condition, the method of eating,* the manner of eating, must be corrected.

Matters disturbed with regard to the stomach, and the consequent food indigestion. A badly fed stomach has no chance of doing its duty.

respect. If the cook do not have her fat properly hot, it soaks through the fish, and the muscular fibres so becoming coated with oil are beyond the reach of the gastric juice. Fatty meat, as the flesh of the goose and duck, are avoided by dyspeptics. Such fat-swathed muscular matters are as burdensome to the stomach as quantities of undigested starch.

These are gross and tangible matters, obvious even to the uneducated eye: matters which merely require to be pointed out in order to be comprehended. But beyond these there are subtler matters, such as the difference betwixt veal and chicken, pork and rabbit.

Meat hard of fibre is indigestible, as pork and veal; then beef; then mutton and lamb. Even parts of the same animal vary. A piece of hard neck of mutton is much less digestible than the loose fibre of the meat on the sheep's head. A veal cutlet contrasts with a sweetbread. A salmon steak contrasts with a whiting; though fish are as a rule more digestible than the flesh of mammals. The flesh of fowls is widely different, as a succulent duck compared to a partridge. Game is very digestible with its short fibre: while goose is not only rich and greasy but its fibres are longer and coarser than is the case with game. A leg of a chicken is less digestible than the breast.

Articles that are flatulent cause indigestion, and consequently many persons of feeble digestion have to eschew the legumes as the bean and the pea. Peas bear an evil reputation: and especially in the dry form. Vegetables as a class are regarded as "windy." Uncooked vegetables are to be avoided by dyspeptics not only on this account, but because uncooked vegetable matter does not readily undergo disintegration.

The presence of starch granules in considerable amount in the stomach is a fertile source of gastric suffering. This is an aspect of the subject far too little considered; and yet it is a matter of great importance. The gastric juice has no action upon starch granules. This is quite a different matter from mere masses of farinaceous matter like pastry or pie crust. The disintegration may be perfect, or fairly so, still the starch granule is as indigestible in the stomach as so many minute particles of glass. Commonly, however, there is a combination of causes in action, as more perfect disintegration by the teeth would necessarily bring the starch granule into contact with the saliva. But putting this matter aside for the moment, and considering the farinaceous matter simply as an aggregation of

starch granules; these last, if sufficiently numerous, may even obstruct the action of the stomach. They absorb fluid, and add to the burden borne by the stomach as to its mass and movement. They incommode it; and therefore are obstructive and injurious.

In order to get rid of any superabundance of starch granules, the dyspeptic ought to take the farinaceous articles of food first, and adopt the rule of the nursery, *viz.*, porridge before meat. If a milk pudding be taken first, then the saliva can act upon the starch in the mouth, in the gullet, and in the stomach before the food is acid. By so doing the starch would be rendered soluble and so escape through the wall of the stomach, and would get out of the way, leaving the stomach free to deal with the meat which comes after it. That is the common-sense view of the subject; and a following of the natural order of digestion. But custom bars the adoption of such measures. The dyspeptic must rise above custom. That is all that can be said.

Then there comes the question of dealing with the starch granules previous to their being taken into the mouth. As has been pointed out before, exposure to heat not only cracks the starch granule, but renders much of it soluble as dextrine. Where the application of heat is made by the maling process, there is also the action of the diastase upon starch. In cases, then, where the presence of raw starch granules offends the stomach, cereal matters which have undergone a preparatory digestive course must be chosen. There are many in the market; before long there will be many more. In the meantime the cook can put her farinaceous matter in the oven by itself for some time before proceeding to heat it in the ordinary manner for a milk pudding. Or starchy matters which have undergone some preliminary digestion can be used for soups or broths. Of food, bread well boiled and put through a sieve was used by me as "bread *jeu*," or added to milk or soups. By so doing, not only are these matters as the first course at dinner, a sufficient supply of grape-sugar would be furnished to the body for combustion and support; while the stomach would be left unembarrassed to go on with its own work.

It is not very wise to point out how much food is lost by the use of this well-known rule; but then it is unavoidable in the system of taking up to working point. The dyspeptic is not aorta to be helped in many ways to be sincerely pated.

Where the stomach is offended by too much oil, or too much poultry fat or mutton, or by the fat of the calf, or by the fat of the

lops," such source of offence must be avoided. Very often, if the fat be hidden from sight, the objection to it is lessened. Where fat is not well tolerated it is prudent to conceal its presence. The stomach if left unprejudiced may be more lenient as to its presence. It may be necessary to eschew the fat of animals and compensate the want of this by butter. Butter mixed with potatoes, butter hidden in milk puddings, butter melted with fish: these are comparatively unobjectionable. Where no such compromise is possible, then fat must not be taken till the work of the stomach is over. If the meal be of digestible character, in from one and a half to two hours the fat may be taken as cream (with maraschino or other liqueur), as cod-liver oil, or as an oil emulsion.

What each individual can do, or must do in order to come to terms with a fctchy stomach, is a matter of experience in each case; and only to be ascertained by trial, and indeed repeated trials. Nor is the stomach always the same as to tolerance. What can be endured at one time is intolerable at another. Whenever the system has been taxed or upset the stomach becomes intolerant. Thus when the system is wearied the stomach is bereft of much of its capacity, and food ordinarily well-borne disagrees; and what is more, the gastric revolt tends to become chronic. An American lady-patient, with dyspepsia, traced her woes back definitely to a certain dinner at Genoa. After a long day's journey she and her fellow travellers arrived at the hotel just as the table d'hôte was being served. They sat down and made a hearty meal. This outrage on a weakened stomach in her case led to persisting indigestion. She had then suffered for seven months; and by the greatest consideration had attempted to atone for the original outrage. Offence and repentance are necessary with an injured stomach—which may be as unforgiving as a Corsican.

Then again any mental perturbation may make itself felt either as by loss of appetite or by indigestion. Bad news, received during or shortly after a meal, substitutes for the usual comfortable sensations that of a stone in the stomach—as the food lies motionless in a palsied viscus. There is no doubt that the digestive powers of the Anglo-Saxon are waning. Our early ancestors were equal to a feast after a hard day's marching or fighting; but not so their nineteenth century descendants, who when their energies are taxed find the stomach impaired in digestive capacity. Dr. Payy says that this fact is recognised by experienced sportsmen.

Then again there are stomachs which resent the presence of

fluids, and, contrary to what is usual, be a liquid of its own accord. In most cases of dyspepsia, and on some occasions of men-struations or milk with some carbo-hydrates, such as the stomach. But in these exceptional cases no other course is possible has to be avoided. If this be rigidly done the digestion goes on normally and without discomfort.

Having thus reviewed the causes of indigestion, and its varieties, it may be well to consider its quantity or extent. In some cases the defect is so slight that if the more indigestible foods be avoided all is well. Pastry, uncooked vegetables, nuts, are to be eschewed; but nothing more under ordinary circumstances. Others again cannot take pork, veal, or cheese without suffering for their indiscretion. Then come those who may eat cold beef, or a beef-steak, but who can manage nothing. For all these, artificial pepsin may be enough to get them over the worst difficulties. Persons who are no worse than this have not much to concern of; they may be put to bed, and here now and then especially when travelling.

Then come those more gravely afflicted, and who really have to be very careful about their dietary. They must pass the vegetables as a rule, take very little animal food at once, and that only of the most digestible character. Fish, especially white fish, may be cold or baked rather than fried; chicken, rabbit, or a sweetbread are the chief articles of food with them. They are restricted to milk-puddings; and other puddings have to be foregone.

With such persons matters are growing serious. They can get on fairly well at home with a cook trained to meet the necessities; but out-of-door dining out is next to impossible, which for many is burdensome to them. It is well for them never to forget to keep some pepsin near at hand when eating a man away from home.

The indigestion of miles rarely goes beyond this, except in the case of consumptives, or persons with some affection of the stomach, or some acute disturbance of a naturally feeble stomach. But with females a further limit is not uncommon, and that is the condition of a sufficiency of food to sustain life becomes necessary. Only liquid and semi-liquid food can be eaten, and that in small quantities at once. Some persons take meat-infusions containing some predigested meat, or take meat without some malted preparation, a boiled potato, or a slice of some stewed fruit: such form the dietary. It is a good plan to "string the changes" when the doctor is so far from hand.

possibly, the appetite is small and the palate dainty. The body-income is necessarily small, and cannot be raised. Then let the body-expenditure be reduced to the minimum. The patient must go to bed early, surrounded by bed-clothes in a warm room, the need for fuel-food is brought down to the lowest point; while perfect quiet reduces the demand for food to supply force. Rigid economy is necessary with small means; and this has a wider application than is commonly attached to it. It applies to the body physiologically as well as to social wants financially. A small income is incompatible with lavish expenditure. Beggary is the alternative. The same with the body: if the outgoings exceed the incomings, then exhaustion of the body-capital is inevitable.

The smaller the bulk of the food which can be taken at once the shorter the interval must be betwixt meals. Even when the wonted fare can be taken, but only in limited quantities at once, a small amount of food betwixt meals is usually a good practice, especially in cold weather. When the patient is reduced to the condition just described, the intervals betwixt meals must be made very short. Passing it should not exceed two hours in some cases. The fast of night is injurious. There is a well-recognised tendency to a lowering of the body-energies in the small hours of the morning. Some feed about 2 A.M. as most desirable. With many ladies of small digestive capacities such nocturnal meals add materially to their comfort and enjoyment of life. Such persons of course are readily misinterpreted as dyspeptics, and this is neither a real nor one of those imaginary ailments whose symptoms are so mad as to possess into their minds a potent for nostrum, and Asylum with such good effects that they need no time save enough to be it longer. Uncontrolled, they can go to some such extent as to require the Asylum nostrum once more. And there are persons whose stomachs resemble these crazy animals. Dyspepsia is a term produced, pronounced, specified, requiring a strict regimen for a time. On this distinct improvement is made, and the worst state is passed, and before long some upset occurs that is the mark of a new stage of the disease. And so on is the history of many a life of despair. And the lesson it teaches is the necessity of a constant watchfulness, and that extreme watchfulness, with the least suggestion of change, is the only protection against the danger of breaking down.

Probably this class of cases finds as many patients than any other, as persons are of a delicate build, if not possessed of delicate organs, and the delicate structure of the delicate body.

income could be madeable to them. They are not intelligent enough to get their food good, and they are not careful, their poverty often preventing them from trying to get something well done. They are not in a position to disagree with them, but when they do not they are doing them wrong.

Of this class of person, Dr. P. says, "In the chronic impairment of power exists, as before, the patient must be gradually weaned from his habit of eating. Whilst avoiding that which is known to offend the stomach, and whatever, though innocuous, may be known to stand to upset the stomach, a maximum of nourishment is to keep the diet as closely to what is natural as the condition of the case will permit. Frequently, a person who is suffering from dyspepsia, he is recommended to avoid rich and stimulating food, and may perhaps, in the course of time be brought to eat exclusively, or almost exclusively, liquid nourishment. Such a diet is sufficient to lower the energy and to deprive the stomach of its work. The organ, getting no employment, becomes weaker and weaker, and is also prejudicially influenced by the impoverished state of the system. The aim of the physician in these cases should be rather to raise by appropriate means the digestive capacity to the level of digestible light but ordinary food, than to reduce the food to an adjustment with a low standard of digestive power."

This aspect of the matter is however, not. Instead of cutting off one by one the different articles of food, and by so doing every way to be able to feed the unfortunate dyspeptic on some article of food which can be taken safely. With a view to getting to the stomach the food requisite and indigestible, and to get it in its best possible shape, the aim is to raise the digestive capacity by appropriate medicinal measures as well as suitable food. When the system at large is badly nourished the stomach shares in the general debilitation. Predigested food may be given to raise the general tone, not to relieve the stomach of its work. The ready resort to peptonised foods may render it impossible to test the capacity of the stomach. On the other hand, the physician may throw the case back and let the labour be thrown on the patient, and start be necessitated. Consequently, the treatment of dyspepsia is not a matter of off-hand decisions, but of the most careful and cool, case with its individual peculiarities, and the physician who dyspeptic leaves off one article of food, from the list of those which

loathes the few remaining articles of diet, and the starving creature often fasts sooner than eat them. The body must be fed until the digestive capacity is regained, if possible. That, by a judicious dietary and proper medicinal measures the digestive organs can be restored to a fairly capable condition, is a matter of fact. But in bad or confirmed cases the loyal co-operation of the patient must back up the wise counsel of a competent physician; any defect in either being fatal to success. When dyspepsia is linked with chronic constipation—a common occurrence—this matter must be pointed out to the patient clearly and forcibly. Such patients rarely possess the perseverance indispensable to success. If the patient does persevere courageously the reward comes. We get what we deserve in life to a very great extent, and the maxim certainly holds good of persons of weak digestive power.

There are two matters dyspeptics must ever attend to. (1), never to touch meat that has been twice cooked, though this holds less against fish than other meat; and (2) fat which is rancid, or which has been exposed too long to heat, or to too high a temperature. One American patient could eat cold butter in any quantity, but a fried whiting was an outrage to his stomach.

When a dyspeptic is reckless (as was in all probability the case with the late Tom Carlyle, the philosopher), he endures a great deal of avoidable suffering. But many dyspeptics are of this class, and especially intellectual dyspeptics. They perpetually commit outrages on their digestive organs, and suffer for them, but all the same they hold on their own way. Experience does not teach them. They are very unsatisfactory patients.

Others again commit these recurrent outrages not from wilfulness but from forgetfulness. While the subject is being discussed in the physician's study they are alive to all that is said, and endorse it all. But when their minds are occupied by other matters all this is forgotten, and the indiscretion committed; and it is only when the unpleasant consequences put in their appearance that they become conscious of their forgetfulness.

The most unfortunate matter is that one such act of forgetfulness may undo a long labor of careful dietary, and throw the patient back to climb the painful ascent. It is the stone of Sisyphus once more.

But one curious well-recognised fact there is in connection with indigestion, viz., when the craving for a certain article of diet is very strong, even if it be of a most objectionable nature as regards

of dyspepsia generally, it commonly happens that the patient does not know how and when it is safe to permit the gratification of his appetite, a matter yet hidden from us.

Still, all the same, modern chemico-physics has done so much to enable us to help the dyspeptic, that too many persons, even when the patient is resolute about doing nothing, are not helped.

Gouty Dyspepsia.—Whatever the difficulties to be overcome with the ordinary dyspeptic, they are not less those of the gouty dyspeptic. The reader will remember what is said in the chapter dealing with gout, as to hepatic powers of formation of uric acid. Whether increased or not, the gastric secretion, or not, is a matter shrouded in darkness. Certainly it is that the gouty dyspeptic is troubled with acidity. Uric solvents are distinctly indicated, and without them no diet, however carefully carried out, is but partially successful. The diet, too, in these dyspepsias is confined to leaning to common sense, and to keep in the stomach from any fat that has had too much of its life. Consequently all fried meats are to be avoided, even when they are lean. Their fish should be boiled, or baked in milk, and the sauce should be served up as fish pie, which must not be cooking for the patient.

Then again they find sweets too many, particularly chocolate, and especially the cane sugar added by the cook to fruit. As a rule a gouty person is not a "sweet tooth," and sweets are seldom liked by them. Still many would like to have stewed fruit, if this could be done with impunity. As much bicarbonate of potassium as will keep a stilling, or a quarter-bit to each pound of fruit, when put into the oven to stew, will neutralise the acidity, after being out of the stomach, the sweetness of the levulose sugar in the fruit will be as good as in the free field. Such stewed fruit will not do any harm, and is useful, but it provides an article so good, and very palatable, in a defensible form. By using canned fruit, stewed fruit, or fruit sauce, or stewed fruit can be had all the year round. On a large scale it can be eaten, or stewed.

Milk puddings made without eggs should form a large part of the ordinary, and these should not be sweetened with cane sugar, or golden sugar, as milk sugar or dextrose. A very small quantity of cream perhaps gives the best result. If the cream is to be used, it should be previously been exposed to heat, as sterilised milk, and should be unflavoured rice, or broken biscuit, or rice or biscuit crumbs, and a moderate quantity of ground or crushed malt, and a very small quantity of yeast. This should be poured milk hot enough to cook the rice, and

sipped. The dish should be placed where it will not get cooler (and not much hotter) for ten minutes, after which it can be put into the oven long enough to become slightly. Such pudding does not require to be kept in the oven till it is "enough"—as is the case with puddings made with farinaceous matters which have had no previous acquaintance with heat—for the materials have already been cooked. Such puddings will not turn sour with most gouty dyspeptics, though the opposite may be the case with a small proportion.

Then a staple article of the dietary will be soups of all kinds, thickened with broken biscuit, malted rice; or cooked farinaceous as the A. B. C. cereals, the farina cocta, shredded maize, baked hominy, etc. If sago or tapioca be used, either should be placed in a dish in the oven for an hour or two hours, care being taken that the farinaceous material is not scorched. Or the sago, rice, or tapioca should be well boiled ere adding them to the soup. By so doing the risk of filling the stomach with quantities of insoluble starch would be avoided; a matter which has been discussed in the early part of the chapter.

Farinaceous matters—supplying the essentially necessary grape sugar to the body—must form a very large portion of the dietary with gouty dyspeptics, and must be so treated as to give as little trouble to the digestive organs as possible. The more such farinaceous matter has been exposed to heat the better, and the less digestion they require within the body.

Soups made with cooked farina, boiled and baked fish, milk puddings prepared as directed, baked apples, and fruit stewed with potash, grapes, bananas, and such natural fruit as agrees, carefully masticated so as not to offend the stomach by tangible lumps, must form the staple dietary of the gouty dyspeptic when the stomach is very irritable. When a careful curative course has led to a better state of affairs then some chicken, rabbit, tripe, sweetbread and like digestible matters may be eaten. Well-mashed potatoes with cream, or butter and milk warmed together, put down before the fire to brown (better than being put into the oven), in order that heat and moisture may still further modify the starch granule, will form a desirable change.

Cold fat is less apt to form fatty acids than warm fat; and biscuits—the plainer the better—and butter should form a considerable portion of each meal. All sweet biscuits contain cane sugar, and so are apt to turn acid. Those which are made with malt are less objectionable. Cream with stewed fruit is also a suitable fat.

As to beverages, a little claret or wine of like character, or some alkaline water, (better if effervescent), or perhaps alcohol, may be desirable. Or a little pure spirit and much water may be given. As to tea, coffee, and cocoa, there seems little objection to them, unless one or the other be found actually to disagree. The sugar added to sweeten them is the cause of much of the acid erroneously attributed to the beverage itself, it seems to me. Milk with equal parts of some mineral water, as seltzer, if milk-alkali be needed, or Vichy, or Vals, if more alkali be desirable, forms a desirable beverage.

Like all dyspeptics the gouty dyspeptic must give his mind to the treatment of his own case. If the dietary is restricted and the regimen irksome, the sufferer must remember that he is the person benefited; and that it is only by the observance of strict dietetic rules that relief is to be obtained. The power to help is limited to the patient's determination to be helped—a matter the dyspeptic must not forget or overlook.

CHAPTER XXXII.

THE FOOD IN BILIOUSNESS.

PROBABLY the contents of this chapter will excite keener criticism than any other in this book. The view taken traverses the ordinary opinions on the subject; and yet an increasing experience bears them out. Some half score years ago a consideration of the chemical composition of the bile acids led me to reflect upon their derivation; and from that again to raise a practical issue, viz., the proper food for bilious subjects.

The bile acids are two. The one Glycocholic acid with a formula $C_{26}H_{43}NO_6$; the other Taurocholic acid, with a formula $C_{27}H_{45}NSO_7$. Both contain nitrogen; and one, also, some sulphur. And these two constituents point to their origin in the albuminoid elements of our food. Then comes the further consideration that persons who are bilious in their early years are apt to become gouty in middle age and advanced life. The whole subject was one of deep interest. Instead of bile being derived from starchy, sugary, and fatty foods, reputed to be "bilious," it evidently had other associations. The dietary of lean meat, dry bread and tea, without sugar or milk, in favour with the bilious, seemed erroneous in theory. But how about clinical facts? The proof of the pudding is the eating, after all. One confirmed bilious subject, of the bilious diathesis, and wearing a look of persistent suffering, came under my care and seemed a most suitable subject for experimentation. With some difficulty she was induced to give up the meat, bread and tea, and take to farinaceous matters and milk puddings, and much to her surprise soon felt very much better for the change. Her case has been alluded to before in the chapter on Food in Affections of the Nervous System, and since writing that chapter an opportunity for going over the facts (as there stated) has offered itself and confirms them. The clinical facts fitted in with the theory so completely that further trials were made. The results of these further trials have pointed unmistakably in the direction that much albuminoid food is not desirable for bilious persons, despite the fact that they find such food sits well upon the stomach. Such food

may be agreeable to the stomach, but it certainly may not be so to the liver. The analogy used in "Indigestion and Biliousness" is as follows: "A young man may find favour in a girl's eyes, but her parents may not be satisfied with him. Acceptable enough to her, the very qualities which weigh favourably with her may be those on which her father specially founds his objections. That food which is acceptable to the stomach may be the very matter which disturbs the liver." Five years' further experience bears out what was written in 1881, when the book appeared.

The function of the bile is, so far as we know it, the emulsifying of fat. If fat be abjured there can exist but little necessity for bile. If a disproportionate amount of albuminoids be eaten, the material for excessive bile is provided. There is a disturbance of balance in the economy. If fat cannot be taken the excess of albuminoids can be avoided at least. And the practical result is relief to the bilious symptoms. In the present state of our knowledge it is impossible to be more explicit. Some day it may be feasible to give a more satisfactory and complete explanation. Our acquaintance with the functions and operations of the liver is recent, and incomplete; but I venture to think the rational outcomes of the above consideration will warrant some practical conclusions.

The food of the native inhabitants of warm countries as taken from Dr. Pavy is as follows: The Mexicans are "accustomed to uniform nourishment of an almost entirely vegetable nature;" while the inhabitants of New Caledonia "subsist chiefly on roots and fish and the bark of a tree;" while in Hindostan: "From the earliest period the most general food in India has been rice, which is still the most common food of nearly all the hottest countries of Asia. Pulse, roots and fruit are also largely eaten." (Ghee is a great delicacy.) In Ceylon they eat "rice seasoned with salt and a few vegetables." In Egypt "vegetables form the principal food of the lower orders." In the Sahara desert "all live on dates." Butter and palm oil are also in demand in various parts of Africa.

The Spartan has his bread and oil and onions. The Italian, maize and macaroni. The Turkish hamas is a vegetable man, but his dietary is almost entirely vegetarian.

"The attention of Herodotus was called to the fact that the Persians, the martiallest and most sporting nation of the east, women had at meals not only several dishes, but several courses of vegetable food, proceeding a very moderate allowance of solid meat." (Chambers.)

Set against these facts the other fact that when the Anglo-Saxon persists in his home-dietary on the plains of Bengal he becomes the subject of biliary trouble, and not infrequently of actual disease in the liver. Whereas men who adopt the dietary of the natives suffer comparatively little.

Indeed the relations of hepatic trouble to an English dietary in tropical countries is a notorious fact.

Regard being paid to what has just been said, the great object of the bilious, like the gouty, is the avoidance of albuminoid articles of food, or the taking of them in very moderate quantities. How many persons, who, when dining out, indulge themselves in meat and drink, find that a calomel pill and a scidlitz powder enable them to escape sundry unpleasant after-consequences? That fact throws some light upon the relations of the food taken to the function of the liver. Loss of appetite follows any excess in food; and what does that tell us? When the liver has more material brought to it by the portal vein than it can deal with, loss of appetite is the result; and by a period of abstinence the liver works round, and all goes on again as usual. The stomach can reject its contents. The bowels can sweep out objectionable matters. The liver can do neither. If not protected by anorexia the liver would have rather a hard time of it with persons who think that the regulation four square meals a day are a part of the Divine Scheme. Loss of appetite comes to its rescue. The bilious disturbance caused by a surfeit, especially with children, is another contribution to the subject. The liver is the furnace in which the waste and superfluous albuminoids are burnt, we are told by physiologists. When there is an accumulation, pyrexia often comes and helps on the oxidation, or in other words "Nature holds a bonfire." By such means the not readily oxidisable nitrogenised matters are burnt up.

Put briefly, with bilious persons it is well to reduce the amount of food taken to the body-needs, or as near this as can be attained. And especially is the moderation to be practised as regards those nitrogenised elements of our food from which, obviously, the bile acids are derived.

Bilious persons, then, should avoid the solid joint at meals. Joints, chops, and steaks, indeed lean meat in every form, must be taken in very small quantities, if at all. Fish, as a less concentrated form of albuminoid material, is to be preferred to flesh. Vegetables should be largely eaten. Fruits do not disagree. Eggs and milk have an evil reputation for encouraging biliousness. Fatty

and sugary matters are reputed to be bilious; but if this be so (and the view has a basis of fact) it must be indirectly and not directly. If the readily oxidisable hydro-carbons attract the oxygen of the body, the less oxidisable albuminoids will be left over, in the ashes or cinders of the body-combustion.

Farinaceous matters and vegetables should form the staple dietary, accompanied by a certain amount of fat: the amount varying with the season, climate, and locality. In temperate climates a certain amount of the concentrated fuel food, the hydro-carbons, may be required. In tropical countries the dietary consists of carbohydrates only, with advantage. The gastronomic proclivities may, however, go in another direction: but reason must be brought to bear upon them.

This view of the dietary appropriate to bilious persons will strike many readers as being novel, and so antagonistic to prevailing opinions as to savour of unsoundness. The testimony furnished by my limited experience may seem insufficient to warrant its being put forward so confidently. If the experience has been limited and recent it has at least been convincing—so far as the writer is concerned. It is one of those matters which can be reasoned upon rather than demonstrated. And if the reader remains incredulous there seems to the writer no means of convincing him. If he himself be plagued with biliousness let him give the matter a fair trial, and note the result. There are many matters which, if they had had to wait till they could be proved to a demonstration before they were launched into the world, would have remained unborn. On the other hand a sweeping change of opinion and practice must have some solid foundation before its propounder can expect the world to accept it. The arguments given in this chapter seem deserving of attention, and the verdict of the world can be awaited with some confidence. My experience has gone in the direction that what is sound in theory works well in practice.

The bilious person should avoid the eggs in the traditional breakfast—bacon and eggs—taking the fat of the bacon by preference. Then if there be any fish some of it should be eaten. In summer some lettuce or salad may be added; in winter some fruit, especially the apple. There exists no real objection to some cream and sugar being added to the tea or coffee. Then the *main* meal should consist of some soup or fish with well-mashed potatoes or other vegetable, and “pulled bread” or biscuits. Dinner should consist of some fish or chicken, a vegetable course, milk pudding, and bis-

omit and butter. Such was the dietary on which a gentleman whose liver gave him much trouble got well, after being under several physicians of repute without avail. On his usual dietary, which embraced a considerable quantity of lean meat, he grew worse and lost flesh. This is a common story. When the liver cannot carry on the proper metabolism of proteids into the serum-albumen of the *liquor sanguinis*, but breaks them down into bile acids or urine solids, the tissues go unfed. In fact the albuminoids consumed "go to feed the disease and not the patient," to use an old-fashioned and time-honoured expression now getting lost sight of too much. Of course it is well to raise the tone of the liver by resort to hepatic stimulants at the same time. But by reducing the demand upon the liver as regards its function of dealing with albuminoids it can perform its work, and normal metabolism goes on. The consequence is the body is once more fed and the biliousness relieved, *i.e.*, the proteid matter is going to its wonted destination—the tissues, and less bile acids are formed. Recently a well-known writer of fiction came under my care with a tongue covered with a thick fur deeply stained with bile, and lithates in her urine. She too had been losing flesh and getting worse on a meat dietary with some alcohol—the proper dietetic means to get strong in popular belief. She adopted my suggestions with quick and prompt relief, and got well most satisfactorily. When the liver is out of order the ordinary dietetic means for getting strong are rather mischievous than useful. It is not what is swallowed but what is digested that is the question. Primary digestion goes on in the alimentary canal, and secondary digestion or elaboration is performed by the liver. Food may be taken and digested, but if the elaborating action is defective nutrition fails, just as much as if food were withheld in the first place. When the body is starving from liver inadequacy, to force down liberal quantities of generous food is to overtax the liver and hamper its action still further. Food rich in carbo-hydrates, and containing but a small proportion of albuminoids, is that which is really indicated and required. The appetite may be feeble and capricious, the palate may be dainty and have its preferences; these may be met, but after all the function of the liver must be remembered; and with it the true body-needs.

Grape sugar, the fuel food of the body, must be supplied in sufficient quantities. Albuminoids for tissue repair, too, are required; but the amount actually necessary for that end is but small. All beyond the bare necessities are superfluous and only hamper and

embarrass the liver; and increase the amount of biliousness, or other solids, as the case may be. That is the lesson taught us by modern advances in physiology and clinical medicine.

This is all rank heresy in the opinion of the people at large. When a person is below par, plenty of animal food and good wine, tonics with iron, are the things to get him up: certainly in most cases; but when the liver is involved there is an error in the calculation. So long as the liver is competent to its work, so long the equation is correct; but when the liver is unequal to its work this regimen but further embarrasses it. The number of stories told of the failure of this plan to get them strong, by delicate patients, is conclusive. A number, too, furnish some curious corroborative testimony. "They bring a specimen of urine for examination in a bottle; and not infrequently that bottle has on it in raised letters: 'Quinine and Iron Wine.'" It has signally failed to do its office. A look at the tongue and a few well-directed questions soon lighten the matter up; and on a different regimen, dietetic and medicinal, improvement sets in. But these persons are often hard to convince. At first sight it does seem an odd way to increase the strength, viz., cutting down the food, and especially the animal food. But the alteration is soon felt. "She stoops to conquer." And this manoeuvre can be practised with satisfactory results in hepatic inadequacy. No chain is stronger than its weakest link. The appetite may be fair, the stomach capable; but the liver is weak. The nutrition is defective usually when the liver is incapable, and the appetite fails. That is Nature's way of attaining a new balance by "leveling down." But an appetite is held to be a man's inalienable right; and so he tries artificial means to create one. He defeats his own ends. He is thwarting and traversing nature's provisions, and so the plan fails.

When the liver is incapable it is well to reduce the demands upon it to its capacities.

So much for the dietary in biliousness.

Now about the drink. A naturally feeble liver cannot get on with malt liquors. Whether it is cholamin or atonic in its character is pernicious; especially with persons of the "And" type, i.e., highly developed nervous system and small viscera. Malt liquors prepared by the English system of brewing are especially injurious to an incapable or sensitive liver. The beers brewed on the German or Lager system are less objectionable, and many "feeble" persons can drink beer on the Continent who cannot tolerate it in

England. Generous wines too have an evil influence. The wine must be poor for most bilious persons. Often even wines must be foresworn and some weak spirits-and-water form the sole beverage or food adjunct.

Abstinence, both as to meat and drink, must be the rule with persons who possess an insufficient liver.

This may seem a hard conclusion, but it is a simple expression of the facts. How far people can inherit an insufficient liver, is a matter about which there is not much doubt: we know gout runs in families, and so does biliousness. Has the advance of civilization a deleterious effect upon the viscera? Does the heir of all the ages find amidst his rich inheritance a waning capacity in the liver? It would seem so, at least, in many instances. There is no apparent impairment in the physique of some families whose history is known, as, for instance, the reigning family in England. But while this is quite true of some families, there is much evidence, and accumulating evidence too, that when the fathers have eaten sour grapes the children's teeth are set on edge. Dr. Murchison follows Dr. Budd in thinking most persons have competent livers, but "in others the liver seems only just capable of performing its functions under the most favourable conditions, and it at once breaks down under adverse circumstances of diet, habit or climate. This innate weakness of the liver is often inherited. The person is born with a tendency to biliary derangements."

When a person, man or woman, inherits an insufficient liver, gastronomic indulgence is thereby forbidden under a penalty.

CHAPTER XXXIII.

FOOD GIVEN OTHERWISE THAN BY THE MOUTH.

THERE are certain morbid conditions in which it becomes necessary to feed the patient otherwise than by the mouth. Especially does this become necessary in acute disturbances of the stomach. There are, however, other circumstances where such alimentation becomes desirable. Sometimes the appetite is so small, or the distaste for food so strong, that the amount taken by the mouth is insufficient for the wants of the system. Under such circumstances it becomes desirable to eke out these scanty meals by some food given *per rectum*. Under other circumstances, especially with marasmatic children, it is well to use enemas with oil.

Before proceeding to describe nutritive enemata and their constituents, it may be well to consider some anatomical and physiological facts which will guide us in practice. When food is given by the mouth it meets with the digestive ferments: first, the saliva; second, the gastric juice; third, the pancreatic secretion. The large intestine has none of these. The secondary digestion in the colon is somewhat of a misnomer, and as such misleading. "Of the particular changes which take place in the large intestine we have no definite knowledge, but it is exceedingly probable that in the voluminous caecum of the herbivora a large amount of digestion of a peculiar kind goes on. We know that in herbivora a considerable quantity of cellulose disappears in passing through the caecum, and even in man himself some is probably digested. We are driven to suppose that this cellulose digestion is carried on in the large intestine, though we know nothing of the nature of the agency by which it is effected. The other digestive changes are probably of a fermentative kind. Be that as it may, when normal digestion, properly so-called, is all but complete at the ileo-caecal valve, or whatever important changes still await the chyme in the larger intestine, the chief characteristic of the work done in the colon is absorption." (M. Foster, F.R.S.)

The only matter we are certain about is the absorption of soluble matter. Starch must then be converted into soluble products before

nutrient enema. Proteids must be converted into peptones for the same reason, *i.e.*, that there is absorption certainly. As to fat, it is highly questionable how far it is of use in a nutritive enema. Fat is emulsionised in digestion and in minute globules passes into the mouths of the lacteals in the intestinal villi; which, as M. Foster puts it, may be said to "eat the fat globules." But the intestinal villi cease at the ilio-cæcal valve; and therefore the fatty matters of nutrient enemata can be of little use. This is an important matter, on which it is well to have clear ideas. Bidder and Schmidt, in a case where there was a duodenal fistula, found that "Fats introduced into the lower part, where they could not be acted upon either by the bile or the pancreatic juice, were but slightly digested" (M. Foster). There must, however, exist some other mode by which fats can leave the alimentary canal (probably as soaps) for "the portal vein during digestion contains more fat than the general venous blood," the same authority tells us. It can, however, not, so far, be shown that fat can pass through the walls of the large intestine. The same authority quotes from Bauer—"Absorption from the large intestine after injection *per anum*, or through a fistula, has been observed not only in the case of soluble peptones and sugar but also in that of the white of egg, caseine, and starch." But there is no mention of absorption of fat. Consequently we may say that fat is practically useless in a nutrient enema because it is not absorbed; and that when the system needs fat it is of no avail to give it by enemata.

Of old a nutrient enema consisted of beef tea, the yolk of an egg, and milk. Of these different materials probably the extractives of the beef tea and the milk-sugar of the milk alone were absorbed—to any extent at least. If brandy were added, probably it, too, was absorbed. Of the other matter, as the casein and egg albumen, some were possibly absorbed.

The bowel really is adapted only to the absorption of matters which have been digested above in the small intestine. If matters be introduced into it from the other extremity, they ought to have been subjected to a like action previous to their occupation of the lower bowel. The ilio-cæcal valve bars their entrance into the small intestine where digestion—as differentiated from mere absorption—takes place. If not predigested they are not fitted for absorption.

Nutrient enemata then can conveniently consist of soluble carbohydrates and peptones. The disagreeable glue-like flavour of most peptones constitutes no objection to their use here. Indeed peptones

and peptonised gruels, are excellent matters for nutritive enemata. Malt preparations, malt foods, and malt extracts are also good material for enemata. Or milk-sugar, or dextro-saccharum, or even levulose sugar, especially as treacle.

When acute gastric disturbance is on foot, then the nutritive enema should consist of soluble carbo-hydrates, meat peptones, and if necessary a little brandy. When there is anorexia and a febrile state co-existent, the utility of albuminoids is less certain: as probably no histogenesis then goes on (as pointed out in a previous chapter). But of the utility of matters furnishing grape-sugar to the system in its hour of need there can be no question. Recently in a case of chronic bronchitis and emphysema the appetite dropped dangerously low even for a lady in bed in a warm bedroom. Here a nutritive enema of Mellins's Food and peptonised milk night and morning just made the difference betwixt slow starvation and improvement. I could have liked to have something to say about the utility of nutritive enemata in the attacks of acute gastric disturbance in phthisis; but unfortunately am not in a position to say much. On *a priori* grounds there is a great deal to say; but how the actual facts will turn out is not so certain. If readers of this work who try nutritive enemata in such cases would put their experience on record we should soon have a body of evidence which would be highly valuable, whether for or against. If favourable, our hands are strengthened in a very trying crisis. If it turn out but a broken reed it will be well to know it, so as not to put faith in an untrustworthy measure at our time of trial.

A large tablespoonful of soluble carbo-hydrate as malt-extract, or Mellins's, or other malt preparation, or milk sugar, or a small teaspoonful of meat peptones in an oz. of warm water, would form something like what is required for a nutritive enema. Such a small bulk would probably be well tolerated by the not very sensitive large intestine. And repeated twice, thrice, or oftener daily would make a substantial addition to the body-income.

How long nutritive enemata may be continued depends upon the condition of the anus. If this become very sore and sensitive they may have to be discontinued after a short trial. (The utility of an opium suppository in such anal sensitiveness suggests itself, or some other local application). If the anus remains insensitive the nutritive enemata can be continued for a longer period. On this matter we have much to learn. When we give an enema to flush out the contents of the large intestine we remain liable to distend the pouches of the colon in which the faeces lodge. But for

nutrient enemata a small bulk is desirable in order that it may be retained and not be cast forth.

We can see a great future for nutritive enemata now that the proper materials for them can readily be procured; and increasing knowledge will teach us many minor points to be attended to, of which probably something can be said in a future edition.

Very recently "rectal alimentation" by nutrient suppositories has been advocated; and in some cases has been found useful. The component factors of nutrient suppositories will be limited by the anatomico-physiological facts of the wall of the colon and its capacities (pointed out in the early part of this chapter), especially what is said about the non-absorption of fat by the bowel. That nutrient suppositories of soluble carbo-hydrates and peptones may come to have a high practical value seems eminently probable. Even if these nutrient bodies have but a limited scope, this may still be enough to give them an incalculable value in times of emergency, when life is quivering in the balance and a trifle will cast the beam. And it is in our capacity to provide these apparently little matters in the nick of time which constitutes a great part of the utility of the medical profession. Sometimes a practical grasp of the exigencies of the case, and the knowledge how to cast the quivering balance, far outweigh in real value an elaborate but sterile diagnosis of minutiae in such hours of need. Emergencies are indeed often the test not only of the possession of knowledge, but of the capacity to make use of it. A nutrient enema at the proper moment may have life or death hanging on it.

Of hypodermic injections of nutritive fluids little can be said. Accounts of such methods of feeding have been published doubtless, chiefly by foreign physicians; but they appear rather to belong to the curiosities of medicine than as part of the working apparatus of daily practice. For matters of small bulk, as potent medicines, the injection syringe is of priceless value; but of concentrated foods adapted for injection purposes we know nothing.

Transfusion of blood scarcely comes under the heading of dietetics.

Then come inunctions of oil. How fat is absorbed by the skin is a matter on which little is known; at least, nothing about it is known to me; but about the fact there can be no question. When attached to the West London Hospital numbers of poor starvelings were among the out-patients. With many of these mites the addition of some cod-liver oil to their scanty fare caused much improvement. With others, however, it became necessary to resort to oil

inunctions. The effects varied: some babies began to be better at once and rapidly, while with others the progress was more gradual. But on the whole the results were satisfactory.

In some instances such inunctions have been found useful in adults where the assimilative organs were unequal to the demand. It is not unusual for phthisical patients to apply cod-liver oil to the chest when it is not readily taken by the mouth. The oil is absorbed, and does good, but the plan is an unsavoury one.

When oil inunctions are indicated the room should be made warm and all draughts excluded; else the exposure may be dangerous. (A poor little infant readily feels any heat loss). Then the skin must be made perfectly clean with hot water and soap, with a piece of flannel or a sponge. After this has been done the oil can be applied. It is well to allow the oil to get warm by the fire while the preliminary operations are going on. About half an ounce (or tablespoonful) is enough at once. About a teaspoonful should be poured into the palm at once, and this should be rubbed well in over the trunk and limbs. How much of this is absorbed I do not know. The undergarment of course becomes oily, and how much absorption goes on after the rubbing I do not know. But, however it comes about, the process gives satisfactory results.

Adults, or children of larger growth, who require oil inunctions are usually—or ought to be—in bed, (in order to reduce the body-expenditure). With these it has been found convenient to anoint one half of the body at once; taking the other moiety in turn. In certain cases this method of feeding proves very useful. In cold weather such coating of oil is found very comfortable. It is not a cleanly method, however, and has nothing to be said for it except its utility in times of need.

Such, then, are the methods of feeding otherwise than by the mouth. They are all objectionable and repugnant to the feelings, and to sensitive persons simply intolerable. Still, there are occasional instances under which these methods are a great good, and it is well to know of them and about them when the emergency arrives. As to the choice of oil, probably an animal oil is more efficacious than a vegetable oil. Of these neats-foot oil is the best offered. When this is offensive then olive oil can be resorted to. Probably this was the oil used by the early Christians, who were anointed with oil and prayed over by the elders of the Church, according to St. James.

Inunctions of oil are less frequently called for than enemata or enemata, which are often of much service.

CONCLUSION.

THE perusal of the foregoing chapters will, it is believed, convince most readers that there is much more to be said for food and feeding in disease,—alike in acute maladies and mere chronic affections,—than at first sight seems probable. To know what food is required in each morbid condition is a matter too little attended to in medical teaching. Medical teaching is conducted in hospitals, and in hospitals for obvious reasons the dietary is arranged in scales of varying liberality. This unavoidably leads to routine—hard and fast lines which are not elastic. Consequently, dietetics are matters picked up by the medical man mostly from his private patients; and probably his first real lessons come from some intelligent patient, or friend of a patient. Beef tea he knows from the hospital, and milk and lime-water or seltzer water; but the rest he acquires as he can. Some men pick up a great deal, others but a little. The Professor of Organic Chemistry tells him little or nothing about the food-value of food; the Professor of Physiology describes digestion to him, but does not describe the various forms of food; the Professor of Practice of Physic gives him directions as to the choice of medicines, but probably never mentions food except when discoursing on diabetes; the Professor of Materia Medica gives a careful account of the different members of the Pharmacopœia, including their actions within the body, but as for food, he has no word to say; while the Professor of Surgery may speak of a bread-poultice or a beef-steak for a black eye; but that is all.

If some systematic lectures were delivered, even if only of an elementary character, the student would have some broad rules to guide him in his choice; instead of being left pretty much like a ship without a rudder. It may be thought by some readers presumptuous for a person not actually engaged in medical teaching to criticise the existing silence, or to make suggestions; but it does seem to me that instead of getting up pharmacology and the appearance of crude drugs it would be well to substitute some account of foods and feeding. When the medical man had to choose and select his raw material, from which, as an apothecary, he prepared extracts, tinctures and infusions, then such knowledge certainly had its value.

But now-a-days a medical man trusts his chemist and his druggist for his medicines, and rarely even makes a fresh brew; consequently, this part of his studies is useless, and money is expended to the student, and an inroad upon time which has many other uses upon it, and which is too precious, in every way, to be wasted. If this time were devoted to some account of food in health and sickness; of the preparation of food in relation to its digestion; of prepared food for infants and invalids and those actually ill; of the predigestion of foods; of the relation of various foods to sundry morbid conditions; it would be much better.

Possibly the absence of such lectures is to some extent due to the fact that a text-book available for such lectures did not exist. Dr. Pavy's book is most interesting and instructive; and the same may be said of King Chambers's *Manual of Diet in Health and Disease*; but neither form systematic treatises suitable for lecturing purposes. The writer has some hope that the present work has some claims to be such a book; but this may be no more than the vanity which is said to cling to authors.

Lectures are delivered yearly before the Royal College of Physicians of London, and we might have thought that the authorities of the College would have found some one to take up the subject of Food in Sickness during the last decade. Their inactivity on the subject is on a par with their indifference in the matter of medical teaching. When a fish smells about the head it is useless to look for freshness at the tail, says a Turkish proverb, and so long as this ruling body in matters medical maintains an attitude of contempt and supine indifference on the matter of food, so long nothing will be done. As an examining body a change would at once be warranted if a question on Dietetics were asked at the higher examination for the membership; but of this there seems as yet no prospect.

There is a better prospect in the United States, where the subject of Dietetics is in a more advanced condition. "The Journal of Reconstructives" is a step forward which affords some satisfaction; while the efforts of the enterprising firm now engaged, to show various forms of food are worthy of a commendation. As it is the knowledge of foods and how to use them, and of the various preparations in the market, that is a revelation to the feeding of the sick and infirm will be wrought. And the concluding sentence is the expression of his belief that the long-desired revolution is close at hand.

THE
PUBLICATIONS
OF
WILLIAM WOOD & COMPANY,
NOS. 56 & 58 LAFAYETTE PLACE,
NEW YORK.

Anatomy.

Weisse, Faneuil D., M.D.,

Professor of Physiology and Anatomy in the Medical Department of the University of the City of New York; and Professor of Anatomy in the New York College of Physicians and Surgeons.

THE DEMONSTRATOR: A Work for the Use of the Students of Human Anatomy. By F. D. WEISSE, M.D. Retrospect for Practitioners of Medicine and Surgery. Illustrated by S. J. COOPER, M.D. Consists of two hundred full-page, three-quarter plates, each containing a fine anatomical preparation.

Quain's

ELEMENTS OF ANATOMY. By RICHARD W. QUAIN, M.D., LL.D., F.R.S., LL.D., Emeritus Professor of Anatomy and Physiology in the University of Cambridge; and ALLEN THOMPSON, M.D., LL.D., F.R.S., LL.D., Professor of Anatomy in the University of Glasgow; and EDWARD ALBERT SMITH, M.D., Assistant Professor of Physiology in the University of Cambridge. New York: Published by William Wood & Company, 56 and 58 Lafayette Place. 1890. 8vo. 744 and 944 pages, illustrated. Price, \$12.00.

It is a pleasure to recommend this work, which has been read and found to be a most valuable and complete work, not only for the student, but also for the practitioner. The illustrations are of the highest quality, and the text is clear and concise. The work is a valuable addition to the library of every student and practitioner. —*The Lancet*, Oct. 1, 1890.

Wood's Pocket Manuals.

THE ANATOMICAL REMEMBRANCER, or, Cyclopedia of Human Anatomy. By W. WOOD, M.D. A complete and concise summary of the facts of human anatomy, arranged in alphabetical order, and adapted for the use of students and practitioners. New York: Published by William Wood & Company, 56 and 58 Lafayette Place. 1890. 207 pages, illustrated. Price, \$1.00.

"A valuable little book, which will be found of great use to the student and practitioner. It is a complete and concise summary of the facts of human anatomy, arranged in alphabetical order, and adapted for the use of students and practitioners. The illustrations are of the highest quality, and the text is clear and concise. The work is a valuable addition to the library of every student and practitioner. —*The Lancet*, Oct. 1, 1890.

"A valuable little companion for the student and practitioner. It is a complete and concise summary of the facts of human anatomy, arranged in alphabetical order, and adapted for the use of students and practitioners. The illustrations are of the highest quality, and the text is clear and concise. The work is a valuable addition to the library of every student and practitioner. —*The Lancet*, Oct. 1, 1890.

FOR INDEX SEE LAST TWO PAGES.

*Anatomy.***Ranney, Ambrose L., A.M., M.D.,**

Assistant Professor of Anatomy, and late Lecturer on Genito-urinary and Minor Surgery in the Medical Department of the University of the City of New York; late Surgeon to the Northern and North-western Dispensaries; Resident Fellow of the New York Academy of Medicine; Member of the Medical Society of the City of New York; Author of "The Applied Anatomy of the Nervous System," "A Practical Treatise on Surgical Diagnosis," "The Essentials of Anatomy," etc.

PRACTICAL MEDICAL ANATOMY. A Guide to the Physician in the Study of the Relations of the Viscera to each other in Health and Disease, and in the Diagnosis of the Medical and Surgical Conditions of the Anatomical Structures of the Head and Trunk. Illustrated. Sold only by subscription. See page 54.

Ellis, George Viner, M.D.,

Professor of Anatomy in University College, London, and

Ford, G. H., Esq.

ILLUSTRATIONS OF DISSECTIONS. In a series of original colored plates, representing the dissections of the human body, with descriptive letter-press. The drawings are from nature by Mr. Ford, from directions by Prof. Ellis. Volumes I. and II. Containing fifty-six full-page chromo-lithographic plates. Sold only by subscription. See page 54.

2. "When, in the second series, we succeeded in presenting our subscribers with 'Savage's Female Pelvic Organs,' with its full-page lithographic plates, we supposed we had reached the extreme limit in reproducing expensive books at so low a price, but these two volumes of Ellis and Ford exceed even that. It is simply wonderful, and cannot fail to *amplify* the knowledge of the value of this series of publications. It would have been impossible to accomplish such results, save in a library such as this, in which *all* the volumes have an *equal* sale."

Bock, C. E., M.D.,

Professor at Leipsic.

ATLAS OF HUMAN ANATOMY. With Explanatory Text. Containing thirty-eight steel plates, most of them hand-colored, of the bones, muscles, vessels, and nerves of the human body; organs of sense, eye, ear, nose, and tongue, respiratory apparatus, abdominal and pelvic viscera, organization of foetus, the teeth, with the embryonary organs of the male and female. One volume, large 4to, maslin. Price, \$15.00. Sold only by subscription. See page 64.

Ziegler, Ernst,

Professor of Pathology, University of Tübingen.

A TEXT BOOK OF GENERAL PATHOLOGICAL ANATOMY AND PATHOGENESIS. Translated and edited for English students by DONALD MACALISTER, A.M., M.B., Member of the Royal College of Physicians; Fellow and Medical Lecturer of St. John's College, Cambridge. Parts I, II, & III. Sold only by subscription. See page 53.

Holden, Luther, M.D.,

Extraordinary and Member of the Council of Physicians in the Royal College of Surgeons of England; Lecturer on Surgery at St. George's Hospital, London; and at the Johns Hopkins Hospital.

Shuter, James, F.R.C.S., M.A., M.B., Cantab.,

Assistant Surgeon to the Royal Free Hospital, and Lecturer on Physiology, and Assistant Lecturer on Anatomy, in St. George's Hospital, London.

HUMAN OSTEOLOGY. Comprising Descriptions of the Bones, with Definitions of the Attachments of the Muscles, the General and Muscular Structure of the Skeleton, Development, &c. Sixth Edition. With fifty-six full-page lithographic plates, and twenty-nine woodcut engravings. Sold only by subscription. See page 51.

2. "This is another of these marvels of bookmaking, for which Wood's Library has been celebrated."

1. not coming.

Delafield, Francis, M D.,

Professor of Pathology and Practical Medicine, University of Physicians and Surgeons, New York City.

Prudden, T. Mitchell, M.D.,

Journal of Polymer Science: Polymer Chemistry Edition, Vol. 19, 1111-1120 (1981)
© 1981 John Wiley & Sons, New York, Inc. 0360-6376/81/0019-1111\$01.00

A HANDBOOK OF PATHOLOGICAL ANATOMY AND HISTOLOGY

Introductory Section on Post-Modern Examinations and the Medical Profession, and Examining Disabled Teachers. One volume, 8 1/2" x 11" paper, 1990, with one hundred and forty-six exercises from oral, written, and audio-tape examinations. Price, \$5.00.

3. The first edition of this work, entitled "Post-mortem Examination," serves as a guide for those persons who might have to perform post-mortem surgery.

In its second edition the scope of the work has been very much extended so that it is now intended to supply all the needs of students and practitioners who wish to make a study of the various reasons of disease, be it that of its clinical symptoms. The point of view of the author is that of the physician who wishes to find out what are the lesions of the body, and he is in the habit of seeing, rather than that of the pure pathologist.

Delafield, Francis, M.D.,

Journal of the American Statistical Association, Vol. 90, No. 430, of Probability and Statistics

STUDIES IN PATHOLOGICAL ANATOMY. Volume L, pp. 219-222.

[illegible]

PART I, Volume II, is now available. Price, \$2.00. Order at once.

Professor DeMelet's *Standards in Plantology of Argentina*, to which we have referred frequently, is the appearance of a number of plants, since this can be collected from a single magnificent volume, now octavo size, and is nicely illustrated with numerous full-voilette page plates. The author's object is to describe and bring to light the various sets of diseases from the various parts of the plant, and the various parts of the plant, and the various parts of the plant.

Results were obtained from a series of 10 experiments. A typical example of a single experiment is shown in Figure 1. The first 10 min of the experiment were devoted to the establishment of a steady state. The following 10 min were devoted to the measurement of the rate of change of the concentration of the product. The rate of change of the concentration of the product was determined by measuring the change in the optical density of the solution at 440 nm. The rate of change of the concentration of the product was determined by measuring the change in the optical density of the solution at 440 nm. The rate of change of the concentration of the product was determined by measuring the change in the optical density of the solution at 440 nm.

Hilton, John, F.R.S., F.R.C.S.

REST AND PAIN 3695

1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 26

1. The first group of people who are interested in the study of the history of the United States are the people who are interested in the history of the United States.

*Pathology, Diseases of Joints, etc.***Wagner, Ernst, M.D.,**

Professor of General Pathology and Pathological Anatomy in the University of Leipzig; Director of the Medical Polytechnic of Leipzig.

A MANUAL OF GENERAL PATHOLOGY. For the Use of Students and Practitioners of Medicine. Translated from the Sixth German Edition by JOHN VAN DYKE, A.M., M.D., Professor of Anatomy and Histology in the Medical School of Syracuse University, N. Y., and E. C. SEGTIN, M.D., Clinical Professor of Diseases of the Mind and Nervous System in the College of Physicians and Surgeons, New York. One volume, Svo, 328 pages. Price, muslin, \$5.50; leather, \$6.50.

"No book in the English language gives such a thorough *resumé* of the elements of medicine. The work, containing over seven hundred closely printed pages, is a storehouse of carefully sifted and well-arranged information in all departments of pathology. Unlike many translations, the text is lucid and pleasant reading, and the work is altogether one of the most valuable which has been issued in the English language. It will be invaluable to the student of pathology, and should be in every physician's library."—*Philadelphia Medical Post*.

"It is a book full of knowledge for the practitioner. A medical student could not better commence work than by reading this book in rough. It has all the modern doctrines and discoveries."—*The Clinic*.

"The views now held by progressive investigators in this branch of medical science are given in a marvelous mass of well-arranged and carefully digested facts, embracing all departments of pathology, and furnishing minute and detailed information on every subject pertaining to the morbid state and disease in all its possible forms. Wagner's systematic arrangement and skill in classification are admirably aided by lucidly elucidating the bearings of facts upon the processes of disease."—*American Journal of the Medical Sciences*.

"Messrs. Wm. Wood & Co. have placed the publication of this country under renewed obligations by the presentation of this immense work. The most popular text book in Germany, having passed through six editions in that country, the only wonder is that it has not been given to the profession of America before this. The most cursory glance through it is sufficient to convince one of the excellence of the work, which bears the imprint of one of

the greatest German minds. The bibliography of the work before us is very copious, and evinces a research rarely entered into by authors. The work consists of four parts, devoted respectively to General Nosology, General Etiology, General Pathology, Anatomy and Physiology, and Pathology of the Blood."—*Pennsylvan Journal of Medicine*.

"The work is one which will amply repay perusal; in fact, no physician should neglect to read it, and the student should make it one of his most cherished text-books."—*Buffalo Medical and Surgical Journal*.

"Clear and concise statements, short and simple sentences, present the condensed thought with which this volume is filled. This is as it should be. Those who desire more than carefully arranged results will find in the copious bibliography and abundant intercalated references the sources from which the information has been derived. An index of authors and another of subjects render invaluable aid. We consider this book as one of the most valuable of the many medical works translated from German into English. Its application is general, for it will be found not only upon the shelves of the surgeon and the chemist, but also in the hands of the medical student, who can rely upon no safer and surer guide through the obscurity which surrounds the causes and phenomena of disease. The modern explanation of 'teaching odd' may be seen, as well as the more elaborate theories with regard to fever. Above all, it contains what is to be found in no other single volume—it is filled with a complete and comprehensive series of systematically arranged abstracts, which form an epitome of general pathology."—*Boston Medical and Surgical Journal*.

Longstreth, Morris, M.D., etc.

RHEUMATISM, GOUT, AND SOME OF THE ALLIED DISEASES. Sold only by subscription. See page 54.

Barwell, Richard, F.R.C.S.

Professor of Clinical Surgery.

A TREATISE ON DISEASES OF THE JOINTS. Illustrated by numerous engravings on wood. Sold only by subscription. See page 55.

Lyman, Henry M., A.M., M.D.,

Professor of Physiology and Neurology in the Medical College of the University of Toronto, and Lecturer on the Diseases of the Nervous System in the University of Michigan.

ARTIFICIAL ANÆSTHESIA AND ANÆSTHETICS. Sold only by subscription. See page 55.

*Diseases of Old Age, Electricity.***Index Rerum**

For the Use of Physicians and Others. Sent by Express, Postpaid, \$1.00. 4

Charcot, J. M., M.D.,

Dr. Jean Martin Charcot, M.D., of Paris, France, is the celebrated Master of the
 Medicine of the Salpêtrière Hospital, Paris, France, and Professor of the
 Faculty of Medicine, Paris, France, and of the Salpêtrière Hospital, Paris, France.

THE DISEASES OF OLD AGE. Translated by L. HARRISON HENRY, M.D., with
 numerous additions by A. L. Loomis, M.D., etc., Professor of Pathology and
 Practical Medicine in the Mount Sinai Department of the University of New York.
 New York: Consulting Physician to the Charity Hospital, and to the
 Out-Door Receiving to the Central Dispensary; Visiting Physician to the
 Hospital to the Mount Sinai Hospital, etc., etc. Sold by Express, Postpaid.
 See page 55.

Beard, Geo. M., A.M., M.D.,

Professor of New York Academy of Medicine, and of the University of New York,
 and of the Academy of Medicine, New York, and of the University of New York.

Rockwell, A. D., A.M., M.D.,

Professor of New York Academy of Medicine, and of the University of New York,
 and of the Academy of Medicine, New York, and of the University of New York.

**A PRACTICAL TREATISE ON THE MEDICAL AND SURGICAL USES OF
 ELECTRICITY, INCLUDING LOCALIZED AND GENERAL FARADIZA-
 TION; LOCALIZED AND GENERAL GALVANIZATION; ELECTROLYSIS
 AND GALVANOCAUTERY.** Fourth Edition, Revised, by A. D. ROCKWELL,
 M.D. One volume, 8vo, 788 pages, with colored plates, and illustrations.
 Price, in cloth, \$5.00; in leather, \$6.00.

"The impression it gives at first sight is
 very favorable; it is beautifully printed, and
 illustrated with woodcuts in two volumes, and
 is a most valuable work for the student."

"The volume is a very good one, and is
 a most valuable work for the student."

"Retreats are numerous, and are of a
 nature which is highly to be desired. We
 are sure to find a most valuable work for
 the student, and a most valuable work for
 the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

that we teach in the Medical Academy of New York."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

"The subject is a most valuable one, and
 is a most valuable work for the student."

*Electricity, Massage.***Rockwell, A. D., A.M., M.D..**

LECTURER on Electricity at the New York State Woman's Hospital; Member of the American Neurological Association; Fellow of the New York Academy of Medicine; Member of the New York County Medical Society, etc.

LECTURES ON ELECTRICITY (DYNAMIC AND FRANKLINIC) IN ITS RELATIONS TO MEDICINE AND SURGERY. One volume, 8vo, 122 pages, 11 striped, muslin. Price, \$1.25.

"It contains all that is practical and useful on this subject to the practitioner; it is concise and interesting to read, and should be in the hands of every practitioner."—*Western Lancet*.

"Practitioners wishing to obtain a knowledge of the methods of applying and indicating the use of electricity in the treatment of diseases cannot find more useful information in any small compass than is contained in these lectures. We take pleasure in commending it to their notice."—*Canada Journal of Medical Science*.

"These lectures embrace the more important practical matters, as the manipulations of apparatus, and the fields of disease where electricity will promise the most good."—*Obstetric Gazette*.

"At the close of the book there are a series of pithy paragraphs, giving, in plain language, the practical applications of the different currents to various affections, as shown by the personal experience of the writer, and those of other competent specialists."—*Medical Review*.

Graham, Douglas, M.D.,

Member of the Massachusetts Medical Society.

A PRACTICAL TREATISE ON MASSAGE. Its History, Mode of Application, and Effects, Indication and Contra-indication. One handsome volume, 8vo, 300 pages. Price, in cloth, \$2.50.

"We commend the work to those who are interested in the merits of friction, kneading, massage, rolling, and percussion of the external tissues of the body, with a view to their curative, palliative, or hygienic effects."—*Medical and Surgical Journal*.

"Coming from the hands of a medical man, the work may be also pronounced authority in explaining this resource to the student as a guide to its practice."—*Medical Review*, December 31, 1884.

"A complete history, from the beginning

until now, will be found in Dr. Graham's excellent volume."—*Baltimore Sun*, December 14, 1884.

"The subject of massage has been one toward which the attention of the profession has been directed more and more of late years. As a therapeutic measure, it has been growing steadily in favor."—*Medical and Surgical Reporter*, October 20, 1884.

"This is a valuable professional book for which, and for the subject to which it is devoted, we venture to predict distinguished success."—*The Astor*, January 1, 1885.

Morgan, Chas. E., A.B., M.D.

ELECTRO-PHYSIOLOGY AND THERAPEUTICS: BEING A STUDY OF THE ELECTRICAL AND OTHER PHYSICAL PHENOMENA OF THE MUSCULAR AND OTHER SYSTEMS DURING HEALTH AND DISEASE, INCLUDING THE PHENOMENA OF THE ELECTRICAL FISHES. One volume, 8vo, 114 pages, illustrated with five colored drawings, muslin. Price, \$6.50.

"This book is a mine of knowledge to the student in the department of science to which it belongs."—*Medical Review*. "It is a most thorough and comprehensive treatise on the subject."—*Medical Review of Medicine*.

"There is nothing in the English language which at all approaches it as regards the scientific treatment of the whole subject of electricity."—*Medical Review of Medicine and Pharmacy*.

Erb, Dr. Wilhelm,

Professor of Physiology at Leipzig.

HAND-BOOK OF ELECTRO-THERAPEUTICS. Illustrated by 39 wood-engravings. Second edition, 1884. 8vo, 120 pages, 75.

*Fever, Diseases of the Liver.***Seguin, E., M.D.**

MEDICAL THERMOMETRY AND HUMAN TEMPERATURE. One volume, 8vo, 445 pages, illustrated with diagrams, muslin. Price, \$4.50.

"The subject-matter of this volume is of the highest importance, and the profession is indebted to its author for the pains he has manifested in bringing it to our notice. The book is worthy of careful study by every medical student and studious practitioner."—*Medical History of Medicine and Surgery.*

"This elaborate, elegantly printed, and illustrated work contains a full description of everything pertaining to thermometry, the various thermometers, their value, uses, etc. Numerous tables are given from various American and foreign authors and writers,

exhibiting the temperature of various parts. Among these are Dr. Quinquaud's on Yellow Fever, where the temperature is noted in two cases, which terminated fatally and fully. The illustrations, temperature, and statistics are rightly put in number, and are carefully and simply executed. We can only commend the work to all learners as a valuable contribution to the study and culture of one of the most important and interesting subjects of investigation and perception, now so generally being and now daily attracting the attention of physicians."—*Charleston Medical Journal and Review.*

Gregory, George, M.D.,

Professor of the College of Physicians of London; Physician to the St. Thomas' Hospital; and Honorary Member of the National Institute of Hygiene, etc.

LECTURES ON THE ERUPTIVE FEVERS: As now in the Course of Delivery at St. Thomas' Hospital, in London. With Notes and Appendix containing the most recent opinions on exanthematic pathology, and also statistical tables and several plates, by H. D. BRIDLEY, M.D., Physician of the New York Hospital; Fellow of the New York College of Physicians and Surgeons, etc., etc. One volume, 8vo, 319 pages, muslin. Price, \$3.00.

"The very best which has yet been published on Eruptive Fevers; and one which it should be the duty of every physician to provide himself with."—*Northern Leader.*

"This work abounds with valuable information in regard to a class of diseases of very frequent occurrence and of fatal mortality."—*St. George's.*

Wilson, James C., M.D.,

Assistant Surgeon to the Philadelphia Hospital; and the Hospital of the University of Pennsylvania; Lecturer on the Principles of Physical Diagnosis at the Jefferson Medical College; Fellow of the American College of Physicians, etc.

A TREATISE ON THE CONTINUED FEVERS. With an Introduction by J. M. DuClos, M.D., Professor of the Practice of Medicine and Clinical Medicine at the Jefferson Medical College; Physician to the Pennsylvania Hospital; Consulting Physician to the Children's Hospital; Fellow of the American College of Physicians, Philadelphia, etc. Sold only by subscription. See page 55.

Murchison, Charles, M.D., LL.D., F.R.S.,

Professor of the Principles of Medicine, the History of the Fever, and the Principles of Pathology in the University of Edinburgh; Fellow of the Royal Society, etc.

CLINICAL LECTURES ON DISEASES OF THE LIVER, JAUNDICE AND ABDOMINAL DROPSY. Including the Chosen Lectures on Pathology, Demonstrations of the Liver, delivered at the Royal College of Physicians, New York, in preparation.

"A book which surpasses in clinical usefulness any work on the diseases of the liver yet published."—*Medical History of Medicine and Surgery.*

"It is the most instructive, the most beautiful, and the most valuable work on the subject in our language."—*St. George's Medical Journal and Review.*

"A monograph on Liver Affections, which may be profitably read by the student and in which the principles of the system of medicine have been put in its application, and

thoroughly identified with the principles which it is based on."—*Western Medical Review.* "Murchison's volume is the best treatise on the diseases of the liver, and is a valuable addition to the medical literature of the day. It is a book of facts, a book of the day, written in a simple, clear, and concise style, and is a most valuable and interesting work. It is a book of facts, a book of the day, written in a simple, clear, and concise style, and is a most valuable and interesting work. It is a book of facts, a book of the day, written in a simple, clear, and concise style, and is a most valuable and interesting work."—*The Medical Review.*

*Diseases of the Liver, Practice of Medicine.***Murchison, Charles, M.D., LL.D., F.R.S.,**

Fellow of the Royal College of Physicians; Physician and Lecturer on the Principles and Practice of Medicine, St. Thomas' Hospital; Vice President and Consulting Physician, London Fever Hospital, &c.

ON FUNCTIONAL DERANGEMENTS OF THE LIVER: Being the Croonian Lectures delivered at the Royal College of Physicians in March, 1871. One volume, 12mo, 182 pages, illustrated, muslin. Price, \$1.75.

"This monograph is an excellent contribution to medical literature, and is worthy of a place in every medical library."—*Chirurgisch-Lincoln Observer*.

"There is not a physician anywhere who will not read with avidity and profit this admirable exposition of this classic and most important subject."—*American Medical Weekly*.

"This work does not need special criticism at our hands, for its author is known to be a distinguished master of the subject on which he has discoursed. His name has long been identified with careful original researches in this department of pathology, and he has

given in this little book, the reprint of the Croonian lectures, a clear and comprehensive summary of the present state of medical knowledge concerning the relation of functional hepatic disorders to the various vital processes."—*Boston Medical and Surgical Journal*.

"This admirable volume, the work of a practitioner who has brought to the investigation of this subject all the light of modern physiological science will exert a decided influence on the minds of medical men, and awaken renewed interest in the complaints of which it treats."—*The American Practitioner*.

Frerichs, Dr. Fried. Theod.

A CLINICAL TREATISE ON DISEASES OF THE LIVER. Translated by CHARLES MURCHISON, M.D. In three volumes, 8vo. Illustrated by a full-page plate and numerous wood-engravings. Sold only by subscription. See page 57.

Bristowe, John Syer, M.D.,

Author of

DISEASES OF THE INTESTINES AND PERITONEUM. Comprising Articles on—Enteralgia, by JOHN RICHARD WARDLELL, M.D.; Enteritis, Obstruction of the Bowels, Ulceration of the Bowels, Cancerous and other Growths of the Intestines, Diseases of the Cecum and Appendix Vermiformis, by JOHN SYER BRISTOWE, M.D.; Colic, Colitis and Dysentery, by J. WARBURTON BLOCH, M.D.; Diseases of the Rectum and Anus, by THOMAS BLIZZARD CURLEW, F.R.S.; Intestinal Worms, by W. H. RANSOM, M.D.; Peritonitis, by JOHN RICHARD WARDLELL, M.D.; Tubercle of the Peritoneum, Carcinoma of the Peritoneum, Affections of the Abdominal Lymphatic Glands, and Ascites, by JOHN SYER BRISTOWE, M.D. Sold only by subscription. See page 57.

Bennett, John Hughes, M.D., F.R.S.E.,

Professor of the Principles of Medicine, and Senior Professor of Clinical Medicine in the University of Edinburgh; Director of the Practice of the Royal Dispensary and Pathologist to the Royal Infirmary, Edinburgh; Honorary Member and Honorary President of the Royal Medical Society of Edinburgh, &c.

CLINICAL LECTURES ON THE PRINCIPLES AND PRACTICE OF MEDICINE.

Fifth American from the Fourth London Edition. With five hundred and thirty-seven illustrations on wood. One volume, 8vo, 1022 pages. Price, mass. in, \$5.00; leather, \$6.00.

"We recommend this volume with the most unqualified praise to the attentive consideration of the practitioner and students. We have not with us now work of late years on the principles of medicine more likely to advance the true and rightful study of our art."—*British and Foreign Medical Review*.

"When a book, especially so large a book as this, becomes a fourth edition, it may be considered to be pretty independent of new views and new works."

"It would be scarcely too much to say that Dr. Bennett's book is one of the soundest and practical of medicine. To him as much as to any other physician, are due the changes

which have come over the practice of medicine, and for this, more than for anything else, we value him and the books which record his views and essays."—*Lancet*. "This is a most valuable book, and records work hard and weary, which will secure for the author a lasting reputation as a physiologist and practitioner."—*Lancet*.

"One of the most valuable books which have yet been issued from the medical press. No student devoted to the profession will fail to peruse these lectures, and acquaint himself with the doctrines of so eminent an authority in the art of medicine."—*New York Journal of Medicine*.

Mammals.

Code of Medical Ethics

ADOPTED BY THE AMERICAN MEDICAL ASSOCIATION, 18
One volume, 32mo, 30 pages, muslin. Price, 10c.

Bigelow, Jacob, M.D.,

He is a Lecturer in 'Contemporary Modernism' in the M.A. programme in History and Literature in Harvard University. Director of the American Studies Centre, and Vice-President of the Massachusetts Modern Society.

BRIEF EXPOSITION OF RATIONAL MEDICINE. To wit: A. S. P. & Co. The
Paradise of Doctors. A Fable. One volume, 12mo, 69 pages, new. Price,
50c.

NATURE IN DISEASE. Illustrated in various Diseases and Passes. Theology and Medical Miscellaneous Writings, chiefly on morbid subjects. The second edition enlarged. One volume, 12mo, 407 pages, new, in P.B. 6, 81, 25.

Hooper's

PHYSICIAN'S VALE MEICUM: A Manual of the Principles and Practice of Physic, as far as Our Line of General Pathology, Therapeutics, and Hygiene is Concerned. By WILLIAM ALGISTUS GUY, M.B., Cantab., F.R.S., and JOHN HENRIE, M.D., Lond., F.R.S. Volumes I. and II. London: Longmans, Green, & Co., 1891. Sold only by subscription. See page 52.

Curtis, Edward, A.M., M.D.,

MANUAL OF GENERAL MEDICINAL TECHNOLOGY, INCLUDING PRESCRIPTION WRITING. One volume, 32mo, 394 pages, 50c. net. Pp. 1-81, 60. Wood's Pocket Manuals.

*The metric system is explained in a manner that is both helpful and attractive, and will undoubtedly be as much in the way of removing the "fear" of the subject as it is of those who are "familiar" with it, as reflected on the *Ph. M.* of 1981, p. 10. The *Ph. M.* of January 1981, p. 10, p. 11, and p. 12, is a tersely composed, but also on the subject, as referred to above, in a commendable and attractive sort of way that reads like a highly conscientious effort to do

patients. It is worth noting that, in contrast to the findings of the present study, the results of the study by Loxton *et al.* (2000) did not show any association between the use of alcohol and the use of tobacco.

[illegible]

Roosa, D. B. St. John, M.D.,

AVIATION ROCKET MEDICAL HANDBOOK, 1968, 1980, 1982, 1984, 1986, 1988, 1990, 1992, 1994, 1996, 1998, 2000, 2002, 2004, 2006, 2008, 2010, 2012, 2014, 2016, 2018, 2020, 2022, 2024, 2026, 2028, 2030, 2032, 2034, 2036, 2038, 2040, 2042, 2044, 2046, 2048, 2050, 2052, 2054, 2056, 2058, 2060, 2062, 2064, 2066, 2068, 2070, 2072, 2074, 2076, 2078, 2080, 2082, 2084, 2086, 2088, 2090, 2092, 2094, 2096, 2098, 2100, 2102, 2104, 2106, 2108, 2110, 2112, 2114, 2116, 2118, 2120, 2122, 2124, 2126, 2128, 2130, 2132, 2134, 2136, 2138, 2140, 2142, 2144, 2146, 2148, 2150, 2152, 2154, 2156, 2158, 2160, 2162, 2164, 2166, 2168, 2170, 2172, 2174, 2176, 2178, 2180, 2182, 2184, 2186, 2188, 2190, 2192, 2194, 2196, 2198, 2200, 2202, 2204, 2206, 2208, 2210, 2212, 2214, 2216, 2218, 2220, 2222, 2224, 2226, 2228, 2230, 2232, 2234, 2236, 2238, 2240, 2242, 2244, 2246, 2248, 2250, 2252, 2254, 2256, 2258, 2260, 2262, 2264, 2266, 2268, 2270, 2272, 2274, 2276, 2278, 2280, 2282, 2284, 2286, 2288, 2290, 2292, 2294, 2296, 2298, 2300, 2302, 2304, 2306, 2308, 2310, 2312, 2314, 2316, 2318, 2320, 2322, 2324, 2326, 2328, 2330, 2332, 2334, 2336, 2338, 2340, 2342, 2344, 2346, 2348, 2350, 2352, 2354, 2356, 2358, 2360, 2362, 2364, 2366, 2368, 2370, 2372, 2374, 2376, 2378, 2380, 2382, 2384, 2386, 2388, 2390, 2392, 2394, 2396, 2398, 2400, 2402, 2404, 2406, 2408, 2410, 2412, 2414, 2416, 2418, 2420, 2422, 2424, 2426, 2428, 2430, 2432, 2434, 2436, 2438, 2440, 2442, 2444, 2446, 2448, 2450, 2452, 2454, 2456, 2458, 2460, 2462, 2464, 2466, 2468, 2470, 2472, 2474, 2476, 2478, 2480, 2482, 2484, 2486, 2488, 2490, 2492, 2494, 2496, 2498, 2500, 2502, 2504, 2506, 2508, 2510, 2512, 2514, 2516, 2518, 2520, 2522, 2524, 2526, 2528, 2530, 2532, 2534, 2536, 2538, 2540, 2542, 2544, 2546, 2548, 2550, 2552, 2554, 2556, 2558, 2560, 2562, 2564, 2566, 2568, 2570, 2572, 2574, 2576, 2578, 2580, 2582, 2584, 2586, 2588, 2590, 2592, 2594, 2596, 2598, 2600, 2602, 2604, 2606, 2608, 2610, 2612, 2614, 2616, 2618, 2620, 2622, 2624, 2626, 2628, 2630, 2632, 2634, 2636, 2638, 2640, 2642, 2644, 2646, 2648, 2650, 2652, 2654, 2656, 2658, 2660, 2662, 2664, 2666, 2668, 2670, 2672, 2674, 2676, 2678, 2680, 2682, 2684, 2686, 2688, 2690, 2692, 2694, 2696, 2698, 2700, 2702, 2704, 2706, 2708, 2710, 2712, 2714, 2716, 2718, 2720, 2722, 2724, 2726, 2728, 2730, 2732, 2734, 2736, 2738, 2740, 2742, 2744, 2746, 2748, 2750, 2752, 2754, 2756, 2758, 2760, 2762, 2764, 2766, 2768, 2770, 2772, 2774, 2776, 2778, 2780, 2782, 2784, 2786, 2788, 2790, 2792, 2794, 2796, 2798, 2800, 2802, 2804, 2806, 2808, 2810, 2812, 2814, 2816, 2818, 2820, 2822, 2824, 2826, 2828, 2830, 2832, 2834, 2836, 2838, 2840, 2842, 2844, 2846, 2848, 2850, 2852, 2854, 2856, 2858, 2860, 2862, 2864, 2866, 2868, 2870, 2872, 2874, 2876, 2878, 2880, 2882, 2884, 2886, 2888, 2890, 2892, 2894, 2896, 2898, 2900, 2902, 2904, 2906, 2908, 2910, 2912, 2914, 2916, 2918, 2920, 2922, 2924, 2926, 2928, 2930, 2932, 2934, 2936, 2938, 2940, 2942, 2944, 2946, 2948, 2950, 2952, 2954, 2956, 2958, 2960, 2962, 2964, 2966, 2968, 2970, 2972, 2974, 2976, 2978, 2980, 2982, 2984, 2986, 2988, 2990, 2992, 2994, 2996, 2998, 3000, 3002, 3004, 3006, 3008, 3010, 3012, 3014, 3016, 3018, 3020, 3022, 3024, 3026, 3028, 3030, 3032, 3034, 3036, 3038, 3040, 3042, 3044, 3046, 3048, 3050, 3052, 3054, 3056, 3058, 3060, 3062, 3064, 3066, 3068, 3070, 3072, 3074, 3076, 3078, 3080, 3082, 3084, 3086, 3088, 3090, 3092, 3094, 3096, 3098, 3100, 3102, 3104, 3106, 3108, 3110, 3112, 3114, 3116, 3118, 3120, 3122, 3124, 3126, 3128, 3130, 3132, 3134, 3136, 3138, 3140, 3142, 3144, 3146, 3148, 3150, 3152, 3154, 3156, 3158, 3160, 3162, 3164, 3166, 3168, 3170, 3172, 3174, 3176, 3178, 3180, 3182, 3184, 3186, 3188, 3190, 3192, 3194, 3196, 3198, 3200, 3202, 3204, 3206, 3208, 3210, 3212, 3214, 3216, 3218, 3220, 3222, 3224, 3226, 3228, 3230, 3232, 3234, 3236, 3238, 3240, 3242, 3244, 3246, 3248, 3250, 3252, 3254, 3256, 3258, 3260, 3262, 3264, 3266, 3268, 3270, 3272, 3274, 3276, 3278, 3280, 3282, 3284, 3286, 3288, 3290, 3292, 3294, 3296, 3298, 3300, 3302, 3304, 3306, 3308, 3310, 3312, 3314, 3316, 3318, 3320, 3322, 3324, 3326, 3328, 3330, 3332, 3334, 3336, 3

“The Lexington men are three and one fourth inches long, in two and three eighths of an inch, and is three to five of an inch in diameter. The width, which is variable, is $\frac{1}{2}$ inch.”

[illegible]

*Manuals, Cyclopadia, etc.***Tidy, Charles Meymott, M.D., F.C.S.,**

Master of Surgery, Professor of Chemistry and of Forensic Medicine and Public Health at the London Hospital, Medical Officer of Health for Islington; late Deputy Medical Officer of Health and Public Analyst for the City of London, etc.

LEGAL MEDICINE. Volume I. With two colored plates. Contents: Evidence—The Signs of Death—Identity—The Causes of Death—The Post mortem. Sold by subscription only. See page 54.

LEGAL MEDICINE. Volume II. Contents: Expectation of Life—Presumption of Death and Survivorship—Heat and Cold—Burns—Ligaturing—Explosives—Starvation—Sex—Menstruities—Hermaphroditism. Sold by subscription only. See page 54.

LEGAL MEDICINE. Volume III. Contents: Legitimacy and Paternity—Pregnancy—Abortion—Rape, Indecent Exposure—Sodomy, Bestiality—Live Birth, Infanticide—Asphyxia, Drowning—Hanging, Strangulation—Suffocation. Sold by subscription only. See page 52.

Castle, Frederick A., M.D., New York.

WOOD'S HOUSEHOLD PRACTICE OF MEDICINE, HYGIENE, AND SURGERY. A Practical Treatise for the Use of Families, Travellers, Seamen, Miners, and others. By Various Authors. In two volumes, royal 8vo, 819 and 942 pages, illustrated by six hundred and thirty-five fine wood-engravings. Price, per volume, muslin, \$5.00; leather, \$6.00; half morocco, \$7.50. Sold by subscription only.

Ziemssen, H. Von, M.D., Munich.

CYCLOPEDIA OF THE PRACTICE OF MEDICINE. By Various Authors. Complete in twenty volumes, royal 8vo. Price per volume, in muslin, \$5.00; in leather, \$6.00; in half morocco, \$7.50. Sold by subscription only. See page 61.

Peabody, George L., M.D., New York.

SUPPLEMENT TO ZIEMSEN'S CYCLOPEDIA OF THE PRACTICE OF MEDICINE. By Various Authors. In one royal 8vo volume, 844 pages, bound to correspond. Price, in cloth, \$6.00; in leather, \$7.00; in half morocco, \$8.00; also extra muslin, not corresponding, \$6.00; and in red leather, \$7.00.

The aim of this work is to take up each subject treated of in Ziemssen's Cyclopadia, and to bring it down to date. Many of the articles will, therefore, embrace the progress of two years, but some, relating to the later volumes of Ziemssen, begin at a more recent period; the whole, however, forms a complete resumé of the progress of medicine mostly for the past five years.

Carpenter, Wesley M., M.D.,

Assistant Professor of Pathology, Lecturer in the Pathological Laboratory of the University of the City of New York; Secretary of the Medical Society of the County of New York; Secretary of the New York Pathological Society, etc.

AN INDEX OF THE PRACTICE OF MEDICINE. Sold only by subscription. See page 64.

Flint, Austin, M.D.

COMPENDIUM OF PERCUSSION AND AUSCULTATION AND OF THE PHYSICAL DIAGNOSIS OF DISEASES AFFECTING THE LUNG AND HEART. One volume, 18mo, 43 pages, muslin. Price, 50 cents.

Paul, Dr. Constantine,

Professor Agrégé to the Faculty of Medicine of Paris, etc.

DIAGNOSIS AND DISEASES OF THE HEART. Illustrated by numerous fine wood-engravings. Sold only by subscription. See page 52.

Practice, Rivers.

Loomis, Alfred L., M.D., LL.D.,

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–406

A TEXT BOOK OF PRACTICAL MEDICINE OF THE LUNGS.

£1,000 per year, distributed by two hundred and forty children, or £2 10s. per child, or £100,000 per year, or \$1,000,000.

The χ^2 test of goodness of fit is applied to compare the observed and expected frequencies and the results are given in Table 2. The arrangement of the data is such that the null hypothesis of no significant difference is based on a value of $\chi^2 = 0.00$ and $\text{d.f.} = 1$ ($\text{d.f.} = 2n - 1$) (Mantel, 1963; Neter et al., 1977, p. 484).

[illegible]

the usual field of integers, the usual field of integers is considered in textbooks on algebra. This is not the case; but it does so

With the exception of the *Journal of the American Medical Association*, which has been the only journal to publish a review of the book, the book has not been reviewed in the medical literature. The *Journal of the American Medical Association* reviewed the book in 1966, and the reviewer wrote that "this book is an excellent source of information for the student of medicine, and it is also a valuable reference work for the physician."¹ The book has been cited in the *Index Medicus*,² the *Journal of the American Medical Association*,³ and the *Journal of the American Osteopathic Association*.⁴ It is included in the *Journal of the American Medical Association* and the *Index Medicus*, and is also included in the *Journal of the American Osteopathic Association*. The book is a valuable reference work for the student of medicine, and it is also a valuable reference work for the physician.

11. **TUBES ON DISEASES OF THE RESPIRATORY ORGANS, HEART AND KIDNEYS.** One volume, 8vo, 501 pages. Price, bound, \$5.00; leather, \$6.00.

the \mathcal{H}^1 -in style, corresponds to arranging the \mathcal{H}^1 -style and partitioning its tree structure into \mathcal{H}^1 -style and \mathcal{H}^2 -style.

We can also systematically method in which
the elements are arranged, and noted from
left to right, very easily.— $C = C_1 + C_2 + \dots$

and the other, which is the ponderous look
of the last word is on the subject of
the author's not to write the author
of the press is to the press is to the press
of the press is to the press is to the press
of the press is to the press is to the press

¹⁰ We have also considered the possibility of a linear relationship between the number of species and the number of genera. However, the results of the regression analysis are not as good as those for the log-log relationship.

[illegible]

By following the Leontief work, we have

FIGURES ON FEVERS. One volume, 8v., 406 pages. Price, 50s. or \$4.00.

*Diseases of Throat and Chest, of the Rectum.***Corson, John W., M.D.,**

Late President of the Class of "Diseases of the Chest and Throat" in the New York and Eastern Dispensary; formerly Physician to the Brooklyn City Hospital; Physician to the Orange Marine Hospital, etc.

ON THE TREATMENT OF PLEURISY. With an Appendix of Cases, showing the Value of Combinations of Croton Oil, Ether, and Iodine, as Counter-irritants in other Diseases. One volume, 16mo, 31 pages, muslin. Price, 50c.

Ingals, E. Fletcher, A.M., M.D.,

Lecturer on Diseases of the Chest and Physical Diagnosis, and on Laryngology in the Postgraduate Course, Rush Medical College; Clinical Professor of Diseases of the Throat and Chest, Washington Medical College; Physician and Surgeon for Diseases of the Throat and Chest, Central Dispensary, Chicago.

LECTURES ON THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE CHEST, THROAT, AND NASAL CAVITIES. With one hundred and thirty-five Illustrations. One volume, 8vo, 437 pages, muslin. Price, \$4.00.

"It, bringing within one pair of book covers a complete epitome of affections of the chest and the entire respiratory tract, Dr. Ingals has recognized the interdependent relations of a group of anatomical regions falling together naturally, both in histology and pathology. His volume aims trenchly at an objective completeness which has not hitherto been attained, to our knowledge, in any single work of physical diagnosis, or on any disease of the chest and air-passages."—*American Journal of the Medical Sciences*.

"For good, practical, and correct teaching, this book has no superior. There is no volume which would be more useful in the general practitioner's daily life."—*American Medical*

Journal.

"This forms a valuable aid, both to the

student and practitioner, in the study of its subject. It is clear and concise in style, systematic and thorough in the consideration of each detail."—*Chicago Medical Journal and Review*.

"Both as to the matter and arrangement of the book, it must be of great service to the ordinary physician as well as to the specialist."—*Chicago Medical Monthly*.

"The work is of the most practical character; it avoids theoretical and unsatisfactory questions; the subjects are presented in a lucid and compact style. We predict for this work a wide field of usefulness, because it seems to us to be exactly adapted to the use of the physician in active practice."—*Medical Student Journal*.

Kelsey, Charles D., M.D.,

Chief Lecturer for Diseases of the Rectum and Anal, for Diseases of the Colon, in the Johns Hopkins Hospital and Dispensary for Women and Children, etc., etc.

THE PATHOLOGY, DIAGNOSIS, AND TREATMENT OF DISEASES OF THE RECTUM AND ANUS. One volume, 8vo, 430 pages, illustrated by two chromo-lithographic plates and many wood-engravings. Price, in cloth, \$4.00.

"The object of the author has been to make this book a text-book for the student and general practitioner, and to furnish that information which is so difficult to obtain without special advantages, such as are obtained by study of."

"In conclusion, we will add that the author has succeeded admirably in giving us a book which will greatly advance the education of the general practitioner, and a number of others who are heretofore shrunk from this district of the body, unsatisfactory branch of surgery, to essay it, and with renewed confidence."—*North American Medical Journal*.

"It forms the best recent work on a class of disease which, although claimed by surgery, has more real relation than most physical ones are aware of."—*The College and Clinical Record*.

"The work is an excellent one and will be fully appreciated."—*The Physician and the Surgeon's Association*.

"The student and busy practitioner will find here in a condensed form all that is practically known, and much that has not before been so widely known, of the diseases of the rectum and anus."—*The Medical Bulletin*, Philadelphia, Pa.

DISEASES OF THE RECTUM AND ANUS. Sold only by subscription. See page 51.

*Materia Medica, Therapeutics.***Fothergill, J. Milner, M.D.,**

M.D., F.R.C.S., Lecturer of the College of London; Senior Assistant-Physician to the City of London Hospital; Lecturer of the College of Victoria Park; The Assistant-Physician to the Westminster Hospital; Associate Lecturer of the College of Physicians, 1 Chancery Lane.

INDIGESTION, BILIOUSNESS, AND GOUT IN ITS PROTEAN ASPECTS.

Part I.—Indigestion and Biliousness. One volume, f2mo, 320 pages, mass.in. Price, \$7.25.

"The relation of digestion to habits of life, to the state of mind, and to the perfect nutrition of the body, is treated in a masterly manner, and bears the practical hints of the greatest possible utility to the practising physician."

"Altogether the work is a remarkably comprehensive study of a subject which is too little understood by the majority of medical men."—*Am. Jour. Med. Sci. & Surg.*

"Dr. Fothergill's writings always command attention, they are spirited and full of interesting facts, drawn mostly from his own long experience. This volume is written from a philosophical standpoint, and becomes a valuable treatise on natural digestion, by way of introduction to the main topic of the book. As the liver is the great store-house of supplies for the use of the system,

four chapters are devoted to its functions and their disturbances. In referring to the influence of mental strain and worry, Dr. Fothergill says: 'Talking one day with Mr. Van Abbott, whose biscuits for diabetes have such a well-deserved renown, I asked him, "Who are your diabetics mostly?" The reply was very significant: "Business men, comparatively old and gray for their years; men who look as if they had a deal on their minds." This was the response. It stands in suggestive relationship to the fact of neuro-diabetes being set up by shock or other mental perturbation, or of its artificial production by the denature of the floor of the fourth ventricle.' The whole book is practical and interesting reading."

Phillips, Charles D. F., M.D., F.R.C.S.E.,

F.R.C.S., M.D., Lecturer, Westminster Hospital, London.

MATERIA MEDICA AND THERAPEUTICS. Inorganic Substances. Adapted to

the United States Pharmacopoeia, by LAWRENCE JOHNSON, M.D. Volumes I. and II. Sent only by subscription. See page 64.

MATERIA MEDICA AND THERAPEUTICS. Vegetable Kingdom. Part I. and II.

Adapted to the U. S. Pharmacopoeia by HENRY G. PHILLIPS, A.M., M.D., Lecturer in Botany, University of the City of New York; Surgeon to the City of New York Hospital, etc. This practical book forms a volume in this series of 12.

Garrod, Alfred Baring, M.D., F.R.S.,

F.R.C.S., Lecturer, University of London.

THE ESSENTIALS OF MATERIA MEDICA AND THERAPEUTICS. One London Size, 8vo, 449 pages, extra mass.in. Price, \$1.00.

"The author has endeavored to present in 'The Essentials' a brief and systematic summary of the most important parts of the Materia Medica and Therapeutics. The work is divided into two parts, the first part being devoted to the study of the essential parts of the Materia Medica, and the second part to the study of the essential parts of the Therapeutics. The work is written in a clear and concise manner, and is adapted to the use of the student and the practitioner. It is a valuable addition to the literature of the subject."—*British Medical Journal*.

Trousseau, A., M.D.,

M.D., F.R.C.S., Lecturer, University of Paris; Professor, University of Paris.

REMARKABLE CASES. The Case of D. P. LINDEN, M.D., from the *Materia Medica et Therapeutica* of N. G. P. LINDEN, M.D., and J. L. LINDEN, M.D. Volume I. H. 8vo, 18 pages, 1881. Price, 50c.

"A volume of the Trousseau series, and is a most valuable addition to the literature of the subject. It is a most valuable addition to the literature of the subject. The work is written in a clear and concise manner, and is adapted to the use of the student and the practitioner. It is a valuable addition to the literature of the subject."—*British Medical Journal*.

*Therapeutics, Cholera, Diagnosis.***Binz, C., M.D.,**

Professor of Pharmacology in the University of Bonn.

THE ELEMENTS OF THERAPEUTICS. A Clinical Guide to the Administration of Medicines. Translated from the Fifth German Edition, by E. J. B. COOPER, M.D., in conformity with the British and American Pharmacopœia, 1898, by J. H. SPARKS, M.A., M.B., Oxon., Member of the Royal College of Physicians, London, Officier de Santé-Alpes Maritimes, France, formerly of the University of Bonn, Fellow. In one handsome 12mo volume, 345 pages, 8s. Published by Wm. Wood & Co., Ltd. Price, \$2.00.

"Fully up to the times as a therapeutic guide."—*Tribune Medical and Surgical Journal*.

"We are much mistaken, however, if the work in its present shape does not become a general favorite with both students and practitioners. It will also help, we think, to ad-

vance the movement to establish an American Pharmacopœia."—*N. Y. Herald*.

"The work appears to have been written with great care and bears the impress of originality, and is a volume that we should like to recommend in the strongest manner."—*The Medical Record*.

Wendt, Edmund C., M.D.,

Consultant, St. Francis' Hospital; Pathologist and Curator of the New York Infirmary for the Poor.

A TREATISE ON CHOLERA. Edited and Prepared in Association with JOHN C. PETERS, M.D., New York, JOHN B. HAMILTON, M.D., Surgeon General, U. S. Marine Hospital Service, and ELY McCLELLAN, M.D., Surgeon, U. S. Army. Illustrated with maps and engravings. Sold only by subscription. See p. 54.

"By availing themselves of the history and experience of cholera epidemics, to the present day, together with the new light thrown upon its mode of propagation, spread, and curability, the past year through the investigations of Prof. Koch and others—the knowledge of the most preventive measures, quarantine, etc., so recently acquired in France and Italy—the authors of this work aim to produce a book which may at least serve to prepare the physicians of America successfully to combat this dreaded scourge, should it unfortunately again be introduced into our country this year."

Burrall, F. A., M.D.,

ASIATIC CHOLERA. One volume, 12mo, 155 pages, fustlin. Price, \$1.50.

"It is a special merit of Dr. Burrall's timely volume, that it so states facts as to instruct the physician most impressively and acceptably in all that relates to preventive measures and prophylaxis. We are met with no writing on cholera in our language that has more happily

achieved this chief end of such a book. For this reason, no less than for the very excellence of the book, we are sure to commend the attention of all to this publication."—*Medical Record*.

Brown, Harvey E., M.D.,

Assistant Surgeon, United States Army.

REPORT ON QUARANTINE on the Southern and Central Coasts of the United States. One volume, 8vo, 145 pages, fustlin. Price, \$1.25.

Delafield, Francis, M.D.; and Stillman, Charles E., M.D.

A MANUAL OF PHYSICAL DIAGNOSIS. Illustrated by 100 original and 100 transparent lithographed plates. One volume, 8vo, 340 pages, fustlin. Price, \$2.00.

"The want of copiousness in the ordinary manuals on physical diagnosis, and the ever-unsatisfied student, and the never-ending trouble connected therewith. This work is an exception to this rule."—*Ohio Medical Record*.

"We are not aware of any other work of the present kind, so complete, so accurate, so up-to-date, and so well illustrated. It is a volume that we should like to recommend in the strongest manner."—*The Medical Record*.

Diagnosis.

Ranney, Ambrose L., A.M., M.D.,

Assistant Professor of Anatomy and Late Lecturer on the Surgical Diseases of the Genito-Urinary Organs at the Medical School of the Medical Department of the University of the City of New York; Late Surgeon of the North German Dispensary; Resident Fellow of the New York Academy of Medicine; Member of the Medical Society of the County of New York, etc.

A PRACTICAL TREATISE ON SURGICAL DIAGNOSIS, DESIGNED AS A MANUAL FOR PRACTITIONERS AND STUDENTS. Third Edition. One volume of 628 pages, illustrated by 31 plates, handsomely bound in muslin. Price, \$1.50.

"Useful account of its systematic arrangement."—*Chicago Medical Journal*.

"We are glad to see how more information can have been compassed in fewer words."—*Chicago Medical Journal* and *Practitioner*.

"The system and arrangement of the volume are commendable, and the author has written them out well."—*Southern Practitioner*.

"A very good and useful surgical diagnosis for the practitioner, student and beginner. As a text-book for students, it is quite valuable."—*St. Louis Medical Record*.

"With the exception of Macleod's 'Outlines,' published simultaneously in England

and in this country, in 1864, this is, so far as we know, the first monograph ever issued on surgical diagnosis."—*Philadelphia Medical Times*.

"The chief source of perplexity in the practice of medicine and surgery is to find out what is the matter with the patient. Uncomfortable, indeed, is the reflection of a practitioner when he has left a case bandaged and dressed for a *fracture*, when, perchance, it may be a *dislocation*." Dr. Ranney has given us a look to assist us in all such states of uncertainty, and he has done well; for in presenting the symptoms of disease in marked contrast, it makes the diagnosis of similar troubles really easy."—*P Toledo Med. and Surg. Journal*.

Guttman, Dr. Paul,

Professor of Medicine, University of Berlin.

A HANDBOOK OF PHYSICAL DIAGNOSIS; COMPRISING THE THROAT, THORAX, AND ABDOMEN. Translation from the Third German Edition, by ALLEN, Norman, M.D., Fellow of the Faculty of Physicians and Surgeons, Glasgow, and American Editor, with a colored plate and numerous illustrations. Second English Edition. 8vo. pp. 56.

Loomis, Alfred L., M.D.,

Professor of Medicine and Lecturer on the Medical Department of the University of the City of New York, and Surgeon of the Dispensary.

LESSONS IN PHYSICAL DIAGNOSIS. One volume, 8vo. 240 pages, illustrated. Price, \$1.00.

"The review of the text is clear and valuable. We are indebted to the author for the clear and concise manner in which he has presented the subject."—*Chicago Medical Journal*.

"We are glad to see how more information can have been compassed in fewer words."—*Chicago Medical Journal* and *Practitioner*.

"The system and arrangement of the volume are commendable, and the author has written them out well."—*Southern Practitioner*.

"A very good and useful surgical diagnosis for the practitioner, student and beginner. As a text-book for students, it is quite valuable."—*St. Louis Medical Record*.

"With the exception of Macleod's 'Outlines,' published simultaneously in England

and in this country, in 1864, this is, so far as we know, the first monograph ever issued on surgical diagnosis."—*Philadelphia Medical Times*.

"The chief source of perplexity in the practice of medicine and surgery is to find out what is the matter with the patient. Uncomfortable, indeed, is the reflection of a practitioner when he has left a case bandaged and dressed for a *fracture*, when, perchance, it may be a *dislocation*." Dr. Ranney has given us a look to assist us in all such states of uncertainty, and he has done well; for in presenting the symptoms of disease in marked contrast, it makes the diagnosis of similar troubles really easy."—*P Toledo Med. and Surg. Journal*.

"The review of the text is clear and valuable. We are indebted to the author for the clear and concise manner in which he has presented the subject."—*Chicago Medical Journal*.

"We are glad to see how more information can have been compassed in fewer words."—*Chicago Medical Journal* and *Practitioner*.

"The system and arrangement of the volume are commendable, and the author has written them out well."—*Southern Practitioner*.

"A very good and useful surgical diagnosis for the practitioner, student and beginner. As a text-book for students, it is quite valuable."—*St. Louis Medical Record*.

"With the exception of Macleod's 'Outlines,' published simultaneously in England

Botany, Climatology, and Physiology,

Johnson, Laurence, A.M., M.D.,

MEDICAL BOTANY: A Treatise on Plants used in Medicine. By J. E. S. *Scientifically colored plates and text arranged as the work of a physician.* 8 pp. 18. Description. See page 52.

Bell, A. N., A.M., M.D.,

1. *Dr. J. A. M. de Vries*, Minister of Agriculture, Ministry of Agriculture, 1000 CA Amsterdam, The Netherlands.
2. *Dr. J. A. M. de Vries*, Minister of the State of New Jersey, Division of Agriculture, 1000 Washington Avenue, Trenton, New Jersey 08646, U.S.A.
3. *Dr. J. A. M. de Vries*, Minister of the Department of Agriculture, 1000 Washington Avenue, Trenton, New Jersey 08646, U.S.A.

CLIMATOLOGY OF THE UNITED STATES AND ADJACENT COUNTRIES
and of such Foreign Ports and Places as have intimate Commerce with the United States, with special reference to Health Resorts, and the Protection of Public Health. Sold only by subscription. See page 54.

2. This work has been written expressly for Wood's Library, by one whose fathering of it
3. has been for many years in this line.

Fothergill, J. Milner, M.D.,

THE PHYSIOLOGICAL FACTOR IN DIAGNOSIS. By W. G. C. YONGE.
 Translated by S. G. B. Second edition. 8vo, 256 pages. Bound in paper. 8s. 7½d.

[illegible]

Ashby, Henry, M.D.,

MEMORANDA OF PHYSIOLOGY, TO WHICH ARE ADDED
THEORY OF VIBRATIONS, by an American, &c. &c. 1837. 12mo.
Price, \$1.00. Wason's Pocket Memoranda. 12mo.

Comstock, J. C.; and Comings, N., M.D.,

Physiology.

Kirkes' Handbook of Physiology.

HANDBOOK OF PHYSIOLOGY. By W. MORRANT BAKER, F.R.C.S., Surgeon to St. Bartholomew's Hospital and Consulting Surgeon to the Evelina Hospital for Sick Children, Lecturer on Physiology at St. Bartholomew's Hospital, and late Member of the Board of Examiners of the Royal College of Surgeons of England; and VINCENT DOHRMER HARRIS, M.D., Lond., Demonstrator of Physiology at St. Bartholomew's Hospital. Eleventh edition. In one volume, with coloured plates and five hundred illustrations. Price, muslin, £4.00; leather, £5.00.

Dr. Keil's Physiology has long enjoyed a high reputation, as one of the best and most practical works of its kind, and in this new edition, just completed by Drs. Baker and Harris, it is again published as could be presented to the practitioners of America.

of the *Philosophy* to the latest developments in the subject which it treats. The illustrations are well selected and will be found very useful to the student in his efforts to comprehend and master even the most intricate aspects of the subject." *The Harmonist*, Philadelphia, Pa., July, 1885.

There is a number of subjects and arrangements of subjects that might be the volume content of a psychology or a handbook, the absence of such hypotheses and argumentation on the part of the author is a real recommendation of the book. The illustrations are hardly be called good, but they are well drawn and the drawing itself is the old kind of a picture of the work on psychology. (*The 100th Anniversary of Psychology*, Ky., June, 1928, p. 10).

of the book before us is a revision and improvement of Kates' *Physiology*. It is a pleasant work to read. To those of us who have not had the opportunity to read the original of Kates it is book comes as a welcome addition to our library and a helpful reference to the student and ourselves. In the preface the author states which have been the sources of his information. We very cordially commend the book to the *Southwestern Medical Journal*, April, 1914.

[illegible]

"As a guide for the student, and ready reference for the practitioner, this work is not excelled by any other in the English language for the clearness of statement of established facts in the science of which it treats." *The Standard*, New York, April, 1885.

"The Messrs. Wood do not intend to forget the juniors and give them therefore the best works at a nominal price. There is hardly a practitioner of some years who did not learn the principles of physiology from Kirkes, and any work which reaches an eleventh edition, shows its intrinsic value."—*The North American Journal of Homoeopathy*, Philadelphia, May, 1885.

"Kirke's Handbook has been a popular text-book of physiology for so many years that its eleventh edition hardly calls for an extended notice, much less a review. It is enough to say that the editors, who are now practically the authors, have maintained the high character of the work, and have kept it fully up to the times in the science of physiology, which has made such vast progress within the last few years." *Continent's Medical Directory*, May, 1889.

"On carefully looking through the first edition of Kinkadee's Physiology, which is, however, really Kinkadee's, it seems to us, only that the courtesy of the editors, we can find scarcely anything to criticize. It seems thoroughly up to the time, and while most points in physiology have not been entirely omitted, they are stated in such a way as to render them easily comprehensible to the youngest student of medicine." *The Therapeutic Gazette*, May, 1886.

Lambert, T. S., M.D.,

PRIMARY SYSTEMATIC HUMAN PHYSIOLOGY, ANATOMY, AND HYGIENE. By J. A. WILSON, M.D., F.R.C.P., Lecturer in Physiology, University of London School of Medicine, and Consultant, both Simple and Compound Fractures, St. George's Hospital, London. Available for the use of young scholars. One volume, 8vo, pp. xiv + 607, illustrated by 19 plates. Price, 5s.

Brodie, Sir Benjamin, Bart., D.C.L.,

Physiology.

Harris, Vincent, M.D.,

M. = $\frac{1}{2} \pi$ and $\frac{3}{2} \pi$ and C = π and 3π . Phys. 1996. Demonstration of the complex plane. <http://www.phy.bme.hu/~mcs>

Power, D'Arcy, M.A., Oxon.,

MANUAL FOR THE PHYSIOLOGICAL LABORATORY. By C. C. L. S. 1914. 128 pp. 40 illustrations, in color. Price, \$1.50.

The paper of Little is voluminous, although of modest proportions and narrow in scope, is a carefully guided look to the student who desires to learn something of the rudiments of history, to the non-historian and history-ministry.¹¹ *—H. C. H.*
Edinburgh, Scotland, 1881.

"The programme given to laboratory work in a well equipped school sets a basis created in detail at the school as the voyage and a notice, which is in every way fitted to

serve the needs of any work or experiment in manipulative study, to furnish the student with the essentials of history, mathematics, astronomy, and physiology." (*Journal of the Assoc. of Voc. Teachers*, October 18, 1882.)

This little book is a collection of papers from the physiological laboratory. It is a very useful addition to the literature of the subject, and is well illustrated by the numerous examples of the work of the laboratory. It is published by the University of California Press, Berkeley, California, 1951.

Satterthwaite, Thomas E., M.D.,

A MANUAL OF HISTOLOGY, Edited and Prepared by FREDERICK STODOLSKY, M.D., of New York. In consultation with Drs. THOMAS D. BROWN, J. COLLIER WARREN, WILLIAM F. WHITNEY, CHARLES I. BRACE, and C. H. WATSON, of Boston; Dr. J. HENRY C. SARGENT, of Philadelphia; Dr. BENJAMIN F. WOODBURY, of Brooklyn; and Drs. EDWARD C. WOODB, ABRAHAM MAYOR, R. W. ARNOLD, A. R. RICHMOND, W. R. BRIDGES, D. BRYSON DEAN, C. L. DODGE, and W. H. PIERCE, of New York City. New Edition, with appendix, 1903. 8vo. 354 pages, previously 320 pp. and 316 pp. respectively. 1903. \$1.75.

It is a privilege to have a relatively well-known author and scholar, who is undoubtedly, as being a trustee of the American Veterinary Epitaphists Association, to be the author of the most recent issue of the *Journal of Veterinary Epitaphists*. In the last issue, the text was a tribute to the deceased, a review of the literature, and a list of the new members of the Association. I am pleased to see

[illegible]

Thudichum, J. L. W., M.D.,

A JOURNAL OF CHEMICAL PHYSICS, Vol. 1, No. 1, 1933
Published by the American Chemical Society, 527 P Street, N. W., Washington, D. C.

Microscopy.

Brocklesby, John, A.M.,

Professor of Mathematics and Natural Philosophy in Trinity College, Hartford.

THE AMATEUR MICROSCOPIST, or, Views of the Microscope. With a Manual of Microscopic Manipulation and Microscopic Objects. Illustrated with two hundred and forty-seven figures on wood and stone. One volume, 144 pages, muslin. Price, \$1.75.

"A little book full of curious and interesting facts regarding the microscopic world."—*New York Herald*.

"Published in very beautiful shape."—*London Mail*.

"It treats of the microscope, how to use it, and how to prepare microscopic objects for

examination."—*London Standard*. "Thoroughly revised, improved, and fully illustrated, and well calculated."—*London Herald*.

"The book is fine, a gotten up carefully, and forgetful to all teachers who desire to extend their knowledge into this most interesting domain."—*Kansas City Daily Star*.

Frey, Heinrich,

Professor of Microscopical Technology at Zurich.

THE MICROSCOPE AND MICROSCOPICAL TECHNOLOGY. A Text-book for Physicians and Students. Translated and Edited by GEO. R. CHAFFIN, M.D., Surgeon New York Eye and Ear Infirmary, Ophthalmic and Aural Surgeon to the St. Catherine and Williamsburg Hospitals, etc., etc. Illustrated with one hundred and eighty-eight engravings on wood. One volume, 8vo, 660 pages. Price, muslin, \$6.00; or cloth or leather, \$7.00.

"In many respects we think this the best work on the microscope."—*Medical Record of New York*.

"A complete exposition of the subject, practically indispensable to the practical microscopist."—*Chicago Medical Journal*.

"The work is presented very modestly, yet we find it not only very accurate in all its details of process, but complete regarding varieties of figures treated. The condensed style of the author, the richness of his matter, together with his unhesitating of histology, permit an abridgement of nearly all questions of microscopic anatomy, and many of obscure pathology. The solid silver stage and reflecting an instrument are especially valuable to one about to purchase."—*New York Journal of Medicine*.

"We cannot but be won, of all others, particularly fitted by its completeness and arrangement to serve the student, whether beginner or one far advanced. The best and most recent methods are here given in detail. The definitions of the author make this part of the work complete to the present time. Each tissue and organ is treated with a completeness limited only by the present progress of microscopic art. The translator and editor deserves the gratitude of the medical profession for placing before an English reading public Dr. Frey's work, rendered still more

valuable by his own additions, found in *Revue Scientifique, Technique et Scientifique, Practical Microscopy*.

"Those who are familiar with Frey's admirable manual will be gratified to find that this work is very readable, and that it enables our American and English students who are unacquainted with the German language to participate in the instruction of the great Zurich professor. This abridgement of Frey's treatise possesses an especial value. As a beginner, observer, or recorder of microscopic matter in which are described the most improved methods and means of studying various structures in their healthy and morbid conditions. To sum up all, we have a neat, handy, and valuable volume in our hands. The microscopist cannot afford to be without it."—*Philadelphia Medical Journal*.

"We advise all chemists and students of the microscope to possess Frey's *Microscopical Technology*."—*Boston Medical and Surgical Journal*.

"It is a peculiar, indeed, a rare, contribution of the profession to the literature of the work. With three or four exceptions, and the expert possessors of the work are not for the purpose of the book, but for the instruction of the student."—*London Standard*.

Carpenter, Wm. B., C.B., M.D., LL.D.

THE MICROSCOPE AND ITS REVELATIONS. Six Parts.

Volume I. Illustrated by one colored and twenty-six pictorial plates, and one hundred and two fine wood engravings.

Volume II. Illustrated with twenty-eight pictorial plates, and one hundred and two wood engravings. Sold only by subscription. See page 34.

*Surgery.***Holmes' System of Surgery.**

A SYSTEM OF SURGERY, THEORETICAL AND PRACTICAL, IN TREATISES OF VARIOUS AUTHORS. Edited by T. Holmes, M.D., Surgeon-in-Chief of the Army, St. George's Hospital, and Surgeon-in-Chief of the Royal College of Physicians. Third Edition, thoroughly revised. Three volumes, containing 2,000 pages. With numerous woodcuts, lithographs, and chromo-lithographs. The cloth-bound set, \$21.00; in leather, \$24.00; in half leather, \$17.00.

It is fully up to the times, and forms a "What ever may be the belief in the efficacy of antiseptics, it will be in practice progressively strong as to the treatment of the various principles of cleanliness, of asepsis, of isolation, and the management of the various forms of these principles are of the highest importance, and the different parts are made to be very judiciously presented."—*The Medical Record*.

The work as a whole, claiming as it does to represent the English school of surgery, is complete in every respect, and really stands without a rival. As it stands, it represents the original and latest Holmes' System of Surgery, and contains the most recent views of the many distinguished authors who first contributed to its pages."—*The Medical Record*.

Clarke, W. Fairlie, M.A. and M.B., Oxon., F.R.C.S.,

Assistant Surgeon, Charing Cross Hospital.

A MANUAL OF SURGERY. A new edition, thoroughly revised, with numerous illustrations by an American surgeon. Nearly two hundred lithographs. 300 pages. Sold only by subscription. See page 57.

Wyeth, John A., M.D., Univ. of Louisville,

Medical Director, New York City; M.D. Schools of N. Y. City, Albany, and Philadelphia; Honorary Member of the College of Physicians and Surgeons of the University of Maryland.

ESSAYS IN SURGICAL ANATOMY AND SURGERY. One volume, 8vo., 292 pages, illustrated, muslin. Price, \$2.00.

A HANDBOOK OF MEDICAL AND SURGICAL REFERENCE. One volume, 8vo., 379 pages, muslin. Price, \$1.25; trucks, \$1.50.

Bauer, Louis, M.D., M.R.C.S., Eng.,

Professor of Pathology and Clinical Surgery, Lecturer at the New York State Medical School, Medical Director, New York Pathological Society, of the American Medical Association, and Lecturer at the London Medical Society, Eng.

LECTURES ON ORTHOPEDIC SURGERY. Delivered at the Brooklyn Medical and Surgical Institute. Revised and augmented. One volume, 8vo., 339 pages, illustrated, muslin. Price, \$3.25.

"We are especially pleased with the chapters on diseases of the spine and joints which occupy a large portion of the book. Dr. Bauer has given us opportunity of acquainting him- self with the resources of information on the subject, and his work will be found as nearly complete as the present advanced condition of our knowledge admits of."—*The Medical Record*.

"As a treatise on orthopedics we have very few words to commend it, and it is one of the most exhaustive and complete of its class in the literature."—*St. Louis Medical Review*.

Carnochan, John Murray, M.D.,

Lecturer in the University of Edinburgh, and Pathologist, Royal Infirmary.

A TREATISE OF THE ETIOLOGY, PATHOLOGY, AND TREATMENT OF CONGENITAL DISLOCATIONS OF THE HEAD OF FEMUR. Edited and published by the author. 25 lithographed plates. One volume, 8vo., 275 pages, muslin. Price, \$2.00.

"By far the most complete and systematic work in the English language and the only one which contains any directions for treatment."—*British Association Medical and Physiological Journal*.

"We are especially pleased to have a treatise on this subject, the only one in the English literature. As a treatise on the subject of congenital dislocations of the head of femur, it is one of the most complete and exhaustive of its class in the literature."—*St. Louis Medical Review*.

*Surgery.***Poulet, Alfred, M.D.,**

Assistant Surgeon-Major, Inspector of the School for Military Medicine at Val-de-Grace.

A TREATISE ON FOREIGN BODIES IN SURGICAL PRACTICE. Illustrated by original wood-engravings. Translated from the French. Volumes I. and II. Sent only by subscription. See page 56.

Taylor, Charles Fayette, M.D.,

Surgeon of the New York Dispensary, and of the Dispensary and Hospital; Fellow of the New York Academy of Medicine, Member of the New York County Medical Society, and Permanent Member of the American Medical Association.

ON THE MECHANICAL TREATMENT OF DISEASE OF THE HIP-JOINT. One volume, 8vo, 62 pages, illustrated, muslin. Price, \$1.25.

Peugnet, Eugene, M.D.,

Surgeon of the New York Dispensary, Member of the New York Pathological Society, of the Medical Society of the City of New York, and of the Medical Society of the County of New York, etc., etc.

THE NATURE OF GUNSHOT WOUNDS OF THE ABDOMEN, AND THEIR TREATMENT: Based on a Review of the Case of the late James Fisk, Jr., in its Medicolegal Aspects. One volume, 8vo, 96 pages, muslin. Price, \$1.25.

Pilcher, Lewis S., A.M., M.D.,

of New York, N.Y.

THE TREATMENT OF WOUNDS. Being a Treatise on the Principles upon which the Treatment of Wounds should be Founded, and on the Best Methods of carrying them into Practice, including a Consideration of the Modifications which Special Injuries may demand. Illustrated by wood-engravings.

Salter, S. James A., M.B., F.R.S.,

M.D., F.R.C.S., Royal College of Surgeons in London; Examiner in Dental Surgery at the College of Dental Surgeons, London, 1884.

DENTAL PATHOLOGY AND SURGERY. One volume, 8vo, 339 pages, illustrated, muslin. Price, \$1.50.

Mr. Salter was educated as a surgeon, and was House Surgeon at King's College Hospital, and practised surgery for the first few years of his professional career; but it has given to the profession a valuable treatise, not only on the pathology of the teeth, but also on dental surgery. In conclusion, we would remark, that not only as a most able and practical surgeon, but also as a writer and pathologist, Dr. Salter's treatise on dental subjects is a valuable treatise, and should be read by every one interested in dental or dental operations. The book is well printed, and is illustrated with many fine illustrations, and is a most valuable work. *Medical and Dental Review.*

"This is a most valuable work, and is a most valuable work, and is a most valuable work."

long been known as one of the most scientific dentists of the day. Mr. Salter had the great advantages of such a complete medical education as is implied by the possession of a degree of the University of London, upon which he turned his attention to dental subjects, and he has continued his labors upon the broad basis thus laid down, with the satisfactory results that he has contributed not a little to both the science and practice of dentistry, the latter in its widest sense. We close Mr. Salter's work well satisfied that it is an honest record of good physiological and practical work, and we congratulate both the surgical and dental professions on possessing such a valuable work of reference."—*London Lancet.*

Goddard, Paul B., M.D., M.R.N.S., M.R.P.S.,

Professor of Anatomy, and Lecturer on the Anatomy of the Human Teeth, at the University of London.

THE ANATOMY, PHYSIOLOGY, AND PATHOLOGY OF THE HUMAN TEETH. With the Most Improved Methods of Treatment, including Operations, and the Methods of Making and Setting Artificial Teeth. With thirty plates. Aided in the preparation by Joseph L. PARKIN, Dentist. One volume, 4to, 227 pages, illustrated. Price, \$6.75.

*Surgery, Diseases of Women.***Owen, Professor Richard,**

THE PRINCIPAL FORMS OF THE SKELETON AND THE FIBRO-CARTILAGINOUS SYSTEM OF NATURAL HISTORY AND COMPARATIVE ANATOMY.—One volume, 64 pages, illustrated, cloth. Price, 75c.

The International Encyclopædia of Surgery.

A Systematic Treatise on the Theory and Practice of Surgery, by the various nations. Edited by JOHN ASHURST, JR., M.D., Professor of Surgery in the University of Pennsylvania. This encyclopædia is illustrated with chromolithographs and woodcut engravings. Published in Russian, \$6.00; leather, \$7.00; half in Russia \$8.00. See page 61.

Mundé, Paul F., M.D.,

Professor of Obstetrics in the New York Polyclinic, and Surgeon-in-Chief of the New York Hospital, and of the Maternity Hospital, New York. Author of *The American Gynaecologist's Manual*.

A TEXT-BOOK OF MINOR SURGICAL GYNECOLOGY. One volume, containing nearly 600 pages, illustrated with over three hundred engravings and photographs. Price, \$5.00.

Dr. Mundé's Manual, which appeared as one of the volumes of the *Library of Standard Medical Authors*, met with success in the market, and the success of the publication arranged with the distinguished author that the preparation of a new and enlarged work which would be based upon the previous one, and incorporate with the new material all the new matter as would be necessary to keep pace with the advance of the science, and the requirements of a book suitable for teaching purposes.

The volume here announced is the result, and the publishers confidently expect that it will support popularity in its field. To such an extent, however, with the publishers, that this new work is an expansion of the scope and number of facts that it is necessary to treat of those minor conditions and maladies commonly encountered in the treatment of diseases of women. As the scope of the work was enlarged to include facts, the logical sequel is not permit the fundamental facts to be omitted, and the student and practitioner should know, and is obliged to know, the facts of the science in the course of his practicing his work, while it is not supposed to apply to the more general principles of the body or operating table. While attempting to give the facts, and clear and concise description of details and manipulations, the requirements of a work of extensive value, the book contains but a minimum of comments. The practical statements of the methods and operations and the detailed facts in description will render the book of the value of those giving especial attention to the treatment of diseases of women, and it should be of general practical use, where it is to be employed as a text-book of the specialty.

Duncan, J. Mathews, A.M., M.D., etc.,

Professor of Obstetrics in the University of Pennsylvania, and Surgeon-in-Chief of the Maternity Hospital, Philadelphia. Author of *Practical Treatise on Perimetritis and Parametritis*, and *On the Mortality of Childbed and Maternity Hospitals*.

ON THE MORTALITY OF CHILDBED AND MATERNITY HOSPITALS.—One volume, 8vo, 172 pages, paper. Price, \$2.50.

A PRACTICAL TREATISE ON PERIMETRITIS AND PARAMETRITIS.—One volume, 12mo, 240 pages, paper. Price, \$2.00.

These books have been long recognized as the best of their kind, and are the result of the author's observations and experience. They are now being reissued, and are being observed, mark the present, as Dr.

Partridge, Edward L., M.D.,

New York, etc.

THE OBSTETRICAL REMEMBRANCER.—A Practical Handbook. Published in English, illustrated with engravings, by the author, J. W. Partridge, M.D., New York. Price, \$1.00.

*Diseases of Women.***Sims, J. Marion, A.B., M.D.,**

Professor of Obstetrics, H. P. S. School of New York; Fellow of the New York Academy of Medicine; of the New York Pathological Society; of the New York Historical Society; of the New York State Academy of Medicine; of the Royal Medical and Chirurgical Society, London; of the London Medical Association; of the Pathological Society, London; Fellow of the Obstetrical Society, London; Honorary Member of the Académie des Sciences et des Lettres, Paris; Hon. Fellow of the Imperial Academy of Medicine, St. Petersburg; and President of the Legion of Honour, etc.

CLINICAL NOTES ON UTERINE SURGERY. With Special Reference to the Management of the Sterile Condition. One volume, 8vo, 401 pages, illustrated, paper, Springfield, Mass. Price, \$1.00.

"It is a grand, far-reaching, persevering effort, the result of a long and arduous action, and a noble example of the way to the development of a specialty that has been a stumbling-block to the general profession. We commend this work to our readers."—*New Orleans Medical and Surgical Journal*.

"The volume should be in the hands of

every practitioner, young and old. While its teachings are so simple that the merest tyro can fully comprehend them, they are replete with valuable lessons to the physician of ripe experience. We have perused the work with much satisfaction, arising from it refreshed rather than satiated."—*Pacific Medical and Surgical Journal*.

Skene, Alexander, J. C., M.D.,

Professor of Obstetrics, Westminster Hospital, and College Hospital; Fellow of the Anatomical Society of Great Britain; Corresponding Member of the Gynecological Society of the County of Kings; and of the Society of New York.

DISEASES OF THE BLADDER AND URETHRA IN WOMEN. One volume, 8vo, 314 pages, illustrated, muslin. Price, \$3.00.

"It is a subject of the sterling practical matter which this work abounds, we have the author's personal observations admirably expounded, particularly in the chapter on urinary analysis."—*Pacific Medical Journal*.

"This work of the distinguished gynecologist, Skene, may be in medical literature. It is undoubtedly the only systematic treatise on this subject in the English language, and it is a treatise that cannot but be laden with the fruits of his long and able country."—*Pacific Medical Journal*.

"Dr. Skene has rendered the profession a real service, while the meagre information on this subject heretofore within the reach of busy practitioners will command for this book the place it should fill in every library."—*St. Louis Medical and Surgical Journal*.

"These lectures are exactly what the author claims for them—a convenient, plain, uncomplicated statement of the principal diseases of the female urethra and bladder."—*Detroit Journal*.

Bedford, Gunning S., A.M., M.D.,

Professor of Obstetrics, H. P. S. School of New York; Lecturer on Diseases of Women and Children, and Clinical Obstetrics, in the H. P. S. School of New York; and of the Clinical Lectures on the Diseases of Women and Children.

THE PRINCIPLES AND PRACTICE OF OBSTETRICS. Carefully revised and completely re-issued. One volume, 8vo, 763 pages, illustrated by four colored plates and ninety-nine wood-engravings. Price, muslin, \$6.00; leather, \$6.50.

"As a text-book, it is a truly excellent one. It is systematic in its arrangement, clear in its statements, and in its teachings."—*American Journal of the Medical Sciences*.

CLINICAL LECTURES ON THE DISEASES OF WOMEN AND CHILDREN. One volume, 8vo, 667 pages. Price, muslin, \$1.00; leather, \$5.00.

"This is a most valuable and useful work. We congratulate the author upon the high compliment paid to his labors in the publication of this work, the new field of uterine pathology, where so many a physician has been vainly seeking for reputation."—*American Journal of the Medical Sciences*.

Garrigues, Henry Jacques, A.M., M.D.,

Professor of Obstetrics, H. P. S. School of New York; Lecturer on Diseases of Women and Children, in the H. P. S. School of New York; and of the Clinical Lectures on the Diseases of Women and Children.

DIAGNOSIS OF OVARIAN CYSTS BY MEANS OF THE EXAMINATION OF THEIR CONTENTS. One volume, 8vo, 112 pages, illustrated, muslin. Price, \$6.25.

*Diseases of Women.***Verrier, E., M.D.**

PRACTICAL MANUAL OF OBSTETRICS. Fourth edition, revised and revised with four Obstetric Tables of Professor Pajot. First American edition, with revision and Annotations by Edward L. Fairbridge, M.D., Professor of Obstetrics in the New York Post-Graduate Medical School. One volume, 8vo, 420 pages. Illustrated by one hundred and two wood engravings. Price, \$1.50. Sold by subscription only. See page 52.

Braun, Dr. Carl R.,

Professor of Midwifery, Vienna.

THE URÆMIC CONVULSIONS OF PREGNANCY, PARTURITION, AND CHILDBED. Translated from the German, with notes, by J. MATTHEWS DUNCAN, F.R.C.P.E., Lecturer on Midwifery, etc. One volume, 12mo, 182 pages, muslin. Price, \$1.00.

"It contains, in a condensed form, the most complete and readable history of this affection yet published."—*New York Journal of Medicine*.

"We advise all who feel interested in the subject to procure it, as it will fully repay the

purchase."—*St. Louis Medical and Surgical Journal*.

"A most valuable essay, and one that will not be easily recalled for its completeness and erudition."—*Dublin Medical Press*.

Byford, William H., A.M., M.D.,

Professor of Obstetrics and Diseases of Women at the University of Maryland College of Physicians, author of "The Practice of Midwifery and Surgery, and of the Diseases and Accidents of Pregnancy, Women," "Chronic Inflammation of the Uterus," and "Gynecology."

A TREATISE ON THE THEORY AND PRACTICE OF OBSTETRICS. Illustrated with one hundred and fifty wood engravings. One volume, 8vo, 490 pages, muslin. Price, \$3.75.

"Professor Byford has been long and favorably known to the professional public by his numerous communications to the medical press, his previously published elaborate books, and by his widely extended private and consulting practice."—"Professor Byford's book is such up to the times, and a successful exposition of the subject."—*Chicago Medical Journal*.

"Byford's Obstetrics affords the student and practitioner the science and practice of the

art in the most available form. It is compact, though not large; it is full and perfect, and still is compressed into comparative small space. It contains what is known, and recommends itself to the profession and especially to medical students, by its clear, well-considered, complete teachings. Everything that can be said in favor of any work on this subject can be said of it."—*Buffalo Medical Journal*.

Klob, Julius M., M.D.,

Professor of Gynecology, Vienna.

PATHOLOGICAL ANATOMY OF THE FEMALE SEXUAL ORGANS. Translated from the German by JOSEPH KAMMERER, M.D., Physician to the German Hospital and Dispensary, New York, and BENJAMIN T. DAWSON, M.D., Assistant to the Chief of Obstetrics in the College of Physicians and Surgeons, New York. One volume, 8vo, 200 pages, muslin. Price, \$1.50.

Chapman, Edwin Nesbit, M.D.,

Lecturer on Gynecology, University of Wisconsin, and Lecturer on Diseases of Women, University of Michigan.

HYSTEROLOGY: A Treatise on Diseases of the Uterus and its Appendages, and Displacements of the Uterus. Illustrated with 100 plates, many of them colored. One volume, 8vo, 604 pages, muslin. Price, \$4.50.

"His book is well worth reading. It is one of the best on the subject that has yet been published."—*Lancet Medical Press*. "A most valuable and instructive treatise on the diseases of the female sex."—*Register and Times*.

*Diseases of Women.***Emmet, Thomas Addis, M.D.,**

Surgeon-in-Chief of the New York State Woman's Hospital, etc., etc.

VESICO-VAGINAL FISTULA FROM PARTURITION AND OTHER CAUSES;
with Cases of Recto-vaginal Fistula. Illustrated with wood-engravings. One
volume, 8vo, 250 pages, muslin. Price, \$2.75.

"A useful and painstaking record of many cases of vesico-vaginal fistula, arising from all sorts of causes. The operations necessary in each case are clearly described."—*Medical Record*, N. Y.

"Certainly, no one is more competent to write on this subject in the matter than Dr. Emmet, for his experience has been great indeed."—*Canadian Medical Repository*.

"No work of its size has so much enriched the literature of gynecology as this one."—*Medical Record*.

"As to the *physique* of the book, if we have any fault to find, it is with the elegance of its workmanship and costume, in which respect it puts to blush its shelf companions."—*Pacific Medical and Surgical Journal*.

Brown, W. Symington, M.D.,

Member of the College of Surgeons, Society of Boston; Fellow of the Massachusetts Medical Society, etc.

A CLINICAL HANDBOOK ON THE DISEASES OF WOMEN. Illustrated with
wood-engravings. One volume, 8vo, 247 pages, muslin. Price, \$2.50.

"The author writes with great simplicity a master of his subject, and chooses, off-hand, the shortest route to the understanding of his subject, and ready clinical lecturer who is his hearers."—*Louisville Medical News*.

Tilt, Edward John, M.D.

A HANDBOOK OF UTERINE THERAPEUTICS AND DISEASES OF WOMEN.
Fourth Edition. Sold only by subscription. See page 55.

Tait, Lawson, M.D.

DISEASES OF WOMEN. A new edition, with considerable additions, prepared by
the Author expressly for Wood's Library. This very compact, useful book makes
a volume of 204 pages, with illustrations. Sold only by subscription. See
page 57.

Fritsch, Heinrich, M.D.,

Professor of Obstetrics and Gynecology at the University of Halle.

THE DISEASES OF WOMEN. A Manual for Physicians and Students. Trans-
lated by J. L. Simon, F.R.S. Illustrated with one hundred and fifty fine wood-
cut engravings. Sold only by subscription. See page 53.

Savage, Henry, M.D.,

Surgeon-General, and Surgeon-in-Chief of England, one of the Consulting Medical Officers of the Sa-
vory Hospital, etc., etc.

**THE SURGERY, SURGICAL PATHOLOGY, AND SURGICAL ANATOMY OF
THE FEMALE PELVIC ORGANS,** in a Series of Plates, taken from Nature,
with Comments, Notes, and Cases. Third Edition, revised and greatly ex-
tended. Sold only by subscription. See page 56.

Hart, D. Berry, M.D.,

Professor of Midwifery, and Lecturer on the Women's School of Medicine, Edinburgh, etc., etc.; and

Barbour, A. H., M.D.,

Professor of Obstetrics, and Lecturer on the Women's School of Medicine, etc., etc.

MANUAL OF GYNECOLOGY. A Manual. Illustrated with eight plates, two of
which are colored, and one hundred and ninety-two fine wood-engravings.
Sold only by subscription. See page 54.

MANUAL OF GYNECOLOGY. A Manual. Illustrated with eight plates, and two hundred and nine fine
wood-engravings. Sold only by subscription. See page 53.

Diseases of Children, of the Eye, Anatomy.

Henoch, Dr. Edward,

Professor of the Pathology and Medicine of Children in the Royal Charité Hospital, and Professor in the same University.

LECTURES ON DISEASES OF CHILDREN. A Handbook for Physicians and Students. Translated from the German. Sold only by subscription. See page 54.

Routh, C. H. F., M.D.

INFANT FEEDING, AND ITS INFLUENCE ON LIFE; or, The Causes and Prevention of Infant Mortality. Third Edition. This unique work forms a volume of 286 pages in Wood's Library. Sold only by subscription. See page 57.

Dwight, Thomas, A.M., M.D.,

Professor of Anatomy and Histology in Harvard University; Fellow of the American Academy of Arts and Sciences; Surgeon at the Carney Hospital.

FROZEN SECTIONS OF A CHILD. Fifteen full-page lithographic plates, drawings from nature by H. P. QUINCY, M.D. One volume, royal 8vo, 66 pages, muslin. Price, \$5.00.

This book possesses the great merit now almost unknown. The plates are pen drawings from nature of the body of a girl said to be three years old, and are beautifully and vigorously executed. They possess peculiar attractions to the anatomist from the fact, that in the picture, that at three years of age, the proportions of the body and of the or-

gans are no longer those of an infant, and not yet those of an adult. The sections from which these drawings are made begin at the neck, and continue, about an inch apart, through the trunk. For the student and for the practitioner this book will prove interesting and practically useful."

Mauthner, Ludwig,

Professor in the University of Vienna.

THE SYMPATHETIC DISEASES OF THE EYE. Translated from the German by WARREN WHEELER, M.D., Surgeon United States Army, and JAMES A. SPALDING, M.D., Member of the American Ophthalmological Society; Ophthalmic Surgeon to the Marine General Hospital. One volume, 12mo, 220 pages, muslin. Price, \$2.00.

This treatise is one of the best parts of this series. It is written in a clear, concise, and reliable manner, and is one of the best works on the subject of sympathetic diseases of the eye that has appeared.

Much of the treatment of these diseases must be done by the physician, and the factors in the treatment are the same in all cases of the disease, and the treatment is the same in all cases.

reliable description of the multifarious symptoms, and the treatment of sympathetic ophthalmia, so that they may at once recognize its presence, and treat it from the outset appropriately and effectually. Although cases of this nature are comparatively rare, their importance is sufficiently great to account for the appearance of this excellent work in an English version."

Noyes, Henry D., M.D.,

Professor of Ophthalmology in the University of Michigan.

Professor of Medicine and Clinical Surgery in the University of Michigan.

DISEASES OF THE EYE. Illustrated by a series of chromo-lithographs and numerous drawings. Sold only by subscription. See page 55.

This treatise is one of the best parts of this series. It is written in a clear, concise, and reliable manner, and is one of the best works on the subject of diseases of the eye that has appeared.

Foote, John, M.D.,

Professor of Ophthalmology in the University of Michigan.

MEMORANDA RESPECTING THOSE DISEASES OF THE EYE WHICH ARE MORE FREQUENTLY MET WITH IN PRACTICE. One volume, 8vo, 100 pages, muslin. Price, 25 cents.

Diseases of the Ear, Throat and Nose.

Roosa, D. B. St. John, M.D.,

[illegible]

A PRACTICAL TREATISE ON THE DISEASES OF THE EAR, INCLUDING THE ANATOMY OF THE ORGAN. Sixth Edition. One vol., 8vo., 314 pages. Illustrated by 140 woodcut engravings and one chromo-lithograph. Paper, 45s.6d.; leather, 55s.6d.

If any one has the right to speak a tentative opinion on theological matters, it is Dr. Roosevelt; for it is he who, more than any other, has made known to the American medical profession the general principles of treatment of mental diseases; and it is safe to say that no book upon any special subject has not more widely distributed throughout the country than has his admirable treatise. If this work be carefully studied, it will be seen that two or three lines run through the whole of what has been written: first, that a skilled

treatment of our troubles involved a complete knowledge of general principles, so that the measure of success is determined by the thoroughness with which the practitioner follows the examination of his cases, before applying his simple, scientific principle.

"It is one of the maxims of the medical profession, that it is to be done without a knowledge of what is required, and that practice in the dark is better than no practice at all. It is less justifiable in the case of medicine than in any other."—*The Medical Record*.

Bosworth Franke Huntington, M.D.,

[illegible]

A MANUAL OF DISEASES OF THE THROAT AND NOSE. One volume, octavo, 418 pages. Illustrated with wood engravings. Muslin. Price, \$3.25.

"The artist has done his work well, and in clear and expressive language gives the reader a new, but not small experience." The work, like most emanating from the other side of the Atlantic, is well written, intelligent, and would not be considered "excellent." *Deliberate Heart* is superb!

As I work on this subject, I intend more especially to bring into practical use the following suggestions, which are wanted and desired. It is worth being fortunate in the country where we live has more talent in the subject. His handling of it, moreover, has been so good and judicious, as I stated when examining a few of his books, as sufficient to sustain it.

M. Henry, Medical Arts.

The book is an excellent introduction to the thinking, and Dr Be worth has been particularly helpful in preparing the writing of "The Philosophy of Mathematics."

The author's goal is to make this book as instructive as possible, without being too mathematical, so that the student, both in the classroom and on his own, will find it as well as *Learning to Read*. *Richard D. Woods*

ATTs are an excellent tool for exposing students to a text, which is an excellent opportunity to provide a more in-depth analysis of a historical document. Students can be asked to write a response to a document, or to be assigned a document to read and then to be asked to write a response to it.

Van Troeltsche, A.

DISEASES OF THE EAR IN CHILDREN. Price, \$1.50.

Salter, Henry Hyde, M.D.,

ON ASTHMA: ITS PATHOLOGY AND TREATMENT 541

Secé, Germain.

DISCUSSION OF THE LUNGS OF A SPECIFIC NON-TUBERCULOUS NATURAL
 A. C. BROWN, JR., D.D., Chief, Pneumology, General Hospital, State of Connecticut, Hartford, Conn.
 and the Lungs of the Tuberculous. The subject of L. P. HUBBARD, M.D., Assistant Surgeon, Medical
 Corps, United States Army, Philadelphia, Pa., and the Lungs of the Tuberculous. The subject of
 SOCIETY. One of the 14. The subject of A. C. BROWN, JR., D.D., Chief, Pneumology, General
 Hospital, State of Connecticut, Hartford, Conn., and the Lungs of the Tuberculous. The subject of
 L. P. HUBBARD, M.D., Assistant Surgeon, Medical Corps, United States Army, Philadelphia, Pa., and the Lungs of the Tuberculous. The subject of

*Diseases of the Nasal Cavities, Larynx, etc.***Semeleder, Dr. Friedrich.**

Professor of Medicine, University of Mexico; Member of the Royal Medical Society of Montpellier in Paris; Formerly Member of the Medical Faculty of the Branch Hospital at Gumpendorf.

RHINOLOGY AND LARYNGOSCOPY: THEIR VALUE IN PRACTICAL MEDICINE. Translated from the German by EDWARD T. CASWELL, M.D. With 100 illustrations, including 10 graphic plates. One volume, octavo, 194 pages. Price, \$3.75.

"This is a most interesting and useful work, of the professional public to whom it comes."
New York Medical Journal.
 "A most valuable and practical work."
New York Medical Journal.

Mackenzie, Morell, M.D., London.

DISEASES OF THE PHARYNX, LARYNX, AND TRACHEA. Illustrated by 112 colored plates. Sold only by subscription. See page 56.

Robinson, Beverley, A.M., M.D. (Paris).

Professor of Medicine, Bellevue Hospital Medical College, New York; Physician to St. Vincent's Hospital, New York.

A PRACTICAL TREATISE ON NASAL CATARRH. One volume, 8vo, illustrated. Price, \$2.50.

"Among the most recent contributions to the literature of the nose, this seems to me to be the best. It is written in a style which is well written, concise, clear, and to the point. The author presents, in good readable style, his opinions as to the diagnosis and treatment of this stubborn and disheartening disease. Aside from the more distinctly professional suggestions, he gives excellent common sense advice on regard to matters not usually dwelt upon in treatises on disease of the nasal cavity." *American Specialist.*

James, Prosser, M.D.

LARYNGOSCOPY AND RHINOSCOPY IN DIAGNOSIS AND TREATMENT OF DISEASES OF THE THROAT AND NOSE. Fourth edition, enlarged, one volume, 8vo, 323 pages. Illustrated with wooden cylinders, and five hand-colored plates. Price, \$2.75.

Jacobi, A., M.D.,

Professor of Medicine, University of Pennsylvania; Physician to the University Hospital, Philadelphia.

THE LARYNX AND RHINOSCOPY. One volume, 8vo, 252 pages, muslin. Price, \$2.50.

"This is a most valuable and practical work, and should be read by all who are interested in the diseases of the throat and nose. For this reason, to comment it to the professional public is to guarantee no one will be disappointed." *The Medical Record.*

Ross, James, M.D.,

Professor of Medicine, University of Pennsylvania; Physician to the University Hospital, Philadelphia.

THE NERVOUS SYSTEM. Second Edition, one volume, 8vo, 1,014 and 1,051 pages, cloth. Price, \$5.00.

"This is a most valuable and practical work, and should be read by all who are interested in the diseases of the nervous system. For this reason, to comment it to the professional public is to guarantee no one will be disappointed." *The Medical Record.*
 "It is a book which the student of medicine should read, and all the physician should read, in a form so convenient and so complete." *The Medical Record.*

*Diseases of the Nervous System.***Rosenthal, M., M.D.,**

Professor of Diseases of the Nervous System at Yonkers.

A CLINICAL TREATISE ON THE DISEASES OF THE NERVOUS SYSTEM. With a Preface by PROFESSOR CHARCOT. Translated from the *Archives Névrologiques* and Enlarged Edition by L. PRIZEN, M.D., Visiting Physician to the Nervous Diseases, Randall's Island Hospital, Physician to the Charity Nervous Diseases, Bellevue Hospital Outdoor Department, and Pathologist to the Hospital Asylum, Blackwell's Island. Illustrated. One volume, 8vo, 555 pages, uniform. Price, \$5.50.

"For a treatise on diseases of the nervous system, there is no work better arranged or more systematically executed. The author is replete with the more advanced and comprehensive researches in this most difficult field of medical science, and we may safely assert that no other book will give more benefit or information on nervous diseases."—*Practical Medical and Surgical Journal*.

"Among the merits of this book worthy of special mention are its form of plan and systematic divisions and subdivisions; the well-considered amount of space and attention which is devoted to each disease; the careful presentation of the subject of symptomatology, diagnosis, and prognosis; and the numerous conscientious reports of original pathological and histological observations."—*Practical Medical Journal*.

"The book has many merits, and much to commend it to the attention of the profession. It is especially true in regard to the classification, the description of many diseases, and, on the whole, in regard to treatment."—*The Clinical Journal*.

"It is systematically arranged, and is written in a style that is plain, clear, and forceful; and is devoid of hypothetical speculation, affectation, and remarkable errors."—*Practical Medical Journal*.

"The great advances that have been made in the diagnosis, pathology, and treatment of nervous diseases of late years have been set forth in this treatise, and we cheerfully commend the book to the profession."—*Sanitary Review*.

"The work is one which neurologists will scan with the interest."—*Medical and Surgical Review*.

"This work of Rosenthal's has been translated and on the diseases of the nervous system in Germany and Austria to which it refers, having, one through several editions. It has been translated into French, and received the commendations of French Neuro-pathologists, especially Professor Charcot, who is the author of a preface to the American translation. The work is, as it purports to be, a clinical one, being especially full in the practical department of symptomatology and treatment. The numerous original papers are also of importance, and are of a useful kind. The work is enriched by clinical observations and pathological anatomy, and exposure, as far as possible, by physiology, but Dr. Rosenthal devotes very little space to theories, general physiological mechanism, differing but in respect from most recent writers on nervous diseases."—*Practical Medical and Surgical Journal*.

Beard, George M., A.M., M.D.,

Professor of Nervous and Mental Diseases, New York University Medical College, New York; and Visiting Physician, Mount Sinai Hospital, New York.

A PRACTICAL TREATISE ON NERVOUS EXHAUSTION, NEURASTHENIA, ITS SYMPTOMS, NATURE, SEQUELAE, AND TREATMENT. New York: William Wood & Co., 1908. 8vo, 198 pages, uniform. Price, \$1.75.

"The book is written in the practical sense, and is a most concise and readable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae. The author is well known to Dr. Beard, and his treatment is based on the latest researches in the field of nervous diseases."—*Practical Medical Journal*.

"Dr. Beard's book is a most valuable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae. It is a most practical and readable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae."—*Practical Medical Journal*.

"The most practical, fully believing in Dr. Beard's view, based on a scientific basis, the existence of neurasthenia and having a true difference between this form of nervousness and that which is the expression of

neurasthenia, and the value of the treatment of the nervous system. We are sure that this book will be a most valuable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae."—*Practical Medical Journal*.

"The book is a most valuable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae. It is a most practical and readable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae."—*Practical Medical Journal*.

"The practical treatment of nervous exhaustion, neurasthenia, and its sequelae, is a most valuable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae."—*Practical Medical Journal*.

"The book is a most valuable treatise on the subject of nervous exhaustion, neurasthenia, and its sequelae."—*Practical Medical Journal*.

Diseases of the Nerves, Apoplexy.

Gowers, W. R., M.D.,

Professor of Clinical Medicine, University College; Senior Assistant Physician, to University College Hospital, and to the National Hospital for the Paralyzed and Epileptic.

EPILEPSY AND OTHER CHRONIC CONVULSIVE DISEASES. Their Causes, Symptoms, and Treatment. Sold by subscription only. See page 51.

DIAGNOSIS OF THE DISEASES OF THE BRAIN AND SPINAL CORD. Sold by subscription only. See page 51.

Bramwell, B.,

DISEASES OF THE SPINAL CORD. Price, \$5.00.

Putzel, L., M.D.,

Professor of Nervous Diseases, Randall's Island Hospital; Physician to the Class for Nervous Diseases, Hospital Out-door Department; and Pathologist to the Lunatic Asylum, Blackwell's Island.

A TREATISE ON COMMON FORMS OF FUNCTIONAL NERVOUS DISEASES. Sold only by subscription. See page 56.

Lidell, John A., A.M., M.D.

A TREATISE ON APOPLEXY, CEREBRAL HEMORRHAGE, CEREBRAL EMBOLISM, CEREBRAL GOUT, CEREBRAL RHEUMATISM, AND EPIDEMIC CEREBRO-SPINAL MENINGITIS. One volume, 8vo, 395 pages, muslin. Price, \$4.00.

"We think the modest hope of the author, as expressed in the preface to this excellent monograph, that it will prove interesting and useful to those who read it, will be fully realized. We cordially recommend the work of Dr. Lidell to the profession as one of very great value."—*Philadelphia Medical Times.*

"To all friends of ours who are engaged in the study or treatment of cerebral diseases, we cordially commend this work as the most complete and satisfactory of any that we have seen. The mechanical execution of the work is excellent."—*Detroit Review.*

Erichsen, John Eric, F.R.S.,

Professor of Surgery to the Queen's Faculty, Professor of Clinical Surgery in University College, Surgeon to the Hospital, Ex-President of the Royal College of Surgeons of England, and President of the Medical Association, Sec. of the

ON CONCUSSION OF THE SPINE, NERVOUS SHOCK, AND OTHER OBSCURE INJURIES OF THE NERVOUS SYSTEM IN THEIR CLINICAL AND MEDICO-LEGAL ASPECTS. One volume, 12mo, 344 pages, muslin. Price, \$2.25.

The author, in the preface to this preface explains that the subject has long been scarcely mentioned in the annals of the profession, and that it was not until the late war, that it was more fully recognized. He then states that he has published this work in the hope that it may be of some service to the profession, and that it may be of some use to the public, in the knowledge of the nature and extent of the injuries of the nervous system, and of the means of their treatment.

They were not peculiar to them, but might be the consequence of any of the more ordinary accidents of civil life. As the work is now presented, it includes eight additional lectures and has been expanded in scope to embrace a wider range of subjects. In view of the medico-legal aspects of this large, obscure, and important class of injuries of the nervous system, this work is highly valuable.

Richet, Chas., A.M., M.D., Ph.D.,

PHYSIOLOGY AND HISTOLOGY OF THE CEREBRAL CONVOLUTIONS; ALSO, POISONS OF THE INTELLIGENCE. Translated by EDWARD P. FOWLER, M.D. One volume, 8vo, 704 pages, muslin. Price, \$4.50.

"The author has given more attention to the cerebral convolutions, and more attention to the poisons of the intelligence, than is given to their structure."—*Medical and Medical News.*

Diseases of the Brain, Lungs, &c.

Benedikt, Moriz.

• • • • •

ANATOMICAL STUDIES UPON BRAINS OF CRIMINALS—A CONTRIBUTION
TO ANTHROPOLOGY, MEDICINE, JURISPRUDENCE, AND PSYCHOLOGY—Translated from the German, by E. P. Fowler, M.D., New York: Department of Translation, New York Medical College, Secretary, Knickerbocker Hotel, New York. One volume, 8vo, 185 pages, price \$1.00. Paper \$0.50.

Charcot, J. M.,

[illegible]

Hamilton, Allan McLane, M.D.

TYPES OF INSANITY IN THE STRAITS SETTLES IN THE LIGHT OF DIAG-
NOSIS OF MENTAL DISEASE, 1901-1905.

Segin, Edward, M.D.

Diseases of the Urinary Organs.

Bartholow, R., A.M., M.D.

SPERMAFORRHŒA: Its Causes, Symptoms, Results, and Treatment. One volume, 8vo, 128 pages, illustrated, muslin. Price, \$1.25.

Harrison, Reginald, F.R.C.S.,

Surgeon, St. George's Hospital, London. Formerly Lecturer on Anatomy and Surgery at the School of Medicine, University of London. Lecturer, Northern Hospital.

LECTURES ON THE SURGICAL DISORDERS OF THE URINARY ORGANS. Delivered at the Liverpool Royal Infirmary. One volume, 8vo, 399 pages, printed on 100,000 copies, with 100 plates and wood-engravings, muslin. Price, \$4.25.

"The lectures of Mr. Harrison's lectures on the diseases of the urinary organs. The work contains a full and complete description of the various methods of treatment of the various diseases of the urinary organs, with a full description of the various methods of treatment of the various diseases of the urinary organs. The work concludes with a full index."

Coulson, W. J., F.R.C.S.

ON THE DISEASES OF THE BLADDER AND PROSTATE GLAND. Sixth edition, revised. One volume, 8vo, 393 pages, handsomely illustrated. Sold separately, 6s. 6d. only. See page 55.

Neubauer, C., M.D.,

Professor of Chemistry, University of Bonn, Germany, and Lecturer in the Chemical Laboratory in the University of Bonn.

Vogel, J., M.D.,

Professor of Chemistry, University of Bonn.

A GUIDE TO THE QUALITATIVE AND QUANTITATIVE ANALYSIS OF THE URINE. Designed for Physicians, Chemists, and Pharmacists. With a Preface by Professor Dr. B. FRIESEN. Translated from the seventh enlarged edition of the German Edition by LEIBOLD, G. CUTLER, M.D., Physician to the Massachusetts General Hospital, Pathologist at the Boston City Hospital, and Assistant in Pathology in the Medical School of Harvard University. Edited by EDWARD S. WOOD, M.D., Professor of Chemistry in the University of Harvard University. In one superb 8vo volume, 551 pages, printed on 100,000 copies, with 100 plates and four fine chromo-lithographic plates. Price, \$1.80 per copy, \$1.00.

"This is a most valuable and useful work, and the most complete and comprehensive work of its kind in any language. The numerous illustrations are unsurpassed in price and execution. The book is a specimen of art. We seldom see a book so well bound with so excellent and so beautiful a binding." *People's Medical and Surgical Journal.*

Belfield, W. T., M.D.,

Professor of Medicine, University of Bonn.

DISEASES OF THE URINARY AND GENITAL ORGANS. Sold by subscription, 10s. 6d. per volume.

Milton, J. L., M.D., M.R.C.S.

ON THE PATHOLOGY AND TREATMENT OF GONORRŒA. One volume, 8vo, 128 pages, illustrated, muslin. Price, 6s. 6d.

Diseases of the Kidneys, etc.

Dickinson, W. Howship, M.D., Cantab.,

Dr. J. C. H. W. is a member of the New York Academy of Medicine, and is a member of the Academy of Medicine of New York.

ON RENAL AND URINARY AFFECTIONS. This volume has just been completed, and comprises the work of which the volume on Albuminuria, published at Wood's Library for 1884, is the first part. Sold by subscription only. See page 51.

Charcot, J. M., M.D.,

(1) *1813-1814* - Member of Paris; Physician to the Salpêtrière; Member of the Academy of Medicine; Member of the Society of London, of the Clinical Society of Buda Pesth, of the Society of Physicians of Vienna; President of the Anatomical Society; former President of the Société de Médecine.

LECTURES ON BRIGHT'S DISEASE OF THE KIDNEYS, DELIVERED AT THE SCHOOL OF MEDICINE OF PARIS. Collected and published by Drs. BOURNVILLE and SEVESTRE, editors of the *Progres Medical*, and translated, with the permission of the author, by HENRY B. MILLARD, M.D., A.M. Illustrated with two colored plates and with wood-engravings. One volume, 8vo, Pp. 488, including. Price, \$1.50.

The merits of the profession are due to the efforts of Dr. Allard, for, too many times, the profession has its portion of the blame for the embarrassments for the client type, and for the prevalence of the book. It should be the duty of one who desires to be informed of the history of Berg's disease, to *consult*

"Whatever may be thought of Professor Charcot's views of the various forms of mental affection, none can refuse to him the merit of a profound thinker and a most sagacious observer, the philosophic character of his views being at once a record of the knowledge of the day and of the genius of their author."

$$t \text{ minutes } / a \text{ col.}$$

There is a large body of research on the pathology of the human brain. The reasons can be attributed to the nature of the obscurely defined concept of mind in the study of psychology. *Principles of General and*

"The Procedure is valuable enough to be well worth the study of every busy practitioner who has a new case of albuminuria to manage, and it will, of course, find a place upon the library shelves of every one." *American Journal of Medicine* 1894.

Stewart, L. Grainger, M.D., F.R.S.E.,

[illegible]

CHRONIC ALBUMINURIA, OR BRIGHT'S DISEASE OF THE KIDNEYS. One volume, 8vo., 1844, illustrated with seven lithographic plates, cloth. Price, 2s.

... the distinct forms with which the name of Bright is inseparably and honorably associated.

Piffard, Henry G., A.M., M.D.,

ANALYSIS OF THE USE OF PHYSICIANS AND
HOSPITALS IN THE UNITED STATES, 1960-1970. Price, \$1.25.

MADE IN MEXICO AND THERAPEUTICS OF THE SKIN. Sold only by sub-

*Syphilis, Skin Diseases.***Vidal, A. (De Cassis).**

Translated by J. S. A. Adams, M.D., of Pe.

A TREATISE ON VENEREAL DISEASES. With a new chapter on the treatment of the same, with annotations, by GEORGE C. BLACKMAN, M.D., Professor of Syphilis in the Medical College of Ohio; Surgeon to the Cincinnati Hospital, &c. &c. &c. Royal Medical and Chirurgical Society of London. One volume, 8vo., 400 pages, in cloth. Price, \$1.50.

Taylor, R. W., M.D.,

Professor of Dermatology, University of Pennsylvania, Philadelphia.

Author of "New York."

SYPHILITIC LESIONS OF THE OSSEOUS SYSTEM IN INFANTS AND YOUNG CHILDREN. One volume, 8vo., 119 pages, in cloth. Price, \$2.50.

Keyes, E. L., A.M., M.D.,

Professor of Dermatology, University of Pennsylvania, Philadelphia. Author of "The Treatment of Syphilis," "Syphilis," "Hereditary Syphilis," "Syphilis in Infants," &c. &c. &c.

VENEREAL DISEASES. Sold long by subscription. See page 56.

Diday, Paul.

ON SYPHILIS IN INFANTS. Translated by DR. G. WHEATLEY. With Notes and Additions by E. R. SURGUS, M.D. With an extra plate. See page 56.

Dr. Diday is an American doctor of France, who writes of Dr. Stages, as follows: "He believes that this method of treatment will serve the purpose of being put into the general treatment of Infants. I speak from the fact that I have more in my report to work."

Busey, Samuel C., M.D.,

Professor of Dermatology, University of Pennsylvania, Philadelphia. Author of "The Treatment of Syphilis," "Syphilis," "Hereditary Syphilis," "Syphilis in Infants," &c. &c. &c.

CONGENITAL OCCLUSION AND DILATATION OF LYMPH CHANNELS. Translated by Dr. J. S. A. Adams, M.D. One volume, 8vo., 119 pages, in cloth. Price, \$2.00.

Cazenave and Schedel.

MANUAL OF DISEASES OF THE SKIN. Translated by FRANK M. CROZIER, M.D., of Philadelphia. With an extra plate. See page 56.

Living, Robert, A.M., and M.D., Cantab., F.R.C.P., Lond.,

Professor of Dermatology, University of Pennsylvania, Philadelphia.

A HANDBOOK ON THE DIAGNOSIS OF SKIN DISEASES. One volume, 8vo., 119 pages, in cloth. Price, \$1.50.

NOTES ON THE TREATMENT OF SKIN DISEASES. One volume, 8vo., 119 pages, in cloth. Price, \$1.50.

Wilson, Erasmus, F.R.S.

THE STUDENT'S BOOK OF CUTANEOUS MEDICINE AND DISEASES OF THE SKIN. One volume, 8vo., 119 pages, in cloth. Price, \$1.50.

*Skin Diseases, U. S. Pharmacopœia.***Fox, Tilbury, M.D., London.**

Physician to the Hospital for Skin Diseases in the University College.

SKIN DISEASES, their Description, Pathology, Diagnosis, and Treatment. Second American from Third London Edition, re-written and enlarged. With a Catalogue of the Medicines and a Crossed Index, and sixty-seven additional illustrations. One volume, 8vo, 532 pages, muslin. Price, \$5.00.

"The author has written a book that abounds with facts, and is well written, and contains a great deal of information, both as to the diseases of the skin, and the treatment of them."—*British and Foreign Medical Review*.

"A valuable work, well organized in the plan, and containing some of the best facts of the subject."—*Med. Rev.*

"We have no doubt that it will be the student's and the practitioner's pocket guide to the diseases of the skin, in which the author has embodied the most important facts of the most recent investigations, and has given a full and complete account of those affections."—*Edinburgh Medical Journal*. "It will prove a most valuable and ready reference."—*Edinburgh Medical Review*.

"It is a most complete and practical. The author has done a more valuable contribution to our stock on the subject of skin diseases than any other work."—*Edinburgh Medical Review*.

"We would impress all practitioners of medicine with the need of this work, and study it."—*Edinburgh Medical Review*.

"This is a most valuable work, and a volume of the kind in the form of a pocket manual is much needed. The illustrations were small, thick

and smoky. Am. Publ. will now find its successor launched forth as an imposing octavo, luxurious both as to paper and type. This is, in our opinion, a great improvement. We must confess to a virtuous horror of pocket manuals as recalling days when medical students were less sensible and industrious than, happily, they now are. Not only is this volume much enlarged, but is also recast in parts and rewritten. One of the most important new features is the addition of many new illustrations, of which there are now nearly one hundred in the volume. The author has taken great pains to include the latest researches in dermatology in this edition; and we can, therefore, enthusiastically recommend the book to our readers. Without question, it is now the most complete and practical work on cutaneous medicine in the English language. The ordinary student will find in it all that he can desire, and will only be led by its tone to wholesome methods and higher flights of research; while the practitioner will fall back upon its resources with satisfaction and with fresh resolves."—Notice of the new edition in the *London Lancet*, February 8, 1873.

The Pharmacopœia of the United States.

Supplied by Revision. By authority of the National Convention for Revising the Pharmacopœia, held at Washington, A.D. 1880. One volume, 8vo, about 400 pages, steel-plate and tissue paper. Price, \$4.00; leather, \$5.00; leather bound, \$6.00. Unbound, printed on one side, \$5.00.

"Of course, it is a new book, which is one of the best. The Pharmacopœia is the best of its kind, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

contributors."—*Medical Counselor*, Grand Rapids.

"The National Pharmacopœia is the standard authority as to medical preparations. The present revised edition represents the best efforts of the best representative men in the pharmaceutical profession of our country."—*Lancet's Medical News*.

"The committee certainly have lived up to their privileges. They have revised it. Compared with 1870, it is almost a revolution, but a simple revision."—*The Medical Advancer*, New York.

"The committee devoted a little more than two years to the work of revision, and the result is a book which is now before us. On examining the work, we are at once struck with the important differences that exist between the old and the new."—*Medical Record*, Jan. 1, 1881.

"It is on the physician's table, this work is consulted more frequently than any other book in the profession, and it represents the results of the most recent researches. The committee have done a most valuable and meritorious work, and it is a book of which we all should be proud."—*Medical Review*, Jan. 20, 1881.

"The new edition of the Pharmacopœia is a most valuable work, and it is a book of which we all should be proud."—*Medical Review*, New York City.

*Effects of Drugs: Chemistry, Pictures.***Lewin, L., Dr.,**

Associate in the Pharmacological Institute, University of Leipzig.

THE INCIDENTAL EFFECTS OF DRUGS, A PHARMACOLOGICAL AND CLINICAL HANDBOOK. Translated by W. T. ALEXANDER, M.D. One volume, 8vo, 239 pages, muslin. Price, \$2.00.

"That individuals vary in their susceptibility to drugs, and that idiosyncrasies of the most unexpected character reveal themselves in the office, are observations of very ancient date. So far, in, are these accidental effects that sometimes their cause is apt to escape attention. Variations in the phenomena of the action of, and the common therapeutic action of, drugs, depending largely, like the disease phenomena, on the acquired or inherited peculiarities of the patient. Facts

belonging to this category, although not very well defined periods and in the unwritten experience of practitioners, but Dr. Lewin, in this book, by collecting and classifying the evidence and adding to them his own observations, stamped by special study. He reserves credit for breaking new ground in the last place, and in the second for material, with admirable judgment, this somewhat neglected branch of knowledge." *London Medical News.*

Witthaus, R. A., A.M., M.D.,

Professor of Medical Chemistry and Toxicology, the University of Vermont, Member of the American Society of Pharmacologists.

A TEXT-BOOK OF MEDICAL CHEMISTRY. Price, muslin, \$3.50.

"We do not hesitate to recommend this work to the profession as a complete text-book of the subjects of which it treats." *Medical Review.*

"The author has, we think, succeeded fairly in presenting what every practitioner should know of the science of chemistry." *Philadelphia Medical and Surgical Reporter.*

ESSENTIALS OF CHEMISTRY, Organic and Inorganic. Wood's Pocket Manuals. Price, \$1.00.**Draper, John C., M.D., LL.D.,**

Professor of Chemistry in the Medical Department, University of New York, and of Hygiene and Sanitary History in the College of the City of New York.

A PRACTICAL LABORATORY COURSE IN MEDICAL CHEMISTRY. One volume, oblong 8mo, 71 pages interleaved, muslin. Price, \$1.00.

"The book, containing, in a concise and scientific form, all upon the above topics that is of practical value to the physician. We cordially recommend it to physicians and students." *Mississippi Valley Medical Monthly.*

"In printing this book every other page has been left blank in order that the student may be enabled to conveniently record in its proper place the results of the experiments he makes, and of additional facts obtained from analytical chemistry." *Chicago Medical Reporter.*

"In this few books combine so much useful information in a compact, correct, convenient, and readable manner." *Chicago Medical Review.*

"This excellent manual is one of the best adapted for the student or practitioner whose

time is limited, and who desires to secure results with the least expenditure of time, labor and without the aid of an instructor." *Chicago Medical Times.*

"This little work comprises what a young doctor ought to know about chemistry for his patients' good." *Indiana Medical Journal.*

"It is bound so as to open from the top so as to easily lie open on the table, and as it lies open, the lower pages have been left blank for any notes and additions when the student desires to refer as he proceeds with the work." *Chicago Medical Journal.*

"This student is indeed to be congratulated in being able to secure a work of this nature prepared by the order of Dr. Draper's experience." *Chicago Medical Journal.*

Pictures for Physicians' Offices and Libraries.

Edward Jenner, the First Inoculation of a Child, May 14th, 1796.

Andreas Vesalius, the Anatomist.

Spencer Perry House.

Dr. S. A. W. S.

Andreas Pavi, Demonstrating the Use of the Forceps.

The Young Mother.

Size of each, 14x24 inches. Price, each \$2.00. Catalogues of these pictures will be sent upon application.

The Village Doctor.

The Roman Physician.

Sir John Astley.

William Harvey Demonstrating the Circulation of the Blood.

The Anatomical Lecture.

The Accoucheur.

Miscellaneous.

Parkes, E., M.D.

A MANUAL OF PRACTICAL HYGIENE. Edited by F. S. B. FRANCOIS DE CHAMONT, M.D., Sixth Edition. With an Appendix. Giving the American practice on matters relating to hygiene. Prepared by and under the supervision of FREDERICK N. OWEN, C.E., and Sanitary Engineer. Two volumes in one, 8vo, 996 pages. Illustrated with nine full-page plates, and fine wood-engravings, and colored charts. Price, \$5.00.

Sternberg, G. M., M.D.

BACTERIA. By DR. ANTOINE MAGNIN, of Paris, and GEORGE M. STERNBERG, M.D., F.R.M.S., Major and Surgeon, U. S. Army. One volume, 8vo, 494 pages. Illustrated with five full-page plates, including heliotype and lithographic reproductions of photomicrographs. Muslin. Price, \$4.00.

Sternberg, G. M., M.D.

MALARIA AND MALARIAL DISEASES. One volume, 8vo, 332 pages. Muslin. Price, \$2.50. Sold by subscription only. See page 52.

Sturgis, F. R., M.D.

MEDICAL TOPICS, concluding: 1. Hints and Suggestions for Reform in Medical Education. 2. A Plan for the State Regulation of Medicine and Surgery. 3. Medical Education: Its Objects and Requirements. One volume, 8vo, 64 pages, paper cover. Price, 25 cents.

Visiting List (Medical Record), or Physician's Diary.

Continued in the improved features of previous publications of this sort. Prices: For thirty patients a week, handsome red or black leather binding, wallet style, 100 sheets of paper, \$1.25; for sixty patients a week, same style, with or without index, \$1.50.

Hun, H.

A GUIDE TO AMERICAN MEDICAL STUDENTS IN EUROPE. Price, \$1.25.

Steel, J. H., M.D.

OF THE PRINCIPLES OF VETERINARY ANATOMY. A Manual for the use of Veterinary Students. By J. H. STEEL, M.D., V.M.C., 12mo, 312 pages. Muslin. Price, \$3.00.

Buck, A. H., M.D.

RECEIPTS ON HYGIENE AND PUBLIC HEALTH. By various authors. Edited by A. H. BUCK, M.D., New York. In two volumes, royal 8vo, 702 pages. Illustrated with 100 plates of wood-engravings. (Subscription.) Cloth, \$5.00; in leather, \$6.00; and in morocco, \$7.00.

Hospital Plans.

Architectural plans for the Construction, Organization, and Management of Hospitals, and the various departments of the Johns Hopkins Hospital. By the architects and engineers who designed and constructed by lithographic plans.

Johnson and Martin.

ON THE INFLUENCE OF CLIMATIC FACTORS ON EUROPEAN CONSTITUTIONS. By JOSEPH JOHNSON, M.D., and JAMES RUSSELL MARTIN, Esq. From the French of JOSEPH JOHNSON, M.D., American physician. One volume, 8vo, 240 pages. Price, \$1.00.

Kirby, F. O., M.D.

A TEXT-BEAST ON VETERINARY MEDICINE. A. Applied to the Diseases and Management of the Horse, Dog, and Cattle. Selected from Modern Authorities. One volume, 8vo, 400 pages. Illustrated with 100 lithographic plates, containing 1,000 figures. Price, \$5.00. Also in leather, \$6.00. In fine wood-engravings. Sold by subscription only. See page 52.


The works enumerated on the following pages are

SOLD BY SUBSCRIPTION ONLY

BY

MESSRS. WILLIAM WOOD AND COMPANY

Or their authorized Agents.

 Special Catalogues, Prospectuses, Circulars, etc., giving full details respecting their Publication, together with Specimens of their pages, etc., will be sent to any address upon application to the Publishers.

MEDICAL JOURNALS

PUBLISHED BY

WILLIAM WOOD AND COMPANY,

56 and 58 Lafayette Place, New York.

THE MEDICAL RECORD.

A Weekly Journal of Medicine and Surgery.

Edited by GEORGE F. SHRADY, A.M., M.D., Surgeon to St. Francis Hospital, Consulting Surgeon to the Hospital for Ruptured and Crippled, New York. Royal octavo. Price \$5.00 a year. Circulation, over half a million copies annually.

THE MEDICAL RECORD is the best known and most popular medical newspaper published in the United States. It presents a faithful weekly record of the existing state of Medical Science throughout the world, and in consequence of its high scientific character and practical value has gained the countenance and support of the profession, to the extent that it may be justly considered the representative medical journal of America.

Issued weekly, its form and arrangement permit a rapid and satisfactory perusal of its contents—a feature in scientific literature which no one appreciates more highly than the physician, and which journals issued at longer intervals seldom afford.

The information contained in its columns is of the most practical and reliable sort, always keeping in view the requirements of the general practitioner, and maintaining that standard of excellence to be possessed only by an independent and thoroughly scientific journal.

Illustrations of a superior quality are introduced in the text whenever necessary, and excellence in typographical execution is earnestly sought by the publishers.

THE AMERICAN JOURNAL OF OBSTETRICS

And Diseases of Women and Children.

Edited by PAUL F. MUNDÉ, M.D., Professor of Gynecology at the New York Polyclinic and at Dartmouth College; Gynecologist to Mt. Sinai Hospital; Obstetric Surgeon to Maternity Hospital; Fellow of the Obstetrical Society of New York, and of the American Gynecological Society, etc., etc. Collaborators: in London, ROBERT BARNES, M.D.; in Berlin, CARL SCHROEDER, M.D.; in Paris, PIERRE BUDIN, M.D. Octavo. Price, \$5.00 a year, in advance.

This Journal is pre-eminently the accepted organ for communications in its scope; the original articles which have appeared in its pages include many of the most important contributions to medical literature. A very available means, which a liberal outlay of money could employ, has been devoted to its copious illustration, and its presentation in a form of typographical excellence, unqualed by any medical journal in the world. Much of the success of the journal is due to the very satisfactory management of the department of Obstetrics and Gynecology by Dr. Paul F. Munde, and his general supervision of all the literary matter in his capacity as editor-in-chief. The department of Diseases of Children receives the attention which that branch merits and which is an important feature of the journal. The assistance of the distinguished gentlemen whose names are associated with that of the editor as collaborators serves as a further assurance, if such need be, of the usefulness of the journal, and is the result of an arrangement which was made in view of the extensive foreign circulation it has acquired.

THE AMERICAN JOURNAL OF OBSTETRICS has 112 pages, each number, or, 1,344 pages in the year, or more than three times the number contained in the first volume.

THE AMERICAN JOURNAL OF OBSTETRICS will be sent to pull-up subscribers for the MEDICAL RECORD for \$4.00 a year.

Journal of Cutaneous and Venereal Diseases

(ISSUED MONTHLY.)

Edited by **PRINCE A. MORROW, A.M., M.D.**, Professor of Venereal Diseases, University of the City of New York; Surgeon to Charity Hospital; Surgeon to the Bureau of Outdoor Relief, Bellevue Hospital. Price, \$2.50 per year.

The diseases which it is the object of this Journal to embrace are of such manifest interest and importance, and occur with sufficient frequency in the practice of every physician to warrant their special consideration in an appropriate channel. At the present time there are five journals devoted to these subjects, this being the only one in the English language. It is intended to serve as a medium of communication between those who have given special attention to these subjects, referred to, and those members of the profession in general practice whose time or opportunities will not permit them to engage in these special lines of investigation. In carrying out this aim it is the intention of the publishers that the Journal shall be eminently practical.

The colored plates and wood engravings that have thus far appeared are the best that can be produced, and this feature will be fully maintained as its practical utility to the subscriber is very generally acknowledged.

THE JOURNAL OF CUTANEOUS AND VENEREAL DISEASES will be sent to paid-up subscribers for the MEDICAL RECORD or the AMERICAN JOURNAL OF OBSTETRICS for \$2.00 a year.

AMERICAN DRUGGIST.

An Illustrated Journal of Pharmacy, Chemistry and Materia Medica.

FRED'K A. CASTLE, M.D., Editor. **CHAS. RICE, Ph.D., Associate Editor.**

Ask any druggist to allow you to examine the latest issue of the AMERICAN DRUGGIST, an Illustrated Monthly Journal of Pharmacy, Chemistry and Materia Medica, or, if you choose, to subscribe for it.

THE PRICE IS ONLY \$1.50 A YEAR.

It contains the latest and most reliable information regarding New Remedies, New Methods of Treatment, New Formulas, valuable information regarding Official and Unofficial Preparations, Analysis of Nostrums, etc., etc., all profusely illustrated, forming a complete record of the progress of Pharmacy, Materia Medica and Therapeutics. Try it for a year. Sample copies sent free upon application.

LIST OF PRICES.

The Medical Record, - - -	\$5.00	Journal of Cutaneous and Ven-	
The American Journal of Obstet-		ereal Diseases, - - -	\$2.50
rics, - - - - -	5.00	American Druggist, - - -	1.50

CLUB DISCOUNTS.

When several of the Journals are subscribed for to be mailed to the same address, the following Club Rates are offered:

Medical Record, {	Full Price	
American Journal of Obstetrics, }	\$10.00	Club Discount, - \$1.00
Medical Record, {	7.50	Club Discount, - .50c
Journal of Cutan. and Ven. Dis., }		
American Journal of Obstetrics, {	7.50	Club Discount, - .50c
Journal of Cut. and Ven. Dis., }		
Medical Record, {	12.50	Club Discount - 1.50
American Journal of Obstetrics, }		
Journal of Cutan. and Ven. Dis., }		

✓ SUBSCRIPTIONS MUST BE ARRANGED TO EXPIRE JUNE 30 OR DECEMBER 31.

At the above prices only when paid in advance to William Wood & Company, 200, Park Avenue, NOT by check.

REFERENCE HANDBOOK OF THE MEDICAL SCIENCES

MESSRS. WILLIAM WOOD & COMPANY beg leave to announce to the Medical Profession of the United States and Canada that they have for several years past been organizing and arranging for the preparation of a work differing from all others heretofore published in this country, and calculated, as they believe, from its character and the high reputation of its very large staff of learned contributors, to be the most practically and generally useful work to all classes of the Medical Profession of any hitherto presented to their notice.

It is designed that this superb work shall cover so wide a field, and embrace such a great variety of topics, as to make it of the greatest practical utility, *not only to the general practitioner, but also to those who are interested more particularly in special departments of medical knowledge.*

This work will consist of a collection of concisely written essays on **all the Important Topics belonging to the Broad Domain of Medicine.** Its subject matter will be alphabetically arranged, thus admitting of instant and ready reference, with no loss of time in reading over page after page to find the precise item desired.

The number of those physicians who take an interest in the more purely scientific branches of medical knowledge is already quite large, and will undoubtedly increase as time goes on. It has, therefore, been thought advisable to *devote considerable space* to the more important topics belonging to the domain of **Anatomy, Embryology, Histology, Physiology, Physiological and Pathological Chemistry, Pathological Anatomy, Climatology, and Medical Botany.** All these topics, as well as those belonging to the more directly practical branches of medical knowledge, such as **General Pathology and Therapeutics, Surgery, Gynecology, Neurology, Otolaryngology, Ophthalmology, Obstetrics, &c., &c., &c.,** have been assigned to writers who are experts in their respective departments.

Woodcut illustrations of the best character and full-page lithographic plates will be a decided feature in the work, and will be introduced whenever the authors themselves think that they will serve to elucidate the text. Special pains have been taken to secure as large a proportion of original cuts as possible, and *no expense has been spared in this or in any other branch of the work to secure the highest order of excellence.*

THE REFERENCE HANDBOOK OF THE MEDICAL SCIENCES, will be limited to a series of **Eight splendid Imperial Octavo Volumes.**

The pages are of large size, in double columns, and the type as small as can be read comfortably by a person of ordinary good eyesight. Each page carrying as many words as three ordinary full octavo pages.

It will be seen that the volumes admit of the introduction of an enormous amount of reading matter equal to the contents of twenty or more of the usual size of octavo volumes, which will be of so practical a nature that the entire work will be literally a handbook for the daily, constant reference of the Profession,—a book which *no physician*, whose time is worth anything at all, *can afford to be without.*

Large treatises on the Practice of Medicine and Surgery, and on other branches of medical knowledge have their appropriate and necessary place,—for long, deliberate and patient study,—but this **Reference Handbook** is designed to furnish not only **Thorough and Exhaustive Information,** but to be a constant daily companion and ready reference on all branches of **Medicine and the Collateral Sciences.**

A large part of the material contained in this HANDBOOK is original work, and upon certain points the information now to be presented has heretofore been inaccessible, by reason of its being scattered through governmental and other manuscript records and volumes.

Prices at which this work will be sold:

In extra English cloth binding, per volume, \$6.00; in fine leather, raised bands, per volume, \$7.00; in extra Turkey morocco, English cloth sides, per volume, \$8.00.

Send for a complete Circular giving list of Contributors (300) and Specimen pages from the Work free.

WOOD'S LIBRARY

or

STANDARD MEDICAL AUTHORS.

IN announcing the volumes in this now celebrated series, it may seem, to those discerning and appreciative gentlemen who, from the first have availed themselves of this project, unnecessary to repeat what we have said at different times in former years respecting the general character and make up of these books. And yet, as many thousands have entered the profession since then, we consider it not amiss briefly to renew some of the more important features of this most successful scheme to supply standard medical literature at low prices. Until 1879 no attempt had ever been made by medical publishers to produce books at less than the large prices, rendered necessary, indeed, by the limited sale attained by most. It was a bold venture to undertake to publish twelve volumes in one year, at but one quarter to one-tenth the prices previously obtained—an undertaking which could only be successful from a sale vastly larger than before attained.

That it was a success, and a grand one, is certain evidence of the wisdom of the plan, and of the sure support which the profession will always accord to enterprises conceived and carried out as this has been.

In the seven years during which this library has been annually published,

EIGHTY-FOUR VOLUMES

of most valuable medical books have been issued, at a cost to the regular subscribers

OF ONE HUNDRED AND FIVE DOLLARS ONLY;

but representing books (as published in other editions) to the

VALUE OF NEARLY FIVE HUNDRED DOLLARS.

It would be impossible for any publisher to issue books of the high character proposed, at such a nominal price, unless a sale could be guaranteed for the series very much larger than is ever attained by books as ordinarily published. We have contracted with *well-known authors*, eminent in their specialties, for *new and original works* upon subjects of present interest to practitioners, and especially conforming to the essentially practical character for which we desire this series to be known.

If the expense of the production of such books as are contemplated were to be borne by the sale of an ordinary edition, the individual cost of such volumes would be from \$5.00 to \$7.00 each, and in several instances much more. By combining these volumes into sets or series of twelve volumes each—by the closest scrutiny of every item that enters into their cost of production and sale—by presenting them in a form so attractive that they will enlist the interest of every lover of books, and by the adoption of a system of distribution by which every book-buying physician can have the opportunity of examination and purchase, we are enabled to supply the series at the uniform price of \$18.00 each, except as hereinafter named.

Seven series have been published, and we shall soon enter upon the publication of the eighth. The very remarkable success that has attended this enterprise from its inception, the widespread support which it has continued to receive, has served as a constant stimulant to further effort in continuation of the libraries upon the same high standard of excellence.

PRESS NOTICES.


"The publishers are to be thanked for their untiring energy and zeal, in thus furnishing the profession with such valuable publications at so low a price—scarcely a nominal consideration."

"We know of no better investment than a subscription to these series."

"Think of it! twelve new books upon the various branches of medicine and surgery, by the best writers of to-day, for only \$18.00!"

WOOD'S LIBRARY OF STANDARD MEDICAL AUTHORS.

"A Valuable Addition to every Physician's Library."



"Continues to Delight and Surprise us."

"This is certainly a bold undertaking, and can only be successful by a liberal support from the profession."

"One of the reasons why medical books are so expensive, as compared with books of general literature, is their comparatively limited sale."

"Of course the only way in which the enterprise can pay its organizers is by an extensive subscription list, and we earnestly ask for a general support along our line."

"The publishers look for the support of the profession in this enterprise."

"The science is one which the profession should endorse."

"This will furnish progressive men with a rare opportunity to procure new books at exceedingly low figures."

"Most of these volumes will be illustrated. Nothing yet offered will at all compare with this proposition."

"As the work is an experiment, we hope it will prove sufficiently remunerative to our contributors during future years. The terms are so reasonable that a library of these volumes is now within the reach of all."

"We have no doubt that every medical man in the profession should enter his name as a contributor."

"We, Wood & Co., wish it possible for every practitioner, however poor his purse, to have a complete library of fine works, and they should receive the heartiest support from the profession."

The above Press Notices show the welcome given to the enterprise at its inception.

"Having for the past year the cooperation of the Press, the following notices are published for the benefit of the public."

"The quality of the material, the paper, type, presswork and binding are all

"of the highest quality, and the artistic work, especially, it is

"of the highest quality, and the artistic work, especially, it is

"of the highest quality, and the artistic work, especially, it is

Catalogue of the Titles of the Works published in WOOD'S LIBRARY OF STANDARD MEDICAL AUTHORS. 1884.

Sixth Series. Price, \$15.00. Volumes not sold separately.

LEGAL MEDICINE. By CHARLES MEYMOTT TIDY, M.D., F.R.S., *Master of Surgery, Professor of Chemistry, and of Forensic Medicine and Public Health at the London Hospital, Medical Officer of Health for Islington, Life Deputy Medical Officer of Health and Public Analyst for the City of London, &c.* Volume III. Contents: Legitimacy and Paternity—Pregnancy, Abortion—Ripe, Inducement—Exposure—Suffocation, Bristling—Live Bath, Infanticide—Asphyxia, Drowning—Hanging, Strangulation—Execution.

PATHOLOGY AND TREATMENT OF GONORRHOEA. By J. L. MILTON, M.D., M.R.C.S., *Lecturer on Diseases of the Skin, St. John's Hospital for Skin Diseases, &c.*
This work is fresh from the author's hands, and treats in a very practical way of this common disease.

DIAGNOSIS AND DISEASES OF THE HEART. By DR. CONSTANTINE PAUL, *Professor Agrégé at the Faculty of Medicine of Paris, &c.* Illustrated by numerous fine wood engravings.

A PRACTICAL MANUAL OF OBSTETRICS. By E. VERRIER, M.D. Translated from the French, and Edited by L. PARTNIDGE, M.D. Profusely illustrated with fine wood engravings.

HUNTER'S PHYSICIAN'S VADE MECUM: A Manual of the Principles and Practice of Physic; with a Synopsis of General Pathology, Therapeutics and Hygiene. Tenth Edition. Revised by WILLIAM ARTHUR'S, M.D., M.B., Cantab., F.R.S., JOHN HARLEY, M.D., Lond., F.R.S. Volume I. Illustrated by numerous engravings.

HUNTER'S PHYSICIAN'S VADE MECUM: A Manual of the Principles and Practice of Physic; with a Synopsis of General Pathology, Therapeutics and Hygiene. Tenth Edition. Revised by WILLIAM ARTHUR'S, M.D., M.B., Cantab., F.R.S., JOHN HARLEY, M.D., Lond., F.R.S. Volume II. Illustrated by numerous engravings.

MALARIA AND MALARIAL DISEASES. By GEORGE M. STERNBERG, M.D., F.R.M.S., *Major and Surgeon, United States Army; Member of the Bohemian Society of Washington; Late Member of the Hungarian Imperial Commission of the National Board of Health, &c.* Illustrated.

DISEASES OF THE ESOPHAGUS, NOSE AND NASO-PHARYNX. By MORRELL MACKENZIE, M.D., F.R.C.S., *Professor of the Practice of the Hospital for the Diseases of the Chest and Throat, Lecturer on Diseases of the Throat at the London Hospital Medical College, &c.* Illustrated by wood engravings.
This is a new and valuable work, namely, "Diseases of the Pharynx, Larynx and Trachea," published in 1883, and earned the warmest commendation from the medical press throughout the world.

A TEXT-BOOK OF GENERAL PATHOLOGICAL ANATOMY AND PATHOGENESIS. By ROBERT LUDLAM, *Professor of Pathology and Anatomy in the University of Glasgow.* Translated by J. M. MACALISTER, M.D., M.B., Cantab., F.R.C.S., DONALD MACALISTER, A.M., M.B., *Member of the Royal College of Physicians, Lecturer in Medicine, Lecturer at St. John's College, Cambridge.* Illustrated by numerous engravings. Second Edition. Profusely Illustrated.

DISEASES OF THE URINARY AND MALE SEXUAL ORGANS. By WM. T. BULFIELD, M.D.

ESOPHAGEAL AND PULMONARY DISEASES. By PROF. SIR JAMES, M.D., *Lecturer on Materia Medica and Surgery, and at the London Hospital; Physician to the Hospital for Diseases of the Throat and Chest, and at the North London Dispensary Hospital, &c.* Illustrated by numerous wood engravings.

MEASLES, RUBEOLY, AND OTHERS ON PEARLS AND IN MEDICINE. By LAURENCE JOHNSON, A.M., M.D., *Professor of Materia Medica, and of the Department of the University of the City of New York; Fellow of the Royal College of Physicians, &c.* Illustrated by nine beautifully colored plates and very numerous engravings.

Wood's LIBRARY OF STANDARD MEDICAL AUTHORS.

Fourth Series. Price, \$18.00. Volumes not sold separately.

ILLUSTRATIONS OF DISSECTIONS. In a series of original colored plates, representing the dissections of the human body, with descriptive letter-press. By GEORGE VINER ELLIS, *Professor of Anatomy, Lecturer on Surgery, London*, and G. H. FORD, Esq. The drawings are from nature by M. Ford, from directions by Prof. Ellis. Volume I. Containing 29 full page chromo-lithographic plates.

ILLUSTRATIONS OF DISSECTIONS. In a series of original colored plates, representing the dissections of the human body, with descriptive letter-press. By GEORGE VINER ELLIS, *Professor of Anatomy, Lecturer on Surgery, London*, and G. H. FORD, Esq. Volume II. Containing 27 full page chromo-lithographic plates.

As the first series, we succeeded in presenting our subscribers with "Savage's Female," a volume which, on the page of the original plates, we supposed we had reached the extreme limit in representing the human body. At a price, but these two volumes of Ellis and Ford have exceeded even this. This series, we believe, cannot fail to command acknowledgment of the value of this series of plates, and we are happy to be able to accomplish such results, save in a library such as this, which is a new and original one, and equal side.

LECTURES ON DISEASES OF CHILDREN. A Hand-book for Physicians and Students. By Dr. EDWARD B. NORTH, *Director of the Clinic and Polytechnic for Diseases of Children in the Royal Children's Hospital, Professor of the Royal University*. Translated from the German. 87. A new and original work, of great practical value.

MATERIA MEDICA AND THERAPEUTICS. Inorganic Substances. By CHARLES D. F. PHILLIPS, M.D., F.R.S.E., *Lecturer on Materia Medica, Westminster Hospital, London*. Adapted from the German of Dr. J. J. ROBERTSON, L. By LAWRENCE JOHNSON, M.D. Volume I.

MATERIA MEDICA AND THERAPEUTICS. Inorganic Substances. By CHARLES D. F. PHILLIPS, M.D., F.R.S.E., *Lecturer on Materia Medica, Westminster Hospital, London*. Adapted from the German of Dr. J. J. ROBERTSON, L. By LAWRENCE JOHNSON, M.D. Volume II.

Since the publication of the learned author's treatise on the Materia Medica and Therapeutics of the Inorganic Substances, in the first series, there has been a continued inquiry for this promised continuation. We are happy to be able now to present it fresh from the hands of Dr. Phillips, and published in a new and original form, by special arrangement with him.

PRACTICAL MEDICAL ANATOMY. A guide to the physician in the Study of the Relations of the Various Organs, either in Health and Disease, and in the Diagnosis of the Medical and Surgical Conditions of the Human Body. By AMBROSE L. RANNEY, A.M., M.D., *Professor of Anatomy and Lecturer on Gynecology and Minor Surgery in the Medical Department of the University of the City of New York; also Surgeon to the Northern and Northern Dispensaries, Resident Physician of the New York Academy of Medicine; Member of the Medical Society of the County of New York; Author of "The Art of Anatomy of the Nervous System," and "The Art of Dissection of the Nervous System," etc., etc.* Illustrated by J. J. ROBERTSON, L.

This is a new and original work, of a new and original character.

MENTAL PATHOLOGY AND THERAPEUTICS. By W. GRIEßINGER, M.D., *Professor of Clinical Medicine and Medical Science at the University of Berlin; Honorary Member of the Medical Association, Assistant Member, Assoc. Lecturer at the Society of Medical Physicians in Berlin, etc.* Translated from the German by C. EDWARD ROBERTSON, M.D., *Chief of the Medical Staff of the Society of Physicians in Berlin, Honorary Member, and JAMES R. HILFORD, M.D., Lecturer.*

This is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority.

DISEASES OF THE RECTUM AND ANUS. By CHARLES D. KELSEY, M.D., *Surgeon to St. Peter's Hospital, for Diseases of the Rectum, Consulting Surgeon for Diseases of the Rectum, to the St. Peter's Hospital Dispensary, for Women and Children, etc., etc.*

ON ASTHMA, ITS PATHOLOGY AND TREATMENT. By HENRY HYDE SALTOR, M.D., *Lecturer on the Pathology of the Lungs, Professor of the Pathology of the Lungs, and Lecturer on the Pathology of the Lungs, at the University of the City of New York, and Lecturer on the Pathology of the Lungs, at the University of the City of New York.*

RHEUMATISM, GOUT, AND SOME OF THE ALLIED DISEASES. By MORRIS LONG, M.D., *Professor of the Pathology of the Lungs, at the University of the City of New York, and Lecturer on the Pathology of the Lungs, at the University of the City of New York.*

This is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority.

LEGAL MEDICINE. By W. W. HARRIS, M.D., F.R.S., *Master of Surgery, Professor of the Pathology of the Lungs, at the University of the City of New York, and Lecturer on the Pathology of the Lungs, at the University of the City of New York.*

LEGAL MEDICINE. By W. W. HARRIS, M.D., F.R.S., *Master of Surgery, Professor of the Pathology of the Lungs, at the University of the City of New York, and Lecturer on the Pathology of the Lungs, at the University of the City of New York.*

This is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority.

This is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority. It is a new and original work, of a new and original character, and of the highest authority.

The stereotype plates of this series are partially destroyed, and after the few sets remaining are sold, no more can be supplied.

Catalogue of the Titles of Works published in

WOOD'S LIBRARY OF STANDARD MEDICAL AUTHORS.

Third Series. Price, \$18.00. Volumes not sold separately.

N ALBUMINURIA. By A. H. F. ... N. M. ...

MATERIA MEDICA AND THERAPEUTICS OF THE SKIN. By H. ...

A TREATISE ON DISEASES OF THE JOINTS. By RICHARD ...

A TREATISE ON THE CONTINUED FEVERS. By JAMES ...

A MEDICAL FORMULARY. By LAWRENCE ...

THE DISEASES OF OLD AGE. By J. M. ...

COULSON ON THE DISEASES OF THE BLADDER AND PROSTATE. By ...

GENERAL MEDICAL CHEMISTRY. By ...

ARTIFICIAL ANAESTHESIA AND ANAESTHETICS. By ...

A TREATISE ON ...

A HANDBOOK OF UTERINE THERAPEUTICS AND DISEASES OF WOMEN. By ...

DISEASES OF THE EYE. By ...

The stereotype plates of this series are partially destroyed, and after the few sets remaining are sold, no more can be supplied.

Catalogue of the Titles of the Works published in WOODS' LIBRARY OF STANDARD MEDICAL AUTHORS.

Second Series. Price, \$18.00. Volumes not sold separately.

VENEREAL DISEASES. By E. L. KYLE, A.M., M.D., *Adjunct Professor of Surgery, and Professor of Clinical Surgery, Boston Hospital Medical College; Consulting Surgeon to the Charity Hospital; Surgeon-in-Chief of the Boston Hospital.*

This is a practical treatise of 161 pages, thoroughly covering the subject. It is written with especial reference to the physician in active practice, and is well illustrated.

A HANDBOOK OF PHYSICAL DIAGNOSIS: Comprising the Thorax, Thorax, and Abdomen. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the Third Edition of the French of ALFRED POULET, M.D., *Prof. Fac. Physicians and Surgeons, etc.* American Edition, by J. C. ALLEN, M.D., with numerous illustrations.

This is a work of the highest authority upon the subject, having passed through several editions in France, and has been translated into French, Italian, Russian, Spanish, Polish, and English. A new edition has been published.

A TREATISE ON FOREIGN BODIES IN SURGICAL PRACTICE. By ALFRED POULET, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French of ALFRED POULET, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French. Volume I.

This is a work of the highest authority upon an entirely new subject, of unusual interest and value. It is written by the author, who has revised and corrected it, with additions, especially in the section on foreign bodies in the eye, and is illustrated by many fine engravings.

A TREATISE ON FOREIGN BODIES IN SURGICAL PRACTICE. By ALFRED POULET, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French of ALFRED POULET, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French. Volume II.

A TREATISE ON COMMON FORMS OF FUNCTIONAL NERVOUS DISEASES. By L. J. KYLE, A.M., M.D., *Adjunct Professor of Surgery, and Professor of Clinical Surgery, Boston Hospital Medical College; Consulting Surgeon to the Charity Hospital; Surgeon-in-Chief of the Boston Hospital.*

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

DISEASES OF THE PHARYNX, LARYNX AND TRACHEA. By MORRELL MAC KENZIE, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French of MORRELL MAC KENZIE, M.D., *Prof. Fac. Physicians and Surgeons, etc.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

THE SURGICAL PATHOLOGY AND SURGICAL ANATOMY OF THE PELVIC ORGANS. By J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French of J. C. ALLEN, M.D., *Professor of Medicine, University of Boston.* Translated from the French.

This is a work of the highest authority upon a subject of general practitioners, and treats in a practical way of the most common forms of functional nervous diseases. It contains a book of 262 pages.

The stereotype plates of this series are partially destroyed, and after the few sets remaining are sold, no more can be supplied.

Catalogue of the Titles of Works published in

Wood's LIBRARY OF STANDARD MEDICAL AUTHORS.

First Series. Price, \$18.00. Volumes not sold separately.

KENT AND PAIN. A Course of Lectures on the Principles of Medicine, as far as they relate to the Treatment of Acute and Surgical Diseases, and the Death of a Variety of Cases. By JOHN KENT, M.D., F.R.S., F.R.C.P., &c. Edited by WILLIAM JACOBSON, F.R.C.S.

DISEASES OF THE INTESTINES AND PERITONEUM. *Chirurgia Abdominis, &c.* By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, and the Principles of the Practice of Surgery, in the University of London. By JOHN STEER BRISTOAL, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By THOMAS WILLIAMS, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By WILLIAM RANSOM, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By JOHN STEER BRISTOAL, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A CLINICAL TREATISE ON DISEASES OF THE LIVER. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A CLINICAL TREATISE ON DISEASES OF THE LIVER. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A CLINICAL TREATISE ON DISEASES OF THE LIVER. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

MATERIA MEDICA AND THERAPEUTICS. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A CLINICAL TREATISE ON THE DISEASES OF THE NERVOUS SYSTEM. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A CLINICAL TREATISE ON THE DISEASES OF THE NERVOUS SYSTEM. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

THE LIVES OF THE MEN. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

ON VITALITY AND ITS INFLUENCE ON LIFE. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A PRACTICAL MANUAL OF THE DISEASES OF CHILDREN, WITH A PREGNANT. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

A MANUAL OF SURGERY. By JOHN RICHARD WARDLE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London. By CHARLES MACKENZIE, M.D., F.R.C.P., &c. Lecturer on the Principles of the Practice of Medicine, in the University of London.

NEARLY ONE HUNDRED
DISTINGUISHED AUTHORS,

HAVE CONTRIBUTED TO

WOODS' LIBRARY

OF

Standard Medical Authors,

INCLUDING

PROFS. A. L. LOOMIS—J. M. DA COSTA—HENRY D. NOYES—E. L. KEYES—A. H. BUCK—
P. F. MUNDT—H. G. PIERARD—R. A. WITTHAUS—H. M. LYMAN—M. ROSENTHAL—
P. T. FRIEDRICH—W. ERB—E. I. PARTRIDGE—A. TROUSSEAU—J. M. CHAR-
COT—E. ZIEGLER—E. A. PARKES—W. GRIESINGER—G. V. ELLIS.

AND

DRS. L. JOHNSON—PROSSER JAMES—J. C. WILSON—R. BARWELL—G. M. STERNBERG—
MORRELL MACKENZIE—L. S. PILCHER—C. M. TIDY—L. PUTZEL—H. SAVAGE—
W. B. CARPENTER—BENJ. HART—C. PAUL—C. B. KELSEY—C. D. F. PHIL-
LIPS—J. HILTON—W. T. BELFIELD—W. J. COULSON—L. TAIT—F. W.
PAVY—P. GUTTMANN—W. F. CLARKE—ETC., ETC., ETC.

THERE HAS BEEN INCLUDED IN THE LIBRARY, IN THE FIRST SIX SERIES,

THIRTY-FOUR

Beautifully Colored Full-page Lithographic Plates,

ONE HUNDRED AND SEVENTY

BLACK AND TINTED

FULL PAGE LITHOGRAPHIC PLATES,

IN ALL

OVER TWO HUNDRED PLATES,

CONTAINING SEVERAL HUNDRED FIGURES,

AND NEARLY

FIVE THOUSAND

OF THE FINEST AND MOSTLY ORIGINAL WOOD ENGRAVINGS,

ALL OF WHICH HAS BEEN SUPPLIED AT THE MERELY
NOMINAL COST OF

FIFTEEN DOLLARS A YEAR
TO THE REGULAR SUBSCRIBER

Ziemssen's Cyclopædia of the Practice of Medicine.

Twenty Royal octavo volumes, including Bielefeld, Hagen, &c., and the German Literature, WITH WHICH SELF IS PRESENTED A COPY OF

ZIEMSEN'S HANDBOOK OF DISEASES OF THE SKIN.

This great work, the most full and complete treatise upon the practice of medicine in any language, is now complete. It is a standard which will long maintain its place as the great storehouse of medical knowledge.

It seems proper upon the conclusion of this great work, to ask attention to a few facts in connection with its translation and reprint. The attention of Wm. Wood & Co. was first directed to the advertisement of the German publisher, in the early part of 1873. After consultation with a number of prominent members of the profession, they concluded to venture upon the translation and publication in English of the work to which it referred, the largest undertaking by far, both as to the editorial labor and the amount of money involved, of any medical publication in the English language.

The estimates of the cost of the volumes, and the price at which the publishers could sell them, were based wholly upon the published plans of Mr. Vogel, thus, while the "copy" came from Germany with most satisfactory regularity, it was soon found that the volumes would much exceed the estimated number of pages; and, after several years of publishing largely increased volumes, it was found that, even with such addition, it would be impossible to complete the work in the fifteen volumes, as at first announced. The alternative then presented was, to increase the number of volumes, or abridge the matter; the latter course would have been much preferred by the publishers, as enabling them to adhere strictly to their original estimates. So many influential subscribers objected, however, that they felt compelled to announce that two volumes would be added to the set. It was hardly to be expected that there would be absolutely no adverse criticism of this course among so large a constituency as the subscribers to this work. It has been very gratifying, however, to find that this change, wholly beyond their control, has been unsatisfactory to less than one per cent. of the subscribers. To show the difference between what they promised and what has actually been given subscribers to Ziemssen's Cyclopædia, the publishers give here with the number of pages of each volume, in comparison with the average number as at first estimated, viz.:

Promised, per volume, 500 to 700 pages.		Average of 600.	
Vol. 1	721 pages.	Vol. 7	1,060 pages.
" 2	763 "	" 8	940 "
" 3	684 "	" 9	936 "
Given, " 4	821 "	" 10	584 "
" 5	726 "	" 11	636 "
" 6	1,038 "	" 12	944 "
Total number of pages <i>actually received</i> by Subscribers, - - -		14,596	
" " " <i>promised</i> by Publishers, - - -		10,260	

Given to Subscribers above their expectations, - - - 4,336 pages, *or the equivalent of 7 volumes of over 600 to 700 pages each.* To the above is now to be added still another extra volume, Diseases of the Skin.

It is, of course, well known that though this extra matter has cost the subscribers nothing, it entailed a very heavy and unlooked for expense upon the publishers. They believe, however, this endeavor to do justly and liberally, by those who had supported this great undertaking, has been appreciated by them.

It is with no little satisfaction that the publishers can look back to the unexampled regularity, in book-publishing, with which the volumes appeared every three months for over three years, and it is with great pleasure that the publishers can now congratulate the subscribers and themselves upon the completion of the work.

WM. WOOD & CO., Publishers.

ZIEMSEN'S CYCLOPÆDIA OF PRACTICE OF MEDICINE.

This great work, the most full and complete treatise upon the practice of medicine in any language, is now complete. It is a standard which will long maintain its place as the great storehouse of medical knowledge.

After a closing the period of the publication of this great work, the Publishers desire again to call attention to the great advantages to be derived from the purchase of the work, after promised by them to the subscribers and the amount that will be finally received from them.

<i>Promised, per Volume, per 1000</i>				<i>Average of 600.</i>		
	VOLUME 1	124 pages.	VOLUME 7	1,060 pages.		
<i>Given</i>	" 2	764 "	" 8	949 "	VOLUME 12	987 pages.
	" 3	644 "	" 9	936 "	" 14	961 "
	" 4	524 "	" 10	582 "	" 15	808 "
	" 5	404 "	" 11	656 "	" 16	1,071 "
	" 6	284 "	" 12	994 "	" 17	982 "
Total, as promised by Subscribers.				14,696		
Total, as promised by Publishers.				10,200		

Given to Subscribers above their expectations, - - 4,496 pages,
which will be received from the subscribers, 700 pages each. To the above is now to be added still another advantage, which will be received from the subscribers.

CONTENTS.

Vol. 1. A. Diseases of the Skin. Part 1.	Vol. 12. Diseases of the Brain and its Membranes.
2. A. Diseases of the Skin. Part 2.	13. Diseases of the Spinal Cord and Medulla Oblongata.
3. A. Diseases of the Nervous System. Part 1.	14. Diseases of the Nervous System and Disturbances of Speech.
4. A. Diseases of the Nervous System. Part 2.	15. Diseases of the Kidney.
5. A. Diseases of the Locomotive Apparatus. Part 1.	16. Diseases of the Locomotive Apparatus and General Anomalies of Nutrition.
6. A. Diseases of the Locomotive Apparatus. Part 2.	17. General Anomalies of Nutrition and Poison.
7. A. Diseases of the Locomotive Apparatus. Part 3.	18. Hygiene and Public Health. Part 1.
8. A. Diseases of the Locomotive Apparatus. Part 4.	19. Hygiene and Public Health. Part 2.
9. A. Diseases of the Locomotive Apparatus. Part 5.	20. General Index to the Whole Cyclopædia.

ZIEMSEN'S HANDBOOK OF THE DISEASES OF THE SKIN PEARSON'S SUPPLEMENT TO ZIEMSEN'S CYCLOPÆDIA.

More than 100,000 copies of this work have been ordered upon this enterprise, and the Publishers are now in receipt of the Publishers for Catalogues.
Price per Volume, Cloth, \$5.00; Leather, \$6.00; Half Morocco, \$7.50.

SPECIAL TERMS.

Now that the great work is completed, the set will be sent in full, to parties of **wholesale order, and liberal terms of payment as to time, allowed.**

WILLIAM WOOD & COMPANY, Publishers,

56 & 58 Lafayette Place, New York.

GENERAL INDEX TO ZIEMSEN'S CYCLOPÆDIA.

**Ninety and Ninety-Eight Columns of References to
DISEASES—THEIR COMPLICATIONS AND REMEDIES.**


This is a complete Index to the whole work, and is a valuable set of the references of the physician, recording the progress of the disease, the position of the patient, the treatment, the progress of the disease, and the result of the treatment. It is a complete Index to the whole work, and is a valuable set of the references of the physician, recording the progress of the disease, the position of the patient, the treatment, the progress of the disease, and the result of the treatment.

This is a complete Index to the whole work, and is a valuable set of the references of the physician, recording the progress of the disease, the position of the patient, the treatment, the progress of the disease, and the result of the treatment.

IN THE REPUTATION OF THE WORK ALONE.

This is a complete Index to the whole work, and is a valuable set of the references of the physician, recording the progress of the disease, the position of the patient, the treatment, the progress of the disease, and the result of the treatment.

THE INTERNATIONAL ENCYCLOPÆDIA OF SURGERY.



CONTENTS OF THE VOLUMES.

VOL. I	GENERAL PRINCIPLES OF SURGERY. AMPUTATIONS
VOL. II	WOUNDS AND INJURIES WHICH MAY OCCUR IN ANY PART OF THE BODY. ULCERS
VOL. III	NEURASTHENIA. TUMORS OF VARIOUS TISSUES OF THE BODY
VOL. IV	NEURASTHENIA. TUMORS OF VARIOUS TISSUES OF THE BODY
VOL. V	NEURASTHENIA. TUMORS OF VARIOUS TISSUES OF THE BODY
VOL. VI	NEURASTHENIA. TUMORS OF VARIOUS TISSUES OF THE BODY

VOLUMES I, II, III, IV, V, NOW READY; VOLUME VI, IN DECEMBER.


PRICES.

THE COMPLETE SET OF SIX VOLUMES	\$60.00
THE COMPLETE SET OF SIX VOLUMES	7.00
THE COMPLETE SET OF SIX VOLUMES	8.00

SEND FOR ILLUSTRATED ANNOUNCEMENT FREE.

Index.

[illegible]



A 001 429 930

11 1 000000

RM

216

F82m

